

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 5-(3-Fluorophenyl)-1-phenylpyrazolidin-3-one

Yuan-Yuan Liu, Hong Shi, Qing-Yan Chu and Hong-Jun Zhu\*

Department of Applied Chemistry, College of Science, Nanjing University of Technology, Nanjing 210009, People's Republic of China  
Correspondence e-mail: zhuhy@njut.edu.cn


Received 13 August 2008; accepted 19 August 2008

Key indicators: single-crystal X-ray study;  $T = 294$  K; mean  $\sigma(\text{C}-\text{C}) = 0.007$  Å;  $R$  factor = 0.082;  $wR$  factor = 0.233; data-to-parameter ratio = 14.4.

In the molecule of the title compound,  $\text{C}_{15}\text{H}_{13}\text{FN}_2\text{O}$ , the phenyl and fluorophenyl rings are oriented at a dihedral angle of  $77.92(3)^\circ$ . The pyrazolidine ring adopts an envelope conformation. An intramolecular  $\text{C}-\text{H}\cdots\text{N}$  hydrogen bond results in the formation of a five-membered ring adopting an envelope conformation. In the crystal structure, intermolecular  $\text{N}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds link the molecules. There are  $\text{C}-\text{H}\cdots\pi$  contacts between aromatic H atoms and the phenyl and fluorophenyl rings. A  $\pi-\pi$  contact between phenyl rings [centroid-centroid distance =  $3.926(1)$  Å] is also observed.

## Related literature

For general background, see: Chiara & Garcia (2005). For related literature, see: Jia *et al.* (2008). For bond-length data, see: Allen *et al.* (1987).



## Experimental

## Crystal data

$\text{C}_{15}\text{H}_{13}\text{FN}_2\text{O}$   
 $M_r = 256.27$   
Monoclinic,  $P2_1/c$   
 $a = 10.265(2)$  Å  
 $b = 7.3130(15)$  Å  
 $c = 17.822(4)$  Å  
 $\beta = 92.39(3)^\circ$   
 $V = 1336.7(5)$  Å<sup>3</sup>  
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.09$  mm<sup>-1</sup>  
 $T = 294(2)$  K  
 $0.30 \times 0.20 \times 0.10$  mm

## Data collection

Enraf-Nonius CAD-4 diffractometer  
Absorption correction:  $\psi$  scan (North *et al.*, 1968)  
 $T_{\min} = 0.973$ ,  $T_{\max} = 0.991$   
2393 measured reflections  
2393 independent reflections  
1231 reflections with  $I > 2\sigma(I)$   
3 standard reflections  
frequency: 120 min  
intensity decay: none

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.081$   
 $wR(F^2) = 0.232$   
 $S = 1.07$   
2393 reflections  
166 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.34$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.75$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2}-\text{H2A}\cdots\text{O}^i$	0.86	1.92	2.777 (4)	172
$\text{C4}-\text{H4A}\cdots\text{N1}$	0.93	2.48	2.836 (6)	103
$\text{C8}-\text{H8A}\cdots\text{O}^{ii}$	0.97	2.59	3.422 (5)	143
$\text{C6}-\text{H6A}\cdots\text{Cg3}^{iii}$	0.93	2.96	3.866 (3)	165
$\text{C12}-\text{H12A}\cdots\text{Cg2}^{iv}$	0.93	3.05	3.751 (3)	134
$\text{C14}-\text{H14A}\cdots\text{Cg3}^v$	0.93	2.80	3.654 (3)	153

Symmetry codes: (i)  $-x, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $-x, y + \frac{1}{2}, -z + \frac{1}{2}$ ; (iii)  $x, y + 1, z$ ; (iv)  $-x + 1, -y, -z + 1$ ; (v)  $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$ . Cg2 is the centroid of the C1-C6 ring and Cg3 is the centroid of the C10-C15 ring.

Data collection: *CAD-4 Software* (Enraf-Nonius, 1985); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the Center of Testing and Analysis, Nanjing University, for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2512).

## References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1-19.  
Chiara, J. L. & Garcia, A. (2005). *Synlett*, pp. 2607-2610.  
Enraf-Nonius (1985). *CAD-4 Software*. Enraf-Nonius, Delft, The Netherlands.  
Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.  
Jia, H.-S., Li, Y.-F., Liu, Y.-Y., Liu, S. & Zhu, H.-J. (2008). *Acta Cryst.* **E64**, o855.  
North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351-359.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112-122.

**supplementary materials**

*Acta Cryst.* (2008). E64, o1886 [ doi:10.1107/S1600536808026706 ]

## 5-(3-Fluorophenyl)-1-phenylpyrazolidin-3-one

Y.-Y. Liu, H. Shi, Q.-Y. Chu and H.-J. Zhu

### Comment

Pyrazolidin-3-one and its derivatives used as medicines and herbicides (Chiara & Garcia, 2005) have been developed most quickly, such as anodyne and antipyretic. We report herein the crystal structure of the title compound.

In the molecule of the title compound (Fig. 1), the bond lengths (Allen *et al.*, 1987) and angles are within normal ranges. Rings A (C1-C6) and C (C10-C15) are, of course, planar, and they are oriented at a dihedral angle of  $A/C = 77.92(3)^\circ$ . Ring B (N1/N2/C7-C9) adopts envelope conformation, with C8 atom displaced by  $0.277(3) \text{ \AA}$  from the plane of the other ring atoms. The intra-molecular C-H $\cdots$ N hydrogen bond (Table 1) results in the formation of a five-membered ring D (N1/C4/C5/C7/H4A) adopting envelope conformation, with N1 atom displaced by  $-0.326(3) \text{ \AA}$  from the plane of the other ring atoms.

In the crystal structure, intermolecular N-H $\cdots$ O and C-H $\cdots$ O hydrogen bonds link the molecules (Fig. 2), in which they may be effective in the stabilization of the structure. The C—H $\cdots$  $\pi$  contacts (Table 1) between the phenyl rings and the aromatic H atoms and a  $\pi$ — $\pi$  contact between phenyl rings  $Cg2 \cdots Cg2^i$  [symmetry code: (i)  $-x, 1-y, 1-z$ , where  $Cg2$  is centroid of the ring A (C1-C6)] further stabilize the structure, with centroid-centroid distance of  $3.926(1) \text{ \AA}$ .

### Experimental

The title compound was prepared according to the literature method (Jia *et al.*, 2008). The crystals were obtained by dissolving the title compound (1.5 g) in ethyl acetate (25 ml) and evaporating the solvent slowly at room temperature for about 10 d.

### Refinement

H atoms were positioned geometrically, with N-H =  $0.86 \text{ \AA}$  (for NH) and C-H =  $0.93, 0.98$  and  $0.97 \text{ \AA}$  for aromatic, methine and methylene H, respectively, and constrained to ride on their parent atoms with  $U_{iso}(H) = 1.2U_{eq}(C,N)$ .

### Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. Hydrogen bond is shown as dashed line.




Fig. 2. A partial packing diagram of the title compound. Hydrogen bonds are shown as dashed lines.

## 5-(3-fluorophenyl)-1-phenylpyrazolidin-3-one

### Crystal data

$C_{15}H_{13}FN_2O$

$M_r = 256.27$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 10.265$  (2) Å

$b = 7.3130$  (15) Å

$c = 17.822$  (4) Å

$\beta = 92.39$  (3)°

$V = 1336.7$  (5) Å<sup>3</sup>

$Z = 4$

$F_{000} = 536$

$D_x = 1.273$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation

$\lambda = 0.71073$  Å

Cell parameters from 25 reflections

$\theta = 9\text{--}12^\circ$

$\mu = 0.09$  mm<sup>-1</sup>

$T = 294$  (2) K

Needle, colorless

$0.30 \times 0.20 \times 0.10$  mm

### Data collection

Enraf-Nonius CAD-4  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 294$ (2) K

$\omega/2\theta$  scans

Absorption correction:  $\psi$  scan  
(North *et al.*, 1968)

$T_{\min} = 0.973$ ,  $T_{\max} = 0.991$

2393 measured reflections

2393 independent reflections

1231 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.0000$

$\theta_{\max} = 25.2^\circ$

$\theta_{\min} = 2.0^\circ$

$h = -12 \rightarrow 12$

$k = 0 \rightarrow 8$

$l = 0 \rightarrow 21$

3 standard reflections

every 120 min

intensity decay: none

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.081$

$wR(F^2) = 0.232$

$S = 1.07$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.1065P)^2 + 0.1962P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

2393 reflections  $\Delta\rho_{\max} = 0.34 \text{ e } \text{\AA}^{-3}$   
 166 parameters  $\Delta\rho_{\min} = -0.75 \text{ e } \text{\AA}^{-3}$   
 Primary atom site location: structure-invariant direct methods Extinction correction: none

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor wR and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O	0.0023 (3)	0.2565 (4)	0.22989 (17)	0.0567 (8)
F	0.2688 (3)	0.7468 (5)	0.56248 (19)	0.098
N1	0.2364 (3)	0.0858 (4)	0.35618 (19)	0.0407 (8)
C1	0.2186 (8)	0.5785 (10)	0.5378 (4)	0.104 (2)
N2	0.1332 (3)	0.0777 (4)	0.3021 (2)	0.0477 (9)
H2A	0.0965	-0.0235	0.2886	0.057*
C2	0.1419 (7)	0.4819 (12)	0.5800 (4)	0.095 (2)
H2B	0.1174	0.5252	0.6264	0.114*
C3	0.0997 (6)	0.3176 (11)	0.5537 (3)	0.0873 (18)
H3A	0.0426	0.2492	0.5816	0.105*
C4	0.1399 (5)	0.2505 (7)	0.4868 (3)	0.0650 (13)
H4A	0.1112	0.1361	0.4704	0.078*
C5	0.2215 (4)	0.3492 (6)	0.4436 (2)	0.0488 (11)
C6	0.2666 (6)	0.5199 (7)	0.4691 (3)	0.0790 (16)
H6A	0.3245	0.5898	0.4424	0.095*
C7	0.2639 (4)	0.2853 (5)	0.3683 (2)	0.0421 (10)
H7A	0.3576	0.3073	0.3647	0.051*
C8	0.1895 (4)	0.3791 (5)	0.3012 (2)	0.0452 (10)
H8A	0.1435	0.4870	0.3174	0.054*
H8B	0.2487	0.4137	0.2626	0.054*
C9	0.0967 (4)	0.2358 (5)	0.2735 (2)	0.0433 (10)
C10	0.3413 (3)	-0.0333 (5)	0.3443 (2)	0.0385 (9)
C11	0.4505 (4)	-0.0252 (6)	0.3943 (2)	0.0509 (11)
H11A	0.4529	0.0611	0.4327	0.061*
C12	0.5555 (4)	-0.1441 (7)	0.3875 (3)	0.0592 (13)
H12A	0.6282	-0.1359	0.4203	0.071*
C13	0.5499 (5)	-0.2750 (6)	0.3310 (3)	0.0660 (14)
H13A	0.6185	-0.3569	0.3265	0.079*

## supplementary materials

---

C14	0.4453 (4)	-0.2840 (6)	0.2825 (3)	0.0573 (12)
H14A	0.4430	-0.3718	0.2447	0.069*
C15	0.3405 (4)	-0.1633 (5)	0.2880 (2)	0.0423 (10)
H15A	0.2699	-0.1706	0.2537	0.051*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O	0.0586 (18)	0.0318 (16)	0.079 (2)	0.0078 (14)	-0.0066 (17)	0.0028 (15)
F	0.098	0.098	0.098	0.000	0.004	0.000
N1	0.0375 (18)	0.0253 (17)	0.059 (2)	0.0030 (14)	-0.0036 (16)	-0.0029 (15)
C1	0.125 (6)	0.083 (5)	0.101 (5)	0.018 (4)	-0.022 (5)	-0.058 (4)
N2	0.0402 (19)	0.0242 (17)	0.078 (3)	0.0014 (15)	-0.0117 (18)	0.0065 (16)
C2	0.089 (5)	0.132 (7)	0.063 (4)	0.044 (5)	0.007 (3)	-0.022 (4)
C3	0.081 (4)	0.117 (6)	0.064 (4)	0.015 (4)	0.002 (3)	0.008 (4)
C4	0.055 (3)	0.067 (3)	0.072 (3)	0.008 (3)	-0.002 (3)	0.000 (3)
C5	0.043 (2)	0.045 (3)	0.058 (3)	0.010 (2)	-0.004 (2)	-0.005 (2)
C6	0.109 (4)	0.063 (3)	0.065 (3)	-0.010 (3)	0.006 (3)	-0.027 (3)
C7	0.040 (2)	0.0212 (19)	0.066 (3)	-0.0020 (17)	0.0037 (19)	-0.0048 (19)
C8	0.040 (2)	0.030 (2)	0.066 (3)	-0.0010 (18)	0.007 (2)	-0.003 (2)
C9	0.042 (2)	0.031 (2)	0.057 (2)	0.0024 (19)	0.005 (2)	-0.013 (2)
C10	0.032 (2)	0.031 (2)	0.052 (2)	0.0004 (17)	-0.0015 (18)	0.0085 (19)
C11	0.047 (2)	0.049 (3)	0.056 (3)	0.014 (2)	-0.004 (2)	-0.009 (2)
C12	0.051 (3)	0.063 (3)	0.062 (3)	0.006 (2)	-0.017 (2)	0.009 (3)
C13	0.052 (3)	0.043 (3)	0.105 (4)	0.012 (2)	0.023 (3)	0.001 (3)
C14	0.055 (3)	0.042 (3)	0.076 (3)	0.006 (2)	0.024 (2)	-0.017 (2)
C15	0.038 (2)	0.039 (2)	0.050 (2)	-0.0045 (19)	0.0006 (18)	-0.0047 (19)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

O—C9	1.226 (5)	C6—H6A	0.9300
F—C1	1.398 (7)	C7—C8	1.552 (6)
N1—N2	1.404 (4)	C7—H7A	0.9800
N1—C10	1.408 (5)	C8—C9	1.487 (5)
N1—C7	1.500 (4)	C8—H8A	0.9700
C1—C2	1.317 (9)	C8—H8B	0.9700
C1—C6	1.405 (9)	C10—C15	1.381 (5)
N2—C9	1.311 (5)	C10—C11	1.404 (5)
N2—H2A	0.8600	C11—C12	1.394 (6)
C2—C3	1.354 (9)	C11—H11A	0.9300
C2—H2B	0.9300	C12—C13	1.389 (6)
C3—C4	1.370 (8)	C12—H12A	0.9300
C3—H3A	0.9300	C13—C14	1.353 (6)
C4—C5	1.366 (7)	C13—H13A	0.9300
C4—H4A	0.9300	C14—C15	1.398 (6)
C5—C6	1.400 (6)	C14—H14A	0.9300
C5—C7	1.502 (6)	C15—H15A	0.9300
N2—N1—C10	115.5 (3)	C8—C7—H7A	109.2

N2—N1—C7	105.8 (3)	C9—C8—C7	103.4 (3)
C10—N1—C7	118.9 (3)	C9—C8—H8A	111.1
C2—C1—F	120.9 (7)	C7—C8—H8A	111.1
C2—C1—C6	125.0 (6)	C9—C8—H8B	111.1
F—C1—C6	113.9 (7)	C7—C8—H8B	111.1
C9—N2—N1	115.1 (3)	H8A—C8—H8B	109.0
C9—N2—H2A	122.4	O—C9—N2	124.1 (4)
N1—N2—H2A	122.4	O—C9—C8	127.0 (4)
C1—C2—C3	117.9 (6)	N2—C9—C8	108.9 (3)
C1—C2—H2B	121.0	C15—C10—C11	118.0 (4)
C3—C2—H2B	121.0	C15—C10—N1	123.6 (3)
C2—C3—C4	121.1 (6)	C11—C10—N1	118.3 (3)
C2—C3—H3A	119.5	C12—C11—C10	121.2 (4)
C4—C3—H3A	119.5	C12—C11—H11A	119.4
C5—C4—C3	121.0 (5)	C10—C11—H11A	119.4
C5—C4—H4A	119.5	C13—C12—C11	119.0 (4)
C3—C4—H4A	119.5	C13—C12—H12A	120.5
C4—C5—C6	119.4 (4)	C11—C12—H12A	120.5
C4—C5—C7	123.0 (4)	C14—C13—C12	120.3 (4)
C6—C5—C7	117.6 (4)	C14—C13—H13A	119.8
C5—C6—C1	115.6 (6)	C12—C13—H13A	119.8
C5—C6—H6A	122.2	C13—C14—C15	121.1 (4)
C1—C6—H6A	122.2	C13—C14—H14A	119.4
N1—C7—C5	111.8 (3)	C15—C14—H14A	119.4
N1—C7—C8	103.6 (3)	C10—C15—C14	120.3 (4)
C5—C7—C8	113.6 (3)	C10—C15—H15A	119.8
N1—C7—H7A	109.2	C14—C15—H15A	119.8
C5—C7—H7A	109.2		
C10—N1—N2—C9	-128.7 (4)	C6—C5—C7—C8	-76.9 (5)
C7—N1—N2—C9	5.0 (5)	N1—C7—C8—C9	17.4 (4)
F—C1—C2—C3	-178.1 (5)	C5—C7—C8—C9	-104.2 (4)
C6—C1—C2—C3	-3.9 (11)	N1—N2—C9—O	-174.3 (4)
C1—C2—C3—C4	2.6 (9)	N1—N2—C9—C8	7.0 (5)
C2—C3—C4—C5	-1.4 (8)	C7—C8—C9—O	166.0 (4)
C3—C4—C5—C6	1.2 (7)	C7—C8—C9—N2	-15.3 (4)
C3—C4—C5—C7	-177.3 (4)	N2—N1—C10—C15	-6.1 (5)
C4—C5—C6—C1	-2.1 (7)	C7—N1—C10—C15	-133.5 (4)
C7—C5—C6—C1	176.5 (5)	N2—N1—C10—C11	176.8 (3)
C2—C1—C6—C5	3.6 (10)	C7—N1—C10—C11	49.4 (5)
F—C1—C6—C5	178.2 (5)	C15—C10—C11—C12	0.2 (6)
N2—N1—C7—C5	108.7 (3)	N1—C10—C11—C12	177.4 (4)
C10—N1—C7—C5	-119.4 (4)	C10—C11—C12—C13	-1.3 (7)
N2—N1—C7—C8	-13.9 (4)	C11—C12—C13—C14	1.4 (7)
C10—N1—C7—C8	117.9 (4)	C12—C13—C14—C15	-0.4 (7)
C4—C5—C7—N1	-15.1 (5)	C11—C10—C15—C14	0.8 (6)
C6—C5—C7—N1	166.3 (4)	N1—C10—C15—C14	-176.2 (4)
C4—C5—C7—C8	101.7 (4)	C13—C14—C15—C10	-0.8 (7)

## supplementary materials

---

### Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N2—H2A···O <sup>i</sup>	0.86	1.92	2.777 (4)	172
C4—H4A···N1	0.93	2.48	2.836 (6)	103
C8—H8A···O <sup>ii</sup>	0.97	2.59	3.422 (5)	143
C6—H6A···Cg3 <sup>iii</sup>	0.93	2.96	3.866 (3)	165
C12—H12A···Cg2 <sup>iv</sup>	0.93	3.05	3.751 (3)	134
C14—H14A···Cg3 <sup>v</sup>	0.93	2.80	3.654 (3)	153

Symmetry codes: (i)  $-x, y-1/2, -z+1/2$ ; (ii)  $-x, y+1/2, -z+1/2$ ; (iii)  $x, y+1, z$ ; (iv)  $-x+1, -y, -z+1$ ; (v)  $-x+1, y-1/2, -z+1/2$ .

Fig. 1




Fig. 2

