

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bis(diethylenetriamine- κ^3N,N',N'')-nickel(II) bis(1,2-dicyanoethene-1,2-dithiolato- κ^2S,S')nickel(II)

Dao-Peng Zhang, Hai-Long Wang, Li-Fang Zhang* and Zhong-Hai Ni

School of Chemistry & Chemical Technology, Shandong University, Jinan 250100, People's Republic of China

Correspondence e-mail: Zhanglf@sdu.edu.cn


Received 20 August 2008; accepted 8 September 2008

Key indicators: single-crystal X-ray study; $T = 273$ K; mean $\sigma(C-C) = 0.006$ Å; R factor = 0.041; wR factor = 0.109; data-to-parameter ratio = 15.4.

The title compound, $[\text{Ni}(\text{C}_4\text{H}_{13}\text{N}_3)_2][\text{Ni}(\text{C}_4\text{N}_2\text{S}_2)_2]$, has been synthesized by the reaction of $\text{Ni}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$, diethylenetriamine (deta) and $\text{Na}_2[\text{Ni}(\text{mnt})_2]$ [$\text{mnt} = \text{maleonitriledithiolate}(2-)$] in methanol. The structure is composed of a $[\text{Ni}(\text{deta})_2]^{2+}$ cation and a $[\text{Ni}(\text{mnt})_2]^{2-}$ anion. The coordination geometry of the Ni^{II} ion in the cation is slightly distorted octahedral, defined by six N atoms from two deta ligands, while the Ni^{II} ion in the anion is four-coordinated by four S atoms from two mnt ligands in a slightly distorted square-planar geometry. The cations and anions are connected by $\text{N} \cdots \text{H} \cdots \text{N}$ hydrogen bonds.

Related literature

For related literature, see: Bois *et al.* (1998); Keum *et al.* (1992); Miller *et al.* (1989); Ren *et al.* (2001); Robertson & Cronin (2002); Simmons *et al.* (1962).



Experimental

Crystal data

$[\text{Ni}(\text{C}_4\text{H}_{13}\text{N}_3)_2][\text{Ni}(\text{C}_4\text{N}_2\text{S}_2)_2]$
 $M_r = 604.13$
 Monoclinic, $P2_1/n$
 $a = 9.589$ (3) Å
 $b = 16.910$ (5) Å
 $c = 16.146$ (4) Å
 $\beta = 97.491$ (4)°

$V = 2595.8$ (13) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.80$ mm⁻¹
 $T = 273$ (2) K
 $0.19 \times 0.17 \times 0.15$ mm

Data collection

Bruker SMART APEXII CCD
 area-detector diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.717$, $T_{\text{max}} = 0.766$

13610 measured reflections
 5065 independent reflections
 3393 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.109$
 $S = 0.99$
 5065 reflections
 329 parameters
 10 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 0.86$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.26$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Ni1—S1	2.1739 (12)	Ni2—N6	2.065 (3)
Ni1—S2	2.1617 (12)	Ni2—N7	2.150 (4)
Ni1—S3	2.1732 (12)	Ni2—N8	2.145 (3)
Ni1—S4	2.1658 (12)	Ni2—N9	2.071 (4)
Ni2—N5	2.164 (3)	Ni2—N10	2.151 (3)
S2—Ni1—S4	87.98 (5)	N8—Ni2—N7	95.96 (15)
S2—Ni1—S3	168.77 (5)	N6—Ni2—N10	98.69 (14)
S4—Ni1—S3	92.72 (4)	N9—Ni2—N10	81.72 (14)
S2—Ni1—S1	92.58 (4)	N8—Ni2—N10	163.04 (16)
S4—Ni1—S1	170.10 (4)	N7—Ni2—N10	89.25 (16)
S3—Ni1—S1	88.65 (4)	N6—Ni2—N5	81.47 (14)
N6—Ni2—N9	177.56 (15)	N9—Ni2—N5	100.96 (15)
N6—Ni2—N8	98.00 (14)	N8—Ni2—N5	91.32 (13)
N9—Ni2—N8	81.73 (14)	N7—Ni2—N5	162.93 (16)
N6—Ni2—N7	82.24 (16)	N10—Ni2—N5	88.20 (14)
N9—Ni2—N7	95.37 (17)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
N5—H5A \cdots N4 ⁱ	0.86 (3)	2.30 (4)	3.098 (6)	154 (3)
N5—H5B \cdots N2 ⁱⁱ	0.86 (3)	2.48 (3)	3.186 (5)	140 (4)
N7—H7A \cdots N3 ⁱⁱⁱ	0.86 (4)	2.56 (3)	3.207 (7)	134 (3)
N8—H8B \cdots N3 ⁱⁱⁱ	0.86 (3)	2.48 (4)	3.164 (6)	138 (3)
N9—H9A \cdots N1 ^{iv}	0.86 (2)	2.58 (3)	3.387 (6)	156 (5)
N10—H10C \cdots N2 ⁱⁱ	0.87 (3)	2.34 (3)	3.198 (5)	173 (5)

Symmetry codes: (i) $-x + 1, -y + 1, -z$; (ii) $x - \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (iii) $x + 1, y, z$; (iv) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT-Plus (Bruker, 2007); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This work was supported by the Natural Science Foundation of China (No. 20701027) and the Postdoctoral Scientific Foundation of China (No. 200704211076).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2151).

References

- Bois, H., Connelly, N. G., Crossley, J. G., Guillorit, J., Lewis, G. R., Orpen, A. G. & Thornton, P. (1998). *J. Chem. Soc. Dalton Trans.* pp. 2833–2838.
- Bruker (2007). *APEX2* and *SAINT-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Keum, C., Kim, C., Kim, C., Kwak, H., Kwon, M. & Namgung, H. (1992). *Bull. Korean. Chem. Soc.* **13**, 695–699.
- Miller, J. S., Calabrese, J. C. & Epstein, A. J. (1989). *Inorg. Chem.* **28**, 4230–4238.
- Ren, X.-M., Duan, C.-Y., Zhu, H.-Z., Meng, Q.-J., Hu, C.-J., Lu, C.-S. & Liu, Y.-J. (2001). *Transition Met. Chem.* **26**, 295–299.
- Robertson, N. & Cronin, L. (2002). *Coord. Chem. Rev.* **227**, 93–127.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Simmons, H. E., Vest, R. D., Blomstrom, D. C., Roland, J. R. & Cairns, T. L. (1962). *J. Am. Chem. Soc.* **84**, 4746–4756.

supplementary materials

Acta Cryst. (2008). E64, m1264-m1265 [doi:10.1107/S1600536808028663]

Bis(diethylenetriamine- κ^3N,N',N'')nickel(II) bis(1,2-dicyanoethene-1,2-dithiolato- κ^2S,S')nickel(II)

D.-P. Zhang, H.-L. Wang, L.-F. Zhang and Z.-H. Ni

Comment

Bidentate dithiolate ligands form well square-planar complexes with nickel ions in different oxidation states. Due to their unique properties and potential applications in such areas as conducting and magnetic materials, nearinfrared dyes, nonlinear optical materials (Robertson & Cronin, 2002), the ion-pair complexes formed from $[M(\text{mnt})_2]^{n-}$ ($M = \text{Ni}, \text{Pd}, \text{Pt}$ or Cu) and transition metal complex cations have been intensively studied (Bois *et al.*, 1998; Miller *et al.*, 1989; Ren *et al.*, 2001). We report here a new ion-pair complex.

The title compound is composed of a $[\text{Ni}(\text{deta})_2]^{2+}$ cation and a $[\text{Ni}(\text{mnt})_2]^{2-}$ anion [deta = diethylenetriamine; mnt = maleonitriledithiolate(2-)] (Fig. 1). In the cation, the Ni^{II} ion has a slightly distorted octahedral geometry, formed by six N atoms from two deta ligands, with the Ni—N distances in a range from 2.065 (3) to 2.164 (3) Å (Table 1), which are consistent with the corresponding values in $[\text{Ni}(\text{en})_3][\text{Ni}(\text{mnt})_2]$ (en = ethylenediamine) (Keum *et al.*, 1992). The Ni^{II} ion in the anion is four-coordinated by four S atoms and these five atoms form a square plane with a mean deviation of 0.161 (6) Å. The Ni—S bond lengths [2.1617 (12)–2.1739 (12) Å] are also in agreement with those found in the above complex. The cations and anions are connected by N—H \cdots N hydrogen bonds (Table 2).

Experimental

The synthesis procedure of the title compound was as following: $\text{Ni}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ (0.037 g, 0.10 mmol) was dissolved in methanol (10 ml) at room temperature with stirring and then deta (0.021 g, 0.20 mmol) was added. A solution of $\text{Na}_2[\text{Ni}(\text{mnt})_2]$ (0.033 g, 0.10 mmol) (Simmons *et al.*, 1962) in methanol (10 ml) was slowly added to the above solution and the mixture was stirred for another 30 min. After filtering, the filtrate was undisturbed for about two weeks at room temperature in air to produce blue crystals suitable for X-ray diffraction (yield 61.75%, 0.037 g). Analysis, calculated for $\text{C}_{16}\text{H}_{26}\text{N}_{10}\text{Ni}_2\text{S}_4$: C 31.81, H 4.34, N 23.19%; found: C 31.76, H 4.31, N 23.24%.

Refinement

H atoms bound to N atoms were found in difference Fourier maps and refined isotropically, with a restraint of N—H = 0.86 (1) Å. H atoms bound to C atoms were positioned geometrically and refined as riding atoms, with C—H = 0.97 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. Structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level. H atoms have been omitted for clarity.

Bis(diethylenetriamine- $\kappa^3\text{N},\text{N}',\text{N}''$)nickel(II) bis(1,2-dicyanoethene-1,2-dithiolato- $\kappa^2\text{S},\text{S}'$)nickel(II)

Crystal data

$[\text{Ni}(\text{C}_4\text{H}_{13}\text{N}_3)_2][\text{Ni}(\text{C}_4\text{N}_2\text{S}_2)_2]$

$M_r = 604.13$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 9.589$ (3) Å

$b = 16.910$ (5) Å

$c = 16.146$ (4) Å

$\beta = 97.491$ (4)°

$V = 2595.8$ (13) Å³

$Z = 4$

$F_{000} = 1248$

$D_x = 1.546$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 3163 reflections

$\theta = 2.4$ – 23.8 °

$\mu = 1.80$ mm⁻¹

$T = 273$ (2) K

Block, blue

$0.19 \times 0.17 \times 0.15$ mm

Data collection

Bruker SMART APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 273$ (2) K

φ and ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.717$, $T_{\max} = 0.766$

13610 measured reflections

5065 independent reflections

3393 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.037$

$\theta_{\max} = 26.0$ °

$\theta_{\min} = 2.3$ °

$h = -11 \rightarrow 11$

$k = -20 \rightarrow 19$

$l = -17 \rightarrow 19$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.041$

$wR(F^2) = 0.109$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.058P)^2]$

$S = 0.99$

5065 reflections

329 parameters

10 restraints

Primary atom site location: structure-invariant direct methods

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.86 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.26 \text{ e } \text{\AA}^{-3}$

Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.72467 (5)	0.23459 (3)	0.01993 (3)	0.04722 (16)
Ni2	0.97276 (5)	0.40148 (3)	0.25321 (3)	0.04194 (15)
S1	0.78132 (10)	0.13475 (6)	0.10229 (6)	0.0555 (3)
S2	0.90403 (11)	0.21706 (7)	-0.04639 (7)	0.0638 (3)
S3	0.52234 (11)	0.23887 (7)	0.06673 (7)	0.0598 (3)
S4	0.68871 (10)	0.34610 (6)	-0.04535 (7)	0.0576 (3)
N1	1.0808 (4)	0.0021 (3)	0.1686 (3)	0.0828 (12)
N2	1.2281 (4)	0.0992 (2)	-0.0397 (2)	0.0727 (11)
N3	0.1991 (4)	0.3603 (3)	0.0519 (3)	0.0980 (15)
N4	0.3914 (5)	0.4842 (3)	-0.1113 (3)	0.0902 (13)
N5	0.7588 (4)	0.4323 (2)	0.2690 (2)	0.0532 (8)
N6	0.8844 (4)	0.2901 (2)	0.2388 (2)	0.0573 (9)
N7	1.1630 (4)	0.3364 (3)	0.2450 (3)	0.0670 (10)
N8	0.9396 (4)	0.4370 (2)	0.1246 (2)	0.0551 (9)
N9	1.0692 (4)	0.5111 (2)	0.2663 (2)	0.0576 (9)
N10	1.0222 (4)	0.4021 (3)	0.3870 (2)	0.0552 (9)
C1	0.9425 (4)	0.1059 (2)	0.0742 (2)	0.0487 (9)
C2	0.9942 (4)	0.1410 (2)	0.0098 (2)	0.0498 (9)
C3	0.4468 (4)	0.3230 (2)	0.0193 (2)	0.0546 (10)
C4	0.5168 (4)	0.3686 (2)	-0.0311 (2)	0.0532 (10)
C5	1.0189 (4)	0.0466 (3)	0.1249 (3)	0.0587 (11)
C6	1.1252 (4)	0.1169 (3)	-0.0162 (2)	0.0564 (11)
C7	0.3083 (5)	0.3441 (3)	0.0363 (3)	0.0676 (12)
C8	0.4494 (5)	0.4331 (3)	-0.0751 (3)	0.0656 (12)
C9	0.6713 (4)	0.3606 (3)	0.2543 (3)	0.0690 (12)
H9C	0.6357	0.3564	0.1954	0.083*
H9B	0.5916	0.3639	0.2854	0.083*
C10	0.7577 (5)	0.2887 (3)	0.2815 (3)	0.0711 (13)
H10A	0.7841	0.2894	0.3416	0.085*
H10B	0.7039	0.2410	0.2668	0.085*
C11	0.9946 (5)	0.2337 (3)	0.2698 (3)	0.0759 (14)
H11A	0.9634	0.1802	0.2563	0.091*
H11B	1.0161	0.2381	0.3301	0.091*
C12	1.1225 (6)	0.2523 (3)	0.2293 (3)	0.0850 (16)
H12A	1.1994	0.2181	0.2518	0.102*
H12B	1.1030	0.2429	0.1696	0.102*
C13	1.0004 (5)	0.5157 (3)	0.1168 (3)	0.0748 (13)

supplementary materials

H13A	0.9436	0.5450	0.0730	0.090*
H13B	1.0944	0.5106	0.1013	0.090*
C14	1.0069 (5)	0.5602 (3)	0.1974 (3)	0.0748 (13)
H14A	1.0627	0.6078	0.1945	0.090*
H14B	0.9129	0.5758	0.2066	0.090*
C15	1.0661 (5)	0.5377 (3)	0.3519 (3)	0.0694 (12)
H15A	0.9721	0.5556	0.3585	0.083*
H15B	1.1303	0.5818	0.3639	0.083*
C16	1.1076 (5)	0.4719 (3)	0.4112 (2)	0.0704 (13)
H16A	1.2063	0.4595	0.4109	0.084*
H16B	1.0943	0.4879	0.4674	0.084*
H8A	0.8544 (16)	0.441 (2)	0.101 (2)	0.051 (11)*
H8B	0.971 (4)	0.4005 (18)	0.095 (2)	0.073 (15)*
H7A	1.219 (4)	0.352 (3)	0.211 (2)	0.090 (17)*
H10C	0.944 (3)	0.406 (3)	0.408 (3)	0.091 (17)*
H5A	0.727 (5)	0.4692 (19)	0.235 (2)	0.093 (18)*
H6A	0.860 (4)	0.281 (2)	0.1865 (8)	0.055 (12)*
H7B	1.217 (4)	0.341 (3)	0.2916 (17)	0.12 (2)*
H10D	1.066 (6)	0.362 (3)	0.409 (4)	0.16 (3)*
H9A	1.1581 (18)	0.503 (4)	0.267 (4)	0.16 (3)*
H5B	0.769 (5)	0.448 (3)	0.3200 (12)	0.101 (18)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.0377 (3)	0.0536 (3)	0.0497 (3)	0.0057 (2)	0.0033 (2)	-0.0112 (2)
Ni2	0.0429 (3)	0.0502 (3)	0.0324 (2)	-0.0003 (2)	0.00373 (19)	-0.0015 (2)
S1	0.0412 (5)	0.0638 (7)	0.0642 (6)	0.0081 (5)	0.0164 (5)	-0.0001 (5)
S2	0.0514 (6)	0.0822 (8)	0.0599 (6)	0.0189 (6)	0.0160 (5)	0.0104 (6)
S3	0.0503 (6)	0.0652 (7)	0.0663 (7)	0.0126 (5)	0.0161 (5)	-0.0012 (5)
S4	0.0451 (6)	0.0601 (7)	0.0667 (7)	0.0028 (5)	0.0040 (5)	-0.0026 (5)
N1	0.062 (2)	0.098 (3)	0.093 (3)	0.027 (2)	0.024 (2)	0.027 (2)
N2	0.055 (2)	0.110 (3)	0.055 (2)	0.029 (2)	0.0172 (18)	0.016 (2)
N3	0.078 (3)	0.134 (4)	0.089 (3)	0.048 (3)	0.038 (2)	0.009 (3)
N4	0.097 (3)	0.094 (3)	0.077 (3)	0.032 (3)	0.000 (2)	0.008 (2)
N5	0.048 (2)	0.067 (2)	0.044 (2)	0.0015 (18)	0.0033 (16)	-0.0034 (19)
N6	0.067 (2)	0.056 (2)	0.047 (2)	-0.0026 (18)	-0.0011 (18)	-0.0019 (17)
N7	0.055 (2)	0.087 (3)	0.059 (2)	0.014 (2)	0.008 (2)	-0.003 (2)
N8	0.048 (2)	0.077 (3)	0.0393 (19)	0.014 (2)	0.0049 (16)	0.0003 (18)
N9	0.065 (2)	0.059 (2)	0.051 (2)	-0.0045 (19)	0.0138 (17)	-0.0004 (17)
N10	0.042 (2)	0.087 (3)	0.0367 (17)	-0.009 (2)	0.0043 (15)	-0.0018 (18)
C1	0.037 (2)	0.058 (3)	0.051 (2)	0.0068 (18)	0.0077 (17)	-0.0086 (19)
C2	0.037 (2)	0.064 (3)	0.049 (2)	0.0069 (19)	0.0053 (17)	-0.0067 (19)
C3	0.048 (2)	0.068 (3)	0.047 (2)	0.017 (2)	0.0024 (18)	-0.016 (2)
C4	0.050 (2)	0.058 (3)	0.049 (2)	0.013 (2)	-0.0017 (18)	-0.013 (2)
C5	0.043 (2)	0.073 (3)	0.063 (3)	0.007 (2)	0.020 (2)	0.002 (2)
C6	0.051 (2)	0.076 (3)	0.043 (2)	0.016 (2)	0.0078 (18)	0.006 (2)
C7	0.064 (3)	0.087 (3)	0.053 (3)	0.025 (3)	0.014 (2)	-0.003 (2)

C8	0.063 (3)	0.077 (3)	0.054 (3)	0.018 (2)	-0.003 (2)	-0.005 (2)
C9	0.048 (2)	0.092 (4)	0.067 (3)	-0.018 (3)	0.006 (2)	-0.014 (3)
C10	0.073 (3)	0.070 (3)	0.071 (3)	-0.026 (3)	0.014 (2)	-0.002 (2)
C11	0.093 (4)	0.057 (3)	0.073 (3)	0.009 (3)	-0.009 (3)	0.008 (2)
C12	0.096 (4)	0.072 (4)	0.082 (3)	0.032 (3)	-0.005 (3)	-0.009 (3)
C13	0.090 (3)	0.081 (4)	0.056 (3)	0.004 (3)	0.021 (2)	0.023 (2)
C14	0.087 (4)	0.052 (3)	0.085 (3)	-0.001 (2)	0.012 (3)	0.008 (2)
C15	0.070 (3)	0.067 (3)	0.071 (3)	-0.005 (2)	0.010 (2)	-0.028 (2)
C16	0.066 (3)	0.100 (4)	0.045 (2)	-0.025 (3)	0.006 (2)	-0.013 (2)

Geometric parameters (Å, °)

Ni1—S1	2.1739 (12)	N9—C15	1.456 (5)
Ni1—S2	2.1617 (12)	N9—H9A	0.86 (3)
Ni1—S3	2.1732 (12)	N10—C16	1.461 (5)
Ni1—S4	2.1658 (12)	N10—H10C	0.87 (3)
Ni2—N5	2.164 (3)	N10—H10D	0.85 (6)
Ni2—N6	2.065 (3)	C1—C2	1.347 (5)
Ni2—N7	2.150 (4)	C1—C5	1.434 (6)
Ni2—N8	2.145 (3)	C2—C6	1.434 (5)
Ni2—N9	2.071 (4)	C3—C4	1.359 (6)
Ni2—N10	2.151 (3)	C3—C7	1.436 (5)
S1—C1	1.737 (4)	C4—C8	1.412 (6)
S2—C2	1.738 (4)	C9—C10	1.505 (6)
S3—C3	1.730 (4)	C9—H9C	0.9700
S4—C4	1.737 (4)	C9—H9B	0.9700
N1—C5	1.144 (5)	C10—H10A	0.9700
N2—C6	1.143 (5)	C10—H10B	0.9700
N3—C7	1.141 (5)	C11—C12	1.497 (7)
N4—C8	1.146 (5)	C11—H11A	0.9700
N5—C9	1.477 (5)	C11—H11B	0.9700
N5—H5A	0.86 (3)	C12—H12A	0.9700
N5—H5B	0.86 (3)	C12—H12B	0.9700
N6—C11	1.463 (5)	C13—C14	1.497 (6)
N6—C10	1.473 (5)	C13—H13A	0.9700
N6—H6A	0.86 (2)	C13—H13B	0.9700
N7—C12	1.486 (6)	C14—H14A	0.9700
N7—H7A	0.86 (4)	C14—H14B	0.9700
N7—H7B	0.86 (4)	C15—C16	1.488 (6)
N8—C13	1.466 (6)	C15—H15A	0.9700
N8—H8A	0.86 (3)	C15—H15B	0.9700
N8—H8B	0.86 (3)	C16—H16A	0.9700
N9—C14	1.453 (5)	C16—H16B	0.9700
S2—Ni1—S4	87.98 (5)	C5—C1—S1	117.0 (3)
S2—Ni1—S3	168.77 (5)	C1—C2—C6	121.7 (4)
S4—Ni1—S3	92.72 (4)	C1—C2—S2	121.4 (3)
S2—Ni1—S1	92.58 (4)	C6—C2—S2	116.9 (3)
S4—Ni1—S1	170.10 (4)	C4—C3—C7	121.0 (4)
S3—Ni1—S1	88.65 (4)	C4—C3—S3	121.4 (3)

supplementary materials

N6—Ni2—N9	177.56 (15)	C7—C3—S3	117.7 (3)
N6—Ni2—N8	98.00 (14)	C3—C4—C8	120.7 (4)
N9—Ni2—N8	81.73 (14)	C3—C4—S4	120.5 (3)
N6—Ni2—N7	82.24 (16)	C8—C4—S4	118.8 (3)
N9—Ni2—N7	95.37 (17)	N1—C5—C1	176.5 (5)
N8—Ni2—N7	95.96 (15)	N2—C6—C2	177.4 (4)
N6—Ni2—N10	98.69 (14)	N3—C7—C3	178.2 (5)
N9—Ni2—N10	81.72 (14)	N4—C8—C4	178.1 (5)
N8—Ni2—N10	163.04 (16)	N5—C9—C10	109.7 (3)
N7—Ni2—N10	89.25 (16)	N5—C9—H9C	109.7
N6—Ni2—N5	81.47 (14)	C10—C9—H9C	109.7
N9—Ni2—N5	100.96 (15)	N5—C9—H9B	109.7
N8—Ni2—N5	91.32 (13)	C10—C9—H9B	109.7
N7—Ni2—N5	162.93 (16)	H9C—C9—H9B	108.2
N10—Ni2—N5	88.20 (14)	N6—C10—C9	107.8 (3)
C1—S1—Ni1	102.61 (14)	N6—C10—H10A	110.1
C2—S2—Ni1	102.53 (13)	C9—C10—H10A	110.1
C3—S3—Ni1	102.40 (15)	N6—C10—H10B	110.1
C4—S4—Ni1	102.68 (15)	C9—C10—H10B	110.1
C9—N5—Ni2	108.1 (3)	H10A—C10—H10B	108.5
C9—N5—H5A	110 (3)	N6—C11—C12	108.0 (4)
Ni2—N5—H5A	111 (3)	N6—C11—H11A	110.1
C9—N5—H5B	114 (3)	C12—C11—H11A	110.1
Ni2—N5—H5B	101 (3)	N6—C11—H11B	110.1
H5A—N5—H5B	112 (5)	C12—C11—H11B	110.1
C11—N6—C10	115.5 (4)	H11A—C11—H11B	108.4
C11—N6—Ni2	106.8 (3)	N7—C12—C11	109.7 (4)
C10—N6—Ni2	108.3 (3)	N7—C12—H12A	109.7
C11—N6—H6A	108 (3)	C11—C12—H12A	109.7
C10—N6—H6A	109 (3)	N7—C12—H12B	109.7
Ni2—N6—H6A	109 (3)	C11—C12—H12B	109.7
C12—N7—Ni2	107.2 (3)	H12A—C12—H12B	108.2
C12—N7—H7A	110 (3)	N8—C13—C14	110.7 (3)
Ni2—N7—H7A	120 (3)	N8—C13—H13A	109.5
C12—N7—H7B	110 (4)	C14—C13—H13A	109.5
Ni2—N7—H7B	109 (4)	N8—C13—H13B	109.5
H7A—N7—H7B	100 (5)	C14—C13—H13B	109.5
C13—N8—Ni2	109.1 (3)	H13A—C13—H13B	108.1
C13—N8—H8A	105 (3)	N9—C14—C13	110.2 (4)
Ni2—N8—H8A	118 (2)	N9—C14—H14A	109.6
C13—N8—H8B	116 (3)	C13—C14—H14A	109.6
Ni2—N8—H8B	108 (3)	N9—C14—H14B	109.6
H8A—N8—H8B	101 (4)	C13—C14—H14B	109.6
C14—N9—C15	119.5 (4)	H14A—C14—H14B	108.1
C14—N9—Ni2	107.4 (3)	N9—C15—C16	110.1 (4)
C15—N9—Ni2	108.0 (3)	N9—C15—H15A	109.6
C14—N9—H9A	114 (4)	C16—C15—H15A	109.6
C15—N9—H9A	100 (4)	N9—C15—H15B	109.6
Ni2—N9—H9A	107 (5)	C16—C15—H15B	109.6

C16—N10—Ni2	108.4 (3)	H15A—C15—H15B	108.2
C16—N10—H10C	108 (3)	N10—C16—C15	109.8 (3)
Ni2—N10—H10C	108 (3)	N10—C16—H16A	109.7
C16—N10—H10D	108 (5)	C15—C16—H16A	109.7
Ni2—N10—H10D	117 (5)	N10—C16—H16B	109.7
H10C—N10—H10D	107 (5)	C15—C16—H16B	109.7
C2—C1—C5	122.5 (3)	H16A—C16—H16B	108.2
C2—C1—S1	120.5 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N5—H5A \cdots N4 ⁱ	0.86 (3)	2.30 (4)	3.098 (6)	154 (3)
N5—H5B \cdots N2 ⁱⁱ	0.86 (3)	2.48 (3)	3.186 (5)	140 (4)
N7—H7A \cdots N3 ⁱⁱⁱ	0.86 (4)	2.56 (3)	3.207 (7)	134 (3)
N8—H8B \cdots N3 ⁱⁱⁱ	0.86 (3)	2.48 (4)	3.164 (6)	138 (3)
N9—H9A \cdots N1 ^{iv}	0.86 (2)	2.58 (3)	3.387 (6)	156 (5)
N10—H10C \cdots N2 ⁱⁱ	0.87 (3)	2.34 (3)	3.198 (5)	173 (5)

Symmetry codes: (i) $-x+1, -y+1, -z$; (ii) $x-1/2, -y+1/2, z+1/2$; (iii) $x+1, y, z$; (iv) $-x+5/2, y+1/2, -z+1/2$.

Fig. 1

