

2-[(Methylsulfanyl)methyl]-1,2-benz-isothiazol-3(2H)-one 1,1-dioxide

Waseeq Ahmad Siddiqui,^{a*} Saeed Ahmad,^b Hamid Latif Siddiqui,^c Rana Altaf Hussain^c and Masood Parvez^d

^aDepartment of Chemistry, University of Sargodha, Sargodha, Pakistan, ^bDepartment of Chemistry, University of Science and Technology, Bannu, Pakistan, ^cInstitute of Chemistry, University of the Punjab, Lahore, Pakistan, and ^dDepartment of Chemistry, University of Calgary, 2500 University Drive NW, Calgary, Alberta, Canada T2N 1N4

Correspondence e-mail: waseeq_786@yahoo.com


Received 16 August 2008; accepted 3 September 2008

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.036; wR factor = 0.094; data-to-parameter ratio = 17.1.

In the title molecule, $\text{C}_9\text{H}_9\text{NO}_3\text{S}_2$, the essentially planar benzisothiazole ring system and the $\text{C}-\text{S}-\text{C}$ atoms of the methylsulfanyl side chain form an angle of $64.45(7)^\circ$. The structure is devoid of any classical hydrogen bonding. However, weak non-classical inter- and intramolecular hydrogen bonds of the type $\text{C}-\text{H}\cdots\text{O}$ are present.

Related literature

For related literature, see: Bernstein *et al.* (1994); Masashi *et al.* (1999); Nagasawa *et al.* (1995); Siddiqui *et al.* (2007a,b, 2008a,b); Xu *et al.* (2006); Liang (2006).



Experimental

Crystal data

$\text{C}_9\text{H}_9\text{NO}_3\text{S}_2$
 $M_r = 243.29$
Monoclinic, $P2_1/c$
 $a = 7.550(3)\text{ \AA}$
 $b = 17.332(8)\text{ \AA}$
 $c = 9.455(3)\text{ \AA}$
 $\beta = 124.26(2)^\circ$

$V = 1022.6(7)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.51\text{ mm}^{-1}$
 $T = 173(2)\text{ K}$
 $0.18 \times 0.16 \times 0.06\text{ mm}$

Data collection

Nonius KappaCCD diffractometer
Absorption correction: multi-scan (*SORTAV*; Blessing, 1997)
 $T_{\min} = 0.915$, $T_{\max} = 0.970$
3975 measured reflections
2322 independent reflections
2004 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.094$
 $S = 1.05$
2322 reflections
136 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.29\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.45\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C2—H2 \cdots O2 ⁱ	0.95	2.49	3.390 (2)	158
C9—H9B \cdots O3	0.98	2.56	3.383 (3)	142

Symmetry code: (i) $-x + 2, -y, -z + 1$.

Data collection: *COLLECT* (Hooft, 1998); cell refinement: *HKL DENZO* (Otwinowski & Minor, 1997); data reduction: *SCALEPACK* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2683).

References

- Bernstein, J., Etter, M. C. & Leiserowitz, L. (1994). *Structure Correlation*, Vol. 2, edited by H.-B. Bürgi & J. D. Dunitz, pp. 431–507. New York: VCH.
- Blessing, R. H. (1997). *J. Appl. Cryst.* **30**, 421–426.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Hooft, R. (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Liang, X., Hong, S., Ying, L., Suhong, Z. & Mark, L. T. (2006). *Tetrahedron*, **62**, 7902–7910.
- Masashi, K., Hideo, T., Kentaro, Y. & Masataka, Y. (1999). *Tetrahedron*, **55**, 14885–14900.
- Nagasawa, H. T., Kawle, S. P., Elberling, J. A., DeMaster, E. G. & Fukuto, J. M. (1995). *J. Med. Chem.* **38**, 1865–1871.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Siddiqui, W. A., Ahmad, S., Khan, I. U., Siddiqui, H. L. & Parvez, M. (2007a). *Acta Cryst. E* **63**, o4116.
- Siddiqui, W. A., Ahmad, S., Siddiqui, H. L. & Parvez, M. (2008a). *Acta Cryst. E* **64**, o724.
- Siddiqui, W. A., Ahmad, S., Siddiqui, H. L., Parvez, M. & Rashid, R. (2008b). *Acta Cryst. E* **64**, o859.
- Siddiqui, W. A., Ahmad, S., Siddiqui, H. L., Tariq, M. I. & Parvez, M. (2007b). *Acta Cryst. E* **63**, o4001.
- Xu, L., Shu, H., Liu, Y., Zhang, S. & Trudell, M. (2006). *Tetrahedron*, **62**, 7902–7910.

supporting information

Acta Cryst. (2008). E64, o1897 [doi:10.1107/S1600536808028109]

2-[(Methylsulfanyl)methyl]-1,2-benzisothiazol-3(2H)-one 1,1-dioxide

Waseeq Ahmad Siddiqui, Saeed Ahmad, Hamid Latif Siddiqui, Rana Altaf Hussain and Masood Parvez

S1. Comment

1,2-benzisothiazole-3-one 1,1-dioxide (saccharin) has been identified as an important molecular component in various classes of 5-HT_{1A} antagonists, analgesics and human mast cell tryptase inhibitors (Liang *et al.*, 2006). Particularly, the substituted derivatives with e.g. *N*-hydroxy and *N*-alkyl substituents have shown important biological activities (Nagasawa *et al.*, 1995). Various biologically important saccharin skeletons and their *N*-alkyl derivatives were efficiently prepared (Xu *et al.*, 2006) by chromium oxide-catalyzed oxidation of *N*-alkyl(*o*-methyl)arenesulfonamides in acetonitrile besides the already developed methodology utilizing irradiation techniques (Masashi *et al.*, 1999) for similar type of conversions. In continuation of our research program on the synthesis of benzisothiazole derivatives (Siddiqui *et al.*, 2007a,b, 2008a,b), we report the synthesis (see Fig. 3) and crystal structure of the title compound, in this paper.

In the molecular structure (Fig. 1) the benzisothiazole rings system is essentially planar, the maximum deviation of any atom from the mean plane through S1/N1/C1–C7 being 0.0224 (8) Å for atom S1. The side chain comprising of atoms S2/C8/C9 is inclined at an angle 64.45 (7)° with the mean-plane of the benzisothiazole rings system. The structure is devoid of any classical hydrogen bonding. However, non-classical intermolecular hydrogen bond of the type C—H···O are present resulting in dimeric units in an *R*₂²(8) motif (Bernstein *et al.*, 1994). In addition, intramolecular hydrogen bonds of the type C—H···O are also present in the structure resulting in an *S*(7) pattern (Bernstein *et al.*, 1994) (details are in Fig. 2 and Table 1).

S2. Experimental

A suspension of saccharin (I) (1.0 g, 5.46 mmol), sodium sulfite (1.4 g, 10.93 mmol) and an excess of 2-chloro-5-methyl-aniline (5 ml) was first stirred at room temperature (30 min.) and then under reflux (1.5 hrs). The reaction mixture turned orange red after reflux. Cooled the reaction mixture to room temperature and extracted the product with chloroform (3 × 25 ml). Concentrated the organic layer under reduced pressure (11 torr) to get light yellow product (II) (0.6 g, 2.46 mmol), yield = 45%. Recrystallization Solvent: MeOH:CH₃CN (1:1). The solution was subjected to slow evaporation at 313 K to obtain colorless crystals.

S3. Refinement

Though all the H atoms could be distinguished in the difference Fourier map the H-atoms were included at geometrically idealized positions and refined in riding-model approximation with the following constraints: aryl, methyl and methylene C—H distances were set to 0.95, 0.98 and 0.99 Å, respectively; in all these instances $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$. The final difference map was free of any chemically significant features.






Figure 1

ORTEP-3 (Farrugia, 1997) drawing of the title compound with displacement ellipsoids plotted at 50% probability level.

**Figure 2**

Hydrogen bonding interactions in the unit cell of the title compound indicated by dashed lines, H-atoms not involved in H-bonds have been excluded.

**Scheme-I****Figure 3**

Reaction scheme.

2-[(Methylsulfanyl)methyl]-1,2-benzisothiazol-3(2H)-one 1,1-dioxide

Crystal data

$C_9H_9NO_3S_2$
 $M_r = 243.29$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 7.550 (3) \text{ \AA}$

$b = 17.332 (8) \text{ \AA}$
 $c = 9.455 (3) \text{ \AA}$
 $\beta = 124.26 (2)^\circ$
 $V = 1022.6 (7) \text{ \AA}^3$
 $Z = 4$

$F(000) = 504$
 $D_x = 1.580 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Cell parameters from 2322 reflections
 $\theta = 4.0\text{--}27.5^\circ$

$\mu = 0.51 \text{ mm}^{-1}$
 $T = 173 \text{ K}$
Plate, colourless
 $0.18 \times 0.16 \times 0.06 \text{ mm}$

Data collection

Nonius KappaCCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω and φ scans
Absorption correction: multi-scan
(SORTAV; Blessing, 1997)
 $T_{\min} = 0.915$, $T_{\max} = 0.970$

3975 measured reflections
2322 independent reflections
2004 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 4.0^\circ$
 $h = -9 \rightarrow 9$
 $k = -22 \rightarrow 19$
 $l = -12 \rightarrow 12$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.094$
 $S = 1.05$
2322 reflections
136 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.048P)^2 + 0.474P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.29 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.45 \text{ e \AA}^{-3}$

Special details

Experimental. m.p. 405–406 K; IR (KBr, ν_{\max} , cm^{-1}): CO 1731 (*s*), SO₂ 1332 and 1177; ¹H-NMR (400 MHz, DMSO-d₆) δ : 2.28 (s, 3H, CH₃), 4.90 (s, 2H, CH₂), 7.98–8.07 (m, 3H, aromatic), 8.13–8.34 (m, 1H, aromatic); ¹³C-NMR (100 MHz, DMSO-d₆) δ : 158.3, 136.7, 135.9, 135.3, 125.9, 125.1, 121.6, 42.6, 15.5 LRMS (ES⁺): *m/z*: 244 [M]⁺ (63.5%).

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	1.06232 (7)	0.08010 (3)	0.35109 (5)	0.02244 (14)
S2	1.41986 (7)	0.24291 (3)	0.32581 (6)	0.02667 (14)
O1	0.9265 (2)	0.19480 (7)	-0.03395 (16)	0.0267 (3)
O2	0.9634 (2)	0.10390 (9)	0.43595 (17)	0.0340 (3)
O3	1.2696 (2)	0.04531 (9)	0.45436 (16)	0.0328 (3)
N1	1.0689 (2)	0.15451 (9)	0.24102 (18)	0.0235 (3)
C1	0.8872 (3)	0.02719 (10)	0.1645 (2)	0.0197 (3)
C2	0.8048 (3)	-0.04560 (11)	0.1537 (2)	0.0244 (4)
H2	0.8436	-0.0741	0.2530	0.029*

C3	0.6623 (3)	-0.07501 (10)	-0.0102 (2)	0.0255 (4)
H3	0.6023	-0.1248	-0.0231	0.031*
C4	0.6060 (3)	-0.03308 (11)	-0.1554 (2)	0.0249 (4)
H4	0.5069	-0.0543	-0.2655	0.030*
C5	0.6929 (3)	0.03959 (11)	-0.1413 (2)	0.0224 (4)
H5	0.6559	0.0680	-0.2404	0.027*
C6	0.8348 (3)	0.06950 (10)	0.0209 (2)	0.0192 (3)
C7	0.9421 (3)	0.14577 (10)	0.0633 (2)	0.0202 (3)
C8	1.1816 (3)	0.22693 (11)	0.3227 (2)	0.0253 (4)
H8A	1.2215	0.2273	0.4420	0.030*
H8B	1.0817	0.2704	0.2618	0.030*
C9	1.6129 (3)	0.18644 (12)	0.5098 (2)	0.0320 (4)
H9A	1.7529	0.1899	0.5261	0.038*
H9B	1.5664	0.1325	0.4916	0.038*
H9C	1.6244	0.2063	0.6117	0.038*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0225 (2)	0.0275 (2)	0.0162 (2)	-0.00213 (16)	0.01023 (18)	-0.00010 (16)
S2	0.0260 (3)	0.0289 (3)	0.0266 (2)	-0.00508 (18)	0.0157 (2)	-0.00217 (18)
O1	0.0280 (7)	0.0252 (7)	0.0258 (6)	-0.0001 (5)	0.0145 (6)	0.0052 (5)
O2	0.0433 (8)	0.0417 (8)	0.0282 (7)	-0.0045 (7)	0.0270 (6)	-0.0055 (6)
O3	0.0238 (7)	0.0406 (8)	0.0223 (6)	0.0006 (6)	0.0059 (6)	0.0055 (6)
N1	0.0253 (8)	0.0243 (8)	0.0186 (7)	-0.0056 (6)	0.0109 (6)	-0.0024 (6)
C1	0.0178 (8)	0.0236 (9)	0.0176 (7)	0.0014 (6)	0.0099 (7)	0.0003 (6)
C2	0.0266 (9)	0.0235 (9)	0.0255 (8)	0.0020 (7)	0.0160 (8)	0.0046 (7)
C3	0.0251 (9)	0.0205 (9)	0.0324 (9)	-0.0007 (7)	0.0172 (8)	-0.0016 (7)
C4	0.0222 (8)	0.0275 (9)	0.0212 (8)	-0.0011 (7)	0.0099 (7)	-0.0046 (7)
C5	0.0216 (8)	0.0258 (9)	0.0180 (8)	0.0017 (7)	0.0102 (7)	0.0012 (7)
C6	0.0169 (8)	0.0223 (8)	0.0187 (8)	0.0013 (6)	0.0101 (7)	0.0009 (6)
C7	0.0174 (8)	0.0231 (9)	0.0204 (8)	0.0017 (6)	0.0108 (7)	0.0010 (6)
C8	0.0231 (9)	0.0247 (9)	0.0272 (9)	-0.0027 (7)	0.0135 (8)	-0.0067 (7)
C9	0.0227 (9)	0.0401 (11)	0.0274 (9)	-0.0025 (8)	0.0106 (8)	-0.0025 (8)

Geometric parameters (\AA , ^\circ)

S1—O3	1.4304 (15)	C3—C4	1.392 (3)
S1—O2	1.4306 (14)	C3—H3	0.9500
S1—N1	1.6754 (16)	C4—C5	1.392 (3)
S1—C1	1.7537 (18)	C4—H4	0.9500
S2—C8	1.804 (2)	C5—C6	1.385 (2)
S2—C9	1.804 (2)	C5—H5	0.9500
O1—C7	1.208 (2)	C6—C7	1.483 (2)
N1—C7	1.397 (2)	C8—H8A	0.9900
N1—C8	1.468 (2)	C8—H8B	0.9900
C1—C2	1.385 (3)	C9—H9A	0.9800
C1—C6	1.390 (2)	C9—H9B	0.9800

C2—C3	1.393 (3)	C9—H9C	0.9800
C2—H2	0.9500		
O3—S1—O2	117.06 (9)	C6—C5—C4	118.30 (16)
O3—S1—N1	110.18 (8)	C6—C5—H5	120.9
O2—S1—N1	109.50 (9)	C4—C5—H5	120.9
O3—S1—C1	112.44 (9)	C5—C6—C1	120.09 (16)
O2—S1—C1	112.28 (9)	C5—C6—C7	126.71 (15)
N1—S1—C1	92.68 (8)	C1—C6—C7	113.20 (15)
C8—S2—C9	100.92 (9)	O1—C7—N1	123.12 (16)
C7—N1—C8	121.81 (15)	O1—C7—C6	128.06 (15)
C7—N1—S1	115.07 (12)	N1—C7—C6	108.81 (14)
C8—N1—S1	122.85 (12)	N1—C8—S2	114.65 (12)
C2—C1—C6	122.64 (16)	N1—C8—H8A	108.6
C2—C1—S1	127.13 (13)	S2—C8—H8A	108.6
C6—C1—S1	110.22 (13)	N1—C8—H8B	108.6
C1—C2—C3	116.69 (16)	S2—C8—H8B	108.6
C1—C2—H2	121.7	H8A—C8—H8B	107.6
C3—C2—H2	121.7	S2—C9—H9A	109.5
C2—C3—C4	121.45 (17)	S2—C9—H9B	109.5
C2—C3—H3	119.3	H9A—C9—H9B	109.5
C4—C3—H3	119.3	S2—C9—H9C	109.5
C5—C4—C3	120.82 (16)	H9A—C9—H9C	109.5
C5—C4—H4	119.6	H9B—C9—H9C	109.5
C3—C4—H4	119.6		
O3—S1—N1—C7	116.17 (13)	C4—C5—C6—C1	0.0 (3)
O2—S1—N1—C7	-113.73 (13)	C4—C5—C6—C7	-179.11 (16)
C1—S1—N1—C7	1.06 (13)	C2—C1—C6—C5	0.8 (3)
O3—S1—N1—C8	-69.74 (16)	S1—C1—C6—C5	-178.13 (13)
O2—S1—N1—C8	60.36 (16)	C2—C1—C6—C7	-179.96 (15)
C1—S1—N1—C8	175.15 (14)	S1—C1—C6—C7	1.10 (18)
O3—S1—C1—C2	66.76 (18)	C8—N1—C7—O1	4.5 (3)
O2—S1—C1—C2	-67.75 (18)	S1—N1—C7—O1	178.63 (14)
N1—S1—C1—C2	179.89 (16)	C8—N1—C7—C6	-174.75 (14)
O3—S1—C1—C6	-114.36 (13)	S1—N1—C7—C6	-0.59 (17)
O2—S1—C1—C6	111.13 (13)	C5—C6—C7—O1	-0.4 (3)
N1—S1—C1—C6	-1.22 (13)	C1—C6—C7—O1	-179.53 (17)
C6—C1—C2—C3	-0.8 (3)	C5—C6—C7—N1	178.80 (16)
S1—C1—C2—C3	177.98 (13)	C1—C6—C7—N1	-0.37 (19)
C1—C2—C3—C4	0.0 (3)	C7—N1—C8—S2	-76.41 (19)
C2—C3—C4—C5	0.8 (3)	S1—N1—C8—S2	109.89 (14)
C3—C4—C5—C6	-0.8 (3)	C9—S2—C8—N1	-82.19 (15)

Hydrogen-bond geometry (\AA , $^\circ$)

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
C2—H2 \cdots O2 ⁱ	0.95	2.49	3.390 (2)	158

C9—H9 <i>B</i> ···O3	0.98	2.56	3.383 (3)	142
----------------------	------	------	-----------	-----

Symmetry code: (i) $-x+2, -y, -z+1$.