

Redetermination of 9,9'-bianthracene-10,10'(9H,9'H)-dione

Zhi-Gang Wen^{a*} and Jia-Ming Li^b

^aDepartment of Chemistry and Chemical Engineering, Qiannan Normal College for Nationalities, Duyun, Guizhou 558000, People's Republic of China, and

^bDepartment of Chemistry and Biology, Qinzhou University, Qinzhou, Guangxi 535000, People's Republic of China

Correspondence e-mail: ljmimarise@163.com


Received 29 August 2008; accepted 9 September 2008

Key indicators: single-crystal X-ray study; $T = 273\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.040; wR factor = 0.106; data-to-parameter ratio = 12.3.

The crystal structure of the title compound, $\text{C}_{28}\text{H}_{18}\text{O}_2$, was originally determined by Ehrenberg [(1967). *Acta Cryst.* **22**, 482–487] using intensity data obtained from Weissenberg photographs. The current determination provides a crystal and molecular structure with a significantly higher precision and presents standard uncertainties on geometric parameters which are not available from the original work. The molecule lies on a crystallographic twofold rotation axis which bisects the $\text{C}-\text{C}$ bond [1.603 (3) \AA] which joins the two anthracen-9(10H)-one units.

Related literature

For general background, see: Li *et al.* (2002); Shi *et al.* (2004); Müller *et al.* (1996, 1998, 2001); Prinz, Burgemeister & Wiegerebe (1996); Prinz, Wiegerebe & Müller (1996). For related structures, see: Ehrenberg (1967).



Experimental

Crystal data

$\text{C}_{28}\text{H}_{18}\text{O}_2$
 $M_r = 386.42$
Monoclinic, $C2/c$
 $a = 22.295$ (4) \AA
 $b = 7.7297$ (12) \AA
 $c = 13.643$ (2) \AA
 $\beta = 126.768$ (2) $^\circ$

$V = 1883.4$ (5) \AA^3
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.09\text{ mm}^{-1}$
 $T = 273$ (2) K
 $0.22 \times 0.18 \times 0.15\text{ mm}$

Data collection

Bruker SMART CCD
diffractometer
Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)
 $T_{\min} = 0.982$, $T_{\max} = 0.987$

4785 measured reflections
1669 independent reflections
1172 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.105$
 $S = 1.03$
1669 reflections

136 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.12\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.17\text{ e \AA}^{-3}$

Data collection: SMART (Bruker, 2003); cell refinement: SAINT (Bruker, 2003); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This work was supported by a key grant from Qiannan Normal College for Nationalities Foundation of Guizhou Province (grant No. 2007Z15) and the Qinzhou University Foundation of Guangxi Zhuang Autonomous Region of the People's Republic of China (grant No. 2008XJKY-10B).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2689).

References

- Bruker (2003). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Ehrenberg, M. (1967). *Acta Cryst.* **22**, 482–487.
- Li, P. C., Wang, T. S., Lee, G. H., Liu, Y. H., Wang, Y., Chen, C. T. & Chao, I. (2002). *J. Org. Chem.* **67**, 8002–8009.
- Müller, K., Altmann, R. & Prinz, H. (1998). *Eur. J. Med. Chem.* **33**, 209–214.
- Müller, K., Breu, K. & Reindl, H. (2001). *Eur. J. Med. Chem.* **36**, 179–184.
- Müller, K., Huang, H. S. & Wiegerebe, W. (1996). *J. Med. Chem.* **39**, 3132–3128.
- Prinz, H., Burgemeister, T. & Wiegerebe, W. (1996). *J. Org. Chem.* **61**, 2857–2860.
- Prinz, H., Wiegerebe, W. & Müller, K. (1996). *J. Org. Chem.* **61**, 2853–2856.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shi, Z.-W., Li, Y.-Z., Li, Y., Lu, G.-Y. & Liu, S.-H. (2004). *Acta Cryst. E* **60**, o2275–o2277.

supporting information

Acta Cryst. (2008). E64, o1931 [doi:10.1107/S1600536808028833]

Redetermination of 9,9'-bianthracene-10,10'(9H,9'H)-dione

Zhi-Gang Wen and Jia-Ming Li

S1. Comment


Synthesis of anthracenone derivatives have attracted great interest due to their interesting biological activities (Müller *et al.*, 1996, 1998, 2001; Prinz, Burgemeister & Wiegerebe, 1996; Prinz, Wiegerebe & Müller, 1996). Herein, we present a redetermination of the crystal structure of the title compound (I) which was originally refined in the non-conventional space group setting I2/a with unit cell parameters; $a = 13.68$ (4), $b = 7.751$ (3), $c = 17.92$ (4), $\beta = 91.1$ (3) (Ehrenberg, 1967). The current structure is of significantly higher precision than the orginal determination which was refined using intensity data obtained from Weissenberg photographs. The molecular structure of (I) is shown in Fig. 1. The molecule consists of two anthracen-9(10H)-one moieties linked together by a C—C [1.603 (3) Å] bond. A crystallographic twofold rotation axis bisects this bond.

S2. Experimental

Reagents and solvents used were of commercially available quality. The title complex (I) was synthesized according to the method of Shi *et al.* (2004) and Li *et al.* (2002). CF_3COOH (40 ml) was added dropwise with stirring to a solution of anthracene-9,10-dione (5.0 mmol) in 15 ml of anhydrous CH_2Cl_2 . The mixture was then placed in an ice bath and NaBH_4 (0.95 g, 25 mmol) was added in portions. The resulting mixture was stirred for 24 h at room temperature. The reaction mixture was poured into 200 ml ice-water. The organic layer was extracted with CH_2Cl_2 , dried over Na_2SO_4 and evaporated *in vacuo*. The crude product was recrystallized from toluene twice to give the main product 9,9'-bianthracene-10,10'(9H,9'H)-dione.

S3. Refinement

H atoms were positioned geometrically and refined using a riding model, with C—H = 0.93–0.98 Å and with $U_{\text{iso}}(\text{H}) = 1.2$ times $U_{\text{eq}}(\text{C})$.

**Figure 1**

The molecular structure with displacement ellipsoids at the 30% probability level [symmetry code: (A) $-x+2, y, -z+1/2$].

9,9'-bianthracene-10,10'(9H,9'H)-dione

Crystal data

$C_{28}H_{18}O_2$

$M_r = 386.42$

Monoclinic, $C2/c$

Hall symbol: $-C\bar{2}yc$

$a = 22.295 (4)$ Å

$b = 7.7297 (12)$ Å

$c = 13.643 (2)$ Å

$\beta = 126.768 (2)^\circ$

$V = 1883.4 (5)$ Å 3

$Z = 4$

$F(000) = 808$

$D_x = 1.363$ Mg m $^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1069 reflections

$\theta = 2.9\text{--}24.7^\circ$

$\mu = 0.09$ mm $^{-1}$

$T = 273$ K

Block, yellow

$0.22 \times 0.18 \times 0.15$ mm

Data collection

Bruker SMART CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.982$, $T_{\max} = 0.987$

4785 measured reflections
1669 independent reflections
1172 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$
 $\theta_{\max} = 25.1^\circ$, $\theta_{\min} = 2.3^\circ$
 $h = -20 \rightarrow 26$
 $k = -9 \rightarrow 9$
 $l = -16 \rightarrow 8$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.106$
 $S = 1.03$
1669 reflections
136 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0478P)^2 + 0.4019P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.12 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.17 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.86707 (8)	-0.25682 (16)	0.29866 (15)	0.0855 (5)
C1	0.90675 (10)	-0.1382 (2)	0.30813 (16)	0.0503 (4)
C2	0.98793 (9)	-0.16248 (19)	0.37296 (14)	0.0423 (4)
C3	1.02019 (10)	-0.3196 (2)	0.43139 (15)	0.0546 (5)
H3	0.9900	-0.4083	0.4254	0.066*
C4	1.09595 (11)	-0.3451 (2)	0.49762 (16)	0.0616 (5)
H4	1.1169	-0.4513	0.5346	0.074*
C5	1.14056 (10)	-0.2121 (2)	0.50886 (15)	0.0572 (5)
H5	1.1921	-0.2277	0.5553	0.069*
C6	1.10969 (8)	-0.0560 (2)	0.45207 (13)	0.0457 (4)
H6	1.1408	0.0332	0.4616	0.055*
C7	1.03286 (8)	-0.02957 (19)	0.38070 (13)	0.0379 (4)
C8	0.99831 (8)	0.13334 (18)	0.30704 (13)	0.0370 (4)
H8	1.0288	0.2306	0.3599	0.044*
C9	0.91944 (8)	0.16565 (19)	0.26300 (13)	0.0393 (4)
C10	0.88717 (9)	0.3276 (2)	0.21705 (15)	0.0491 (4)

H10	0.9158	0.4170	0.2191	0.059*
C11	0.81329 (10)	0.3570 (2)	0.16860 (16)	0.0591 (5)
H11	0.7926	0.4660	0.1382	0.071*
C12	0.76987 (10)	0.2268 (3)	0.16480 (16)	0.0589 (5)
H12	0.7199	0.2471	0.1314	0.071*
C13	0.80076 (9)	0.0673 (2)	0.21051 (15)	0.0535 (5)
H13	0.7716	-0.0207	0.2086	0.064*
C14	0.87527 (8)	0.0352 (2)	0.25983 (14)	0.0426 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0740 (10)	0.0554 (8)	0.1455 (14)	-0.0116 (7)	0.0756 (10)	0.0060 (8)
C1	0.0571 (11)	0.0449 (10)	0.0648 (11)	-0.0082 (8)	0.0450 (9)	-0.0049 (8)
C2	0.0527 (10)	0.0383 (9)	0.0436 (9)	-0.0017 (7)	0.0331 (8)	-0.0012 (7)
C3	0.0705 (13)	0.0430 (10)	0.0553 (11)	-0.0008 (9)	0.0403 (10)	0.0061 (8)
C4	0.0735 (14)	0.0512 (11)	0.0535 (11)	0.0147 (10)	0.0344 (10)	0.0138 (9)
C5	0.0496 (11)	0.0651 (12)	0.0458 (10)	0.0120 (9)	0.0227 (8)	0.0111 (9)
C6	0.0446 (10)	0.0512 (10)	0.0392 (9)	0.0003 (8)	0.0240 (8)	0.0024 (8)
C7	0.0447 (9)	0.0388 (9)	0.0327 (8)	0.0000 (7)	0.0245 (7)	-0.0022 (7)
C8	0.0394 (9)	0.0327 (8)	0.0406 (9)	-0.0041 (6)	0.0247 (7)	-0.0050 (7)
C9	0.0427 (9)	0.0384 (8)	0.0414 (9)	-0.0003 (7)	0.0276 (7)	-0.0074 (7)
C10	0.0502 (11)	0.0421 (9)	0.0571 (11)	0.0027 (8)	0.0333 (9)	-0.0037 (8)
C11	0.0568 (12)	0.0566 (11)	0.0606 (12)	0.0179 (9)	0.0333 (9)	0.0026 (9)
C12	0.0414 (10)	0.0781 (14)	0.0570 (11)	0.0074 (9)	0.0293 (9)	-0.0030 (10)
C13	0.0464 (10)	0.0645 (12)	0.0558 (11)	-0.0060 (9)	0.0339 (9)	-0.0081 (9)
C14	0.0429 (9)	0.0472 (9)	0.0451 (9)	-0.0033 (7)	0.0303 (8)	-0.0070 (7)

Geometric parameters (\AA , ^\circ)

O1—C1	1.2260 (18)	C8—C9	1.500 (2)
C1—C14	1.473 (2)	C8—C8 ⁱ	1.603 (3)
C1—C2	1.475 (2)	C8—H8	0.9800
C2—C3	1.393 (2)	C9—C10	1.392 (2)
C2—C7	1.394 (2)	C9—C14	1.392 (2)
C3—C4	1.372 (2)	C10—C11	1.378 (2)
C3—H3	0.9300	C10—H10	0.9300
C4—C5	1.374 (2)	C11—C12	1.376 (3)
C4—H4	0.9300	C11—H11	0.9300
C5—C6	1.375 (2)	C12—C13	1.367 (2)
C5—H5	0.9300	C12—H12	0.9300
C6—C7	1.388 (2)	C13—C14	1.391 (2)
C6—H6	0.9300	C13—H13	0.9300
C7—C8	1.504 (2)		
O1—C1—C14	120.85 (16)	C7—C8—C8 ⁱ	110.23 (10)
O1—C1—C2	121.12 (16)	C9—C8—H8	107.4
C14—C1—C2	117.97 (14)	C7—C8—H8	107.4

C3—C2—C7	119.84 (16)	C8 ⁱ —C8—H8	107.4
C3—C2—C1	118.80 (15)	C10—C9—C14	118.18 (15)
C7—C2—C1	121.31 (14)	C10—C9—C8	119.74 (14)
C4—C3—C2	120.84 (17)	C14—C9—C8	121.98 (13)
C4—C3—H3	119.6	C11—C10—C9	120.78 (16)
C2—C3—H3	119.6	C11—C10—H10	119.6
C3—C4—C5	119.33 (17)	C9—C10—H10	119.6
C3—C4—H4	120.3	C12—C11—C10	120.62 (17)
C5—C4—H4	120.3	C12—C11—H11	119.7
C4—C5—C6	120.59 (17)	C10—C11—H11	119.7
C4—C5—H5	119.7	C13—C12—C11	119.45 (17)
C6—C5—H5	119.7	C13—C12—H12	120.3
C5—C6—C7	121.03 (16)	C11—C12—H12	120.3
C5—C6—H6	119.5	C12—C13—C14	120.73 (17)
C7—C6—H6	119.5	C12—C13—H13	119.6
C6—C7—C2	118.28 (14)	C14—C13—H13	119.6
C6—C7—C8	121.01 (14)	C13—C14—C9	120.23 (15)
C2—C7—C8	120.59 (14)	C13—C14—C1	119.33 (15)
C9—C8—C7	114.57 (13)	C9—C14—C1	120.43 (14)
C9—C8—C8 ⁱ	109.65 (14)		
O1—C1—C2—C3	-4.6 (2)	C7—C8—C9—C10	166.38 (13)
C14—C1—C2—C3	172.72 (15)	C8 ⁱ —C8—C9—C10	-69.07 (14)
O1—C1—C2—C7	178.00 (16)	C7—C8—C9—C14	-17.4 (2)
C14—C1—C2—C7	-4.7 (2)	C8 ⁱ —C8—C9—C14	107.14 (13)
C7—C2—C3—C4	0.6 (2)	C14—C9—C10—C11	-0.7 (2)
C1—C2—C3—C4	-176.90 (15)	C8—C9—C10—C11	175.66 (14)
C2—C3—C4—C5	1.7 (3)	C9—C10—C11—C12	0.1 (3)
C3—C4—C5—C6	-1.5 (3)	C10—C11—C12—C13	0.5 (3)
C4—C5—C6—C7	-1.0 (3)	C11—C12—C13—C14	-0.5 (3)
C5—C6—C7—C2	3.2 (2)	C12—C13—C14—C9	-0.1 (2)
C5—C6—C7—C8	-172.91 (15)	C12—C13—C14—C1	-179.91 (16)
C3—C2—C7—C6	-3.0 (2)	C10—C9—C14—C13	0.7 (2)
C1—C2—C7—C6	174.42 (14)	C8—C9—C14—C13	-175.56 (13)
C3—C2—C7—C8	173.14 (14)	C10—C9—C14—C1	-179.51 (14)
C1—C2—C7—C8	-9.5 (2)	C8—C9—C14—C1	4.2 (2)
C6—C7—C8—C9	-164.09 (13)	O1—C1—C14—C13	4.5 (3)
C2—C7—C8—C9	19.91 (19)	C2—C1—C14—C13	-172.84 (14)
C6—C7—C8—C8 ⁱ	71.68 (18)	O1—C1—C14—C9	-175.31 (16)
C2—C7—C8—C8 ⁱ	-104.32 (17)	C2—C1—C14—C9	7.4 (2)

Symmetry code: (i) $-x+2, y, -z+1/2$.