

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bis(μ -*N,N'*-di-3-pyridyl-2,6-pyridine-2,6-dicarboxamide- κ^2 N:N')bis[dibromido-mercury(II)] *N,N*-dimethylformamide disolvate

Li-hua Huang and Jie Wu*

Department of Chemistry, Zhengzhou University, Zhengzhou 450052, People's Republic of China

Correspondence e-mail: wujie@zzu.edu.cn


Received 26 August 2008; accepted 8 September 2008

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.037; wR factor = 0.067; data-to-parameter ratio = 17.8.

In the dinuclear centrosymmetric title complex, $[\text{Hg}_2\text{Br}_4(\text{C}_{17}\text{H}_{13}\text{N}_5\text{O}_2)_2]\cdot 2\text{C}_3\text{H}_7\text{NO}$, the Hg^{II} atom is coordinated by two Br atoms and two N atoms from two different ligands in a distorted tetrahedral geometry. The solvent molecule is linked to the 28-atom ring by two hydrogen bonds.

Related literature

For related literature, see: Baer *et al.* (2002); Chae *et al.* (2004 and references cited therein); Qin *et al.* (2003).



Experimental

Crystal data

$[\text{Hg}_2\text{Br}_4(\text{C}_{17}\text{H}_{13}\text{N}_5\text{O}_2)_2]\cdot 2\text{C}_3\text{H}_7\text{NO}$
 $M_r = 1505.62$
 Triclinic, $P\bar{1}$
 $a = 7.7609$ (16) Å
 $b = 12.267$ (3) Å
 $c = 13.296$ (3) Å
 $\alpha = 92.27$ (3)°
 $\beta = 105.82$ (3)°

$\gamma = 104.07$ (3)°
 $V = 1173.7$ (4) Å³
 $Z = 1$
 Mo $K\alpha$ radiation
 $\mu = 10.00$ mm⁻¹
 $T = 293$ (2) K
 $0.20 \times 0.18 \times 0.17$ mm

Data collection

Rigaku Saturn724 diffractometer
 Absorption correction: multi-scan
 (*CrystalClear*; Rigaku/MSC,
 2006)
 $T_{\text{min}} = 0.240$, $T_{\text{max}} = 0.281$
 (expected range = 0.156–0.183)

14249 measured reflections
 5337 independent reflections
 4364 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.067$
 $S = 1.03$
 5337 reflections
 299 parameters
 2 restraints

H atoms treated by a mixture of
 independent and constrained
 refinement
 $\Delta\rho_{\text{max}} = 0.61$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.71$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N4}-\text{H22}\cdots\text{O3}^{\text{i}}$	0.859 (10)	2.08 (2)	2.891 (5)	157 (4)
$\text{N2}-\text{H21}\cdots\text{O3}^{\text{i}}$	0.856 (10)	2.34 (2)	3.076 (5)	144 (3)
$\text{N2}-\text{H21}\cdots\text{N3}$	0.856 (10)	2.25 (4)	2.685 (5)	111 (3)

 Symmetry code: (i) $-x + 1, -y + 1, -z + 1$.

Data collection: *CrystalClear* (Rigaku/MSC, 2006); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank Professor Hou Hong-Wei of Zhengzhou University for his help.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2488).

References

- Baer, A. J., Koivisto, B. D., Coté, A. P., Taylor, N. J., Hanan, G. S., Nierengarten, H. & Dorsselaer, A. V. (2002). *Inorg. Chem.* **41**, 4987–4989.
 Chae, H. E., Siberio-Pérez, D. Y., Kim, J., Go, Y., Eddaoudi, M., Matzger, A. J., O'Keefe, M. & Yaghi, O. M. (2004). *Nature (London)*, **427**, 523–527.
 Qin, Z.-Q., Jennings, M. C. & Puddephatt, R. J. (2003). *Inorg. Chem.* **42**, 1956–1965.
 Rigaku/MSC (2006). *CrystalClear*. Rigaku/MSC, The Woodlands, Texas, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, m1263 [doi:10.1107/S1600536808028754]

Bis(μ -*N,N'*-di-3-pyridyl-2,6-pyridine-2,6-dicarboxamide- κ^2 *N:N'*)bis[dibromidomercury(II)] *N,N*-dimethylformamide disolvate

L. Huang and J. Wu

Comment

Metal-organic frameworks (MOFs) with microporous is currently of great interest because of their interesting structures and potential applications. So far, some interesting microporous MOFs have been documented (Chae *et al.* 2004, and references cited therein). One of the popular strategies to fabricate such compounds is to design the rigid ligands which have the ability to bridge the metal centers with big ring by utilizing their coordination sites. The rigid conjugated clamp-like multi-pyridine ligand *N,N'*-bis(pyridin-3-yl)-2,6-pyridinedicarboxamide has been known as a good candidate in the construction of MOFs with big ring (Qin *et al.* 2003; Baer *et al.* 2002). In this work, we selected this ligand as linker, generating a new coordination complex, [HgIIBr₂(C₁₇N₅O₂)](DMF), (I), which is reported here. In compound (I) each HgII atom is four-coordinated by two N atoms from two ligands and two Br atoms in a distorted tetrahedral coordination sphere (Fig. 1). The two HgII atoms are bridged with two *N,N'*-bis(pyridin-3-yl)-2,6-pyridinedicarboxamide ligands to form a microporous MOFs with 28-number ring. The neighbouring units are linked by the interactions to form a two-dimensional network (Fig. 2) and hydrogen bonds arising between the DMF and *N,N'*-bis(pyridin-3-yl)-2,6-pyridinedicarboxamide ligand (Table 2) complete the structure.

Experimental

The ligand *N,N'*-bis(pyridin-3-yl)-2,6-pyridinedicarboxamide (0.05 mmol, 0.016 g) in DMF (5 ml) was added dropwise to a solution of HgBr₂ (0.1 mmol, 0.036 g) in methanol (3 ml). The precipitate was filtered and the resulting solution was allowed to stand at room temperature in the dark. After one week good quality colorless crystals were obtained and dried in air.

Figures




Fig. 1. View of the title complex, showing the labeling of the non-H atoms and 30% probability ellipsoids. H atoms have been omitted.




Fig. 2. A view of the crystal packing along the *a* axis. Hydrogen bonds are shown as dashed lines.




Fig. 3. Supplementary figure.

Bis(μ -*N,N'*-di-3-pyridylpyridine-2,6-dicarboxamide- $\kappa^2N:N'$)bis[dibromidomercury(II)] *N,N*-dimethylformamide disolvate

Crystal data

[Hg₂Br₄(C₁₇H₁₃N₅O₂)₂]·2C₃H₇NO

$M_r = 1505.62$

Triclinic, $P\bar{1}$

$a = 7.7609$ (16) Å

$b = 12.267$ (3) Å

$c = 13.296$ (3) Å

$\alpha = 92.27$ (3)°

$\beta = 105.82$ (3)°

$\gamma = 104.07$ (3)°

$V = 1173.7$ (4) Å³

$Z = 1$

$F_{000} = 712$

$D_x = 2.130$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 3306 reflections

$\theta = 3.2$ – 27.5 °

$\mu = 10.00$ mm⁻¹

$T = 293$ (2) K

Prism, colourless

$0.20 \times 0.18 \times 0.17$ mm

Data collection

Rigaku Saturn724
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 28.5714 pixels mm⁻¹

$T = 293$ (2) K

dtprofit.ref scans

Absorption correction: multi-scan
(CrystalClear; Rigaku/MSC, 2006)

$T_{\min} = 0.240$, $T_{\max} = 0.281$

14249 measured reflections

5337 independent reflections

4364 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.032$

$\theta_{\text{max}} = 27.5$ °

$\theta_{\text{min}} = 3.2$ °

$h = -10 \rightarrow 10$

$k = -15 \rightarrow 15$

$l = -17 \rightarrow 17$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.037$

$wR(F^2) = 0.067$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0274P)^2]$

$S = 1.03$

5337 reflections

299 parameters

2 restraints

Primary atom site location: structure-invariant direct methods

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.61 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.71 \text{ e } \text{\AA}^{-3}$

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Hg1	0.35413 (3)	0.143038 (15)	0.325732 (13)	0.04901 (8)
Br1	0.37301 (7)	-0.05158 (4)	0.27313 (4)	0.05602 (14)
Br2	0.63842 (7)	0.30874 (4)	0.39232 (4)	0.06222 (15)
O1	0.8838 (6)	0.5153 (3)	1.1145 (3)	0.0790 (12)
O2	0.1707 (5)	0.0697 (3)	0.7966 (2)	0.0565 (9)
N1	0.8086 (5)	0.7906 (3)	0.8251 (3)	0.0394 (8)
N2	0.7259 (5)	0.5300 (3)	0.9465 (3)	0.0399 (9)
N3	0.5289 (5)	0.3133 (3)	0.9248 (2)	0.0342 (8)
N4	0.3039 (5)	0.2237 (3)	0.7291 (3)	0.0375 (8)
N5	0.1886 (5)	0.1582 (3)	0.4441 (3)	0.0388 (8)
C1	0.9457 (6)	0.8600 (4)	0.9012 (3)	0.0410 (11)
H1	0.9948	0.9335	0.8888	0.049*
C2	1.0151 (6)	0.8257 (4)	0.9966 (3)	0.0458 (11)
H2	1.1084	0.8766	1.0485	0.055*
C3	0.9496 (6)	0.7169 (4)	1.0172 (3)	0.0429 (11)
H3	0.9979	0.6935	1.0822	0.051*
C4	0.8082 (6)	0.6423 (3)	0.9377 (3)	0.0323 (9)
C5	0.7417 (6)	0.6847 (3)	0.8436 (3)	0.0356 (10)
H5	0.6454	0.6369	0.7907	0.043*
C6	0.7650 (6)	0.4739 (4)	1.0323 (3)	0.0423 (11)
C7	0.6516 (6)	0.3533 (3)	1.0186 (3)	0.0376 (10)
C8	0.6773 (7)	0.2900 (4)	1.1030 (3)	0.0466 (12)
H8	0.7648	0.3211	1.1670	0.056*
C9	0.5708 (7)	0.1802 (4)	1.0900 (3)	0.0477 (12)
H9	0.5851	0.1356	1.1451	0.057*

supplementary materials

C10	0.4421 (6)	0.1373 (4)	0.9935 (4)	0.0421 (11)
H10	0.3679	0.0633	0.9827	0.050*
C11	0.4256 (6)	0.2061 (3)	0.9135 (3)	0.0353 (10)
C12	0.2874 (6)	0.1602 (3)	0.8079 (3)	0.0380 (10)
C13	0.1959 (6)	0.1874 (3)	0.6234 (3)	0.0352 (10)
C14	0.0045 (6)	0.1485 (3)	0.5954 (4)	0.0418 (11)
H14	-0.0582	0.1444	0.6462	0.050*
C15	-0.0910 (6)	0.1160 (4)	0.4913 (4)	0.0468 (12)
H15	-0.2198	0.0901	0.4707	0.056*
C16	0.0035 (7)	0.1218 (4)	0.4176 (4)	0.0463 (11)
H16	-0.0629	0.0998	0.3472	0.056*
C17	0.2825 (6)	0.1910 (3)	0.5445 (3)	0.0343 (9)
H17	0.4111	0.2175	0.5627	0.041*
H22	0.396 (4)	0.283 (2)	0.745 (3)	0.044 (13)*
H21	0.642 (4)	0.491 (3)	0.8924 (19)	0.039 (12)*
O3	0.4162 (5)	0.5774 (3)	0.2838 (2)	0.0568 (9)
N6	0.1662 (6)	0.5374 (3)	0.3463 (3)	0.0465 (10)
C18	-0.0003 (7)	0.5662 (5)	0.3550 (4)	0.0653 (15)
H18A	-0.0186	0.6282	0.3152	0.098*
H18B	0.0132	0.5875	0.4275	0.098*
H18C	-0.1054	0.5019	0.3280	0.098*
C19	0.2125 (9)	0.4419 (4)	0.3969 (4)	0.0719 (17)
H19A	0.3199	0.4287	0.3813	0.108*
H19B	0.1099	0.3759	0.3715	0.108*
H19C	0.2385	0.4576	0.4716	0.108*
C20	0.2735 (8)	0.5963 (4)	0.2962 (3)	0.0503 (12)
H20A	0.2392	0.6583	0.2671	0.060*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Hg1	0.06762 (15)	0.04080 (12)	0.03641 (11)	0.01450 (9)	0.01118 (9)	0.00594 (8)
Br1	0.0631 (3)	0.0433 (3)	0.0679 (3)	0.0189 (2)	0.0251 (3)	0.0047 (2)
Br2	0.0546 (3)	0.0547 (3)	0.0623 (3)	0.0063 (3)	0.0000 (3)	0.0042 (3)
O1	0.105 (3)	0.053 (2)	0.039 (2)	-0.011 (2)	-0.016 (2)	0.0158 (17)
O2	0.067 (2)	0.0389 (18)	0.049 (2)	-0.0133 (17)	0.0173 (17)	0.0083 (16)
N1	0.044 (2)	0.035 (2)	0.0344 (19)	0.0063 (17)	0.0071 (17)	0.0039 (16)
N2	0.047 (2)	0.037 (2)	0.0269 (19)	0.0029 (18)	0.0026 (17)	0.0033 (17)
N3	0.041 (2)	0.0331 (19)	0.0315 (19)	0.0101 (16)	0.0143 (16)	0.0078 (16)
N4	0.042 (2)	0.0298 (19)	0.034 (2)	-0.0031 (17)	0.0113 (17)	0.0018 (16)
N5	0.048 (2)	0.0330 (19)	0.0318 (19)	0.0080 (17)	0.0082 (17)	0.0027 (16)
C1	0.044 (3)	0.031 (2)	0.045 (3)	0.004 (2)	0.013 (2)	0.005 (2)
C2	0.045 (3)	0.043 (3)	0.036 (2)	-0.002 (2)	0.001 (2)	0.001 (2)
C3	0.044 (3)	0.040 (3)	0.035 (2)	0.006 (2)	0.001 (2)	0.004 (2)
C4	0.036 (2)	0.031 (2)	0.028 (2)	0.0082 (18)	0.0086 (18)	0.0045 (17)
C5	0.040 (3)	0.033 (2)	0.027 (2)	0.0048 (19)	0.0043 (19)	0.0013 (18)
C6	0.052 (3)	0.040 (3)	0.030 (2)	0.010 (2)	0.005 (2)	0.010 (2)
C7	0.046 (3)	0.038 (2)	0.033 (2)	0.013 (2)	0.014 (2)	0.011 (2)

C8	0.056 (3)	0.051 (3)	0.036 (2)	0.014 (2)	0.016 (2)	0.012 (2)
C9	0.060 (3)	0.048 (3)	0.038 (3)	0.014 (2)	0.018 (2)	0.018 (2)
C10	0.052 (3)	0.034 (2)	0.050 (3)	0.013 (2)	0.027 (2)	0.012 (2)
C11	0.044 (3)	0.030 (2)	0.036 (2)	0.0066 (19)	0.021 (2)	0.0063 (18)
C12	0.044 (3)	0.032 (2)	0.042 (3)	0.009 (2)	0.020 (2)	0.004 (2)
C13	0.039 (3)	0.027 (2)	0.038 (2)	0.0062 (18)	0.011 (2)	0.0052 (18)
C14	0.041 (3)	0.033 (2)	0.054 (3)	0.010 (2)	0.018 (2)	0.006 (2)
C15	0.040 (3)	0.032 (2)	0.060 (3)	0.007 (2)	0.003 (2)	0.004 (2)
C16	0.057 (3)	0.033 (2)	0.042 (3)	0.014 (2)	0.000 (2)	0.004 (2)
C17	0.036 (2)	0.032 (2)	0.032 (2)	0.0054 (18)	0.0080 (19)	0.0023 (18)
O3	0.059 (2)	0.053 (2)	0.052 (2)	-0.0049 (18)	0.0235 (18)	0.0038 (17)
N6	0.058 (3)	0.038 (2)	0.044 (2)	0.0050 (19)	0.021 (2)	0.0033 (18)
C18	0.063 (4)	0.070 (4)	0.063 (4)	0.018 (3)	0.022 (3)	-0.010 (3)
C19	0.104 (5)	0.049 (3)	0.080 (4)	0.024 (3)	0.048 (4)	0.023 (3)
C20	0.065 (3)	0.041 (3)	0.036 (3)	0.002 (3)	0.010 (3)	0.000 (2)

Geometric parameters (Å, °)

Hg1—N5	2.315 (3)	C7—C8	1.386 (6)
Hg1—N1 ⁱ	2.351 (3)	C8—C9	1.374 (6)
Hg1—Br1	2.5108 (8)	C8—H8	0.9300
Hg1—Br2	2.5289 (12)	C9—C10	1.383 (6)
O1—C6	1.218 (5)	C9—H9	0.9300
O2—C12	1.225 (5)	C10—C11	1.382 (6)
N1—C5	1.335 (5)	C10—H10	0.9300
N1—C1	1.337 (5)	C11—C12	1.503 (6)
N1—Hg1 ⁱ	2.351 (3)	C13—C14	1.384 (6)
N2—C6	1.357 (5)	C13—C17	1.389 (5)
N2—C4	1.394 (5)	C14—C15	1.369 (6)
N2—H21	0.856 (10)	C14—H14	0.9300
N3—C7	1.334 (5)	C15—C16	1.370 (6)
N3—C11	1.342 (5)	C15—H15	0.9300
N4—C12	1.346 (5)	C16—H16	0.9300
N4—C13	1.414 (5)	C17—H17	0.9300
N4—H22	0.859 (10)	O3—C20	1.236 (6)
N5—C17	1.326 (5)	N6—C20	1.306 (6)
N5—C16	1.337 (6)	N6—C19	1.446 (6)
C1—C2	1.361 (6)	N6—C18	1.452 (6)
C1—H1	0.9300	C18—H18A	0.9600
C2—C3	1.374 (6)	C18—H18B	0.9600
C2—H2	0.9300	C18—H18C	0.9600
C3—C4	1.401 (5)	C19—H19A	0.9600
C3—H3	0.9300	C19—H19B	0.9600
C4—C5	1.390 (5)	C19—H19C	0.9600
C5—H5	0.9300	C20—H20A	0.9300
C6—C7	1.502 (6)		
N5—Hg1—N1 ⁱ	103.47 (12)	C8—C9—H9	120.6
N5—Hg1—Br1	117.00 (9)	C10—C9—H9	120.6

supplementary materials

N1 ⁱ —Hg1—Br1	107.93 (9)	C11—C10—C9	118.8 (4)
N5—Hg1—Br2	102.81 (9)	C11—C10—H10	120.6
N1 ⁱ —Hg1—Br2	100.41 (9)	C9—C10—H10	120.6
Br1—Hg1—Br2	122.54 (3)	N3—C11—C10	123.1 (4)
C5—N1—C1	118.4 (4)	N3—C11—C12	117.5 (3)
C5—N1—Hg1 ⁱ	118.6 (3)	C10—C11—C12	119.4 (4)
C1—N1—Hg1 ⁱ	122.0 (3)	O2—C12—N4	123.6 (4)
C6—N2—C4	126.9 (4)	O2—C12—C11	120.7 (4)
C6—N2—H21	116 (3)	N4—C12—C11	115.7 (4)
C4—N2—H21	118 (3)	C14—C13—C17	118.3 (4)
C7—N3—C11	117.1 (3)	C14—C13—N4	122.0 (4)
C12—N4—C13	122.6 (4)	C17—C13—N4	119.8 (4)
C12—N4—H22	116 (3)	C15—C14—C13	118.7 (4)
C13—N4—H22	121 (3)	C15—C14—H14	120.7
C17—N5—C16	118.9 (4)	C13—C14—H14	120.7
C17—N5—Hg1	118.2 (3)	C14—C15—C16	119.9 (4)
C16—N5—Hg1	122.4 (3)	C14—C15—H15	120.0
N1—C1—C2	121.6 (4)	C16—C15—H15	120.0
N1—C1—H1	119.2	N5—C16—C15	121.8 (4)
C2—C1—H1	119.2	N5—C16—H16	119.1
C1—C2—C3	121.0 (4)	C15—C16—H16	119.1
C1—C2—H2	119.5	N5—C17—C13	122.5 (4)
C3—C2—H2	119.5	N5—C17—H17	118.8
C2—C3—C4	118.2 (4)	C13—C17—H17	118.8
C2—C3—H3	120.9	C20—N6—C19	120.5 (4)
C4—C3—H3	120.9	C20—N6—C18	121.7 (4)
C5—C4—N2	117.6 (4)	C19—N6—C18	117.7 (4)
C5—C4—C3	117.2 (4)	N6—C18—H18A	109.5
N2—C4—C3	125.2 (4)	N6—C18—H18B	109.5
N1—C5—C4	123.5 (4)	H18A—C18—H18B	109.5
N1—C5—H5	118.2	N6—C18—H18C	109.5
C4—C5—H5	118.2	H18A—C18—H18C	109.5
O1—C6—N2	124.0 (4)	H18B—C18—H18C	109.5
O1—C6—C7	121.1 (4)	N6—C19—H19A	109.5
N2—C6—C7	114.9 (4)	N6—C19—H19B	109.5
N3—C7—C8	123.5 (4)	H19A—C19—H19B	109.5
N3—C7—C6	117.3 (3)	N6—C19—H19C	109.5
C8—C7—C6	119.2 (4)	H19A—C19—H19C	109.5
C9—C8—C7	118.7 (4)	H19B—C19—H19C	109.5
C9—C8—H8	120.7	O3—C20—N6	126.2 (5)
C7—C8—H8	120.7	O3—C20—H20A	116.9
C8—C9—C10	118.8 (4)	N6—C20—H20A	116.9
N1 ⁱ —Hg1—N5—C17	-141.8 (3)	C6—C7—C8—C9	179.1 (4)
Br1—Hg1—N5—C17	99.7 (3)	C7—C8—C9—C10	-0.1 (7)
Br2—Hg1—N5—C17	-37.6 (3)	C8—C9—C10—C11	0.2 (7)
N1 ⁱ —Hg1—N5—C16	46.8 (3)	C7—N3—C11—C10	-0.3 (6)
Br1—Hg1—N5—C16	-71.7 (3)	C7—N3—C11—C12	-179.8 (3)

Br2—Hg1—N5—C16	151.0 (3)	C9—C10—C11—N3	0.0 (6)
C5—N1—C1—C2	-1.1 (6)	C9—C10—C11—C12	179.5 (4)
Hg1 ⁱ —N1—C1—C2	167.2 (3)	C13—N4—C12—O2	-4.8 (7)
N1—C1—C2—C3	1.6 (7)	C13—N4—C12—C11	174.0 (4)
C1—C2—C3—C4	-0.5 (7)	N3—C11—C12—O2	-169.3 (4)
C6—N2—C4—C5	-177.3 (4)	C10—C11—C12—O2	11.2 (6)
C6—N2—C4—C3	0.9 (7)	N3—C11—C12—N4	11.9 (6)
C2—C3—C4—C5	-1.1 (6)	C10—C11—C12—N4	-167.7 (4)
C2—C3—C4—N2	-179.4 (4)	C12—N4—C13—C14	51.7 (6)
C1—N1—C5—C4	-0.7 (6)	C12—N4—C13—C17	-128.8 (4)
Hg1 ⁱ —N1—C5—C4	-169.4 (3)	C17—C13—C14—C15	-0.4 (6)
N2—C4—C5—N1	-179.9 (4)	N4—C13—C14—C15	179.2 (4)
C3—C4—C5—N1	1.8 (6)	C13—C14—C15—C16	0.5 (6)
C4—N2—C6—O1	-1.7 (8)	C17—N5—C16—C15	-1.0 (6)
C4—N2—C6—C7	179.3 (4)	Hg1—N5—C16—C15	170.4 (3)
C11—N3—C7—C8	0.4 (6)	C14—C15—C16—N5	0.2 (7)
C11—N3—C7—C6	-178.9 (4)	C16—N5—C17—C13	1.0 (6)
O1—C6—C7—N3	-177.7 (4)	Hg1—N5—C17—C13	-170.7 (3)
N2—C6—C7—N3	1.3 (6)	C14—C13—C17—N5	-0.3 (6)
O1—C6—C7—C8	3.0 (7)	N4—C13—C17—N5	-179.9 (4)
N2—C6—C7—C8	-178.0 (4)	C19—N6—C20—O3	2.1 (7)
N3—C7—C8—C9	-0.2 (7)	C18—N6—C20—O3	-179.2 (5)

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N4—H22 \cdots O3 ⁱ	0.859 (10)	2.08 (2)	2.891 (5)	157 (4)
N2—H21 \cdots O3 ⁱ	0.856 (10)	2.34 (2)	3.076 (5)	144 (3)
N2—H21 \cdots N3	0.856 (10)	2.25 (4)	2.685 (5)	111 (3)

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2




Fig. 3

