

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Triphenylbis(2,4,5-trifluoro-3-methoxybenzoato)antimony(V)

Liyuan Wen, Handong Yin,* Li Quan and Daqi Wang


 College of Chemistry and Chemical Engineering, Liaocheng University, Shandong 252059, People's Republic of China
 Correspondence e-mail: handongyin@163.com

Received 11 September 2008; accepted 16 September 2008

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å; R factor = 0.031; wR factor = 0.093; data-to-parameter ratio = 13.0.

In the title compound, $[\text{Sb}(\text{C}_6\text{H}_5)_3(\text{C}_8\text{H}_4\text{F}_3\text{O}_3)_2]$, the Sb atom lies on an inversion centre and exhibits a trigonal bipyramidal geometry with the axial positions occupied by the O atoms of two carboxylate groups and the equatorial positions occupied by C atoms of the phenyl groups. Intramolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds stabilize the molecular conformation. In the crystal structure, molecules are connected by intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen-bonding interactions, forming a layer structure parallel to $(\bar{2}01)$.

Related literature

 For related structures, see: Ferguson *et al.* (1987); Ruether *et al.* (1985).


Experimental

Crystal data

 $[\text{Sb}(\text{C}_6\text{H}_5)_3(\text{C}_8\text{H}_4\text{F}_3\text{O}_3)_2]$
 $M_r = 763.27$

 Monoclinic, $C2/c$
 $a = 12.7970$ (14) Å

 $b = 22.890$ (2) Å
 $c = 12.5131$ (10) Å
 $\beta = 120.107$ (2)°
 $V = 3170.9$ (5) Å³
 $Z = 4$

 Mo $K\alpha$ radiation
 $\mu = 0.95$ mm⁻¹
 $T = 293$ (2) K
 $0.50 \times 0.40 \times 0.35$ mm

Data collection

 Bruker SMART area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.635$, $T_{\max} = 0.718$

 7869 measured reflections
 2791 independent reflections
 2338 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.031$
 $wR(F^2) = 0.092$
 $S = 1.01$
 2791 reflections

 215 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.98$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.37$ e Å⁻³

Table 1
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C10}-\text{H10}\cdots\text{O2}$	0.93	2.36	3.053 (5)	131
$\text{C16}-\text{H16}\cdots\text{O1}$	0.93	2.49	2.979 (5)	113
$\text{C8}-\text{H8B}\cdots\text{O3}^i$	0.96	2.57	3.240 (7)	127
$\text{C11}-\text{H11}\cdots\text{O2}^{ii}$	0.93	2.51	3.255 (5)	138

 Symmetry codes: (i) $-x, -y + 1, -z + 2$; (ii) $x + \frac{1}{2}, -y + \frac{3}{2}, z + \frac{1}{2}$.

Data collection: SMART (Siemens, 1996); cell refinement: SAINT (Siemens, 1996); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

We acknowledge the National Natural Foundation of China (grant No. 20771053) and the Natural Science Foundation of Shandong Province (2005ZX09) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2246).

References

- Ferguson, G., Glidewell, C., Kaitner, B., Lloyd, D. & Metcalfe, S. (1987). *Acta Cryst.* **C43**, 824–826.
 Ruether, R., Huber, F. & Preut, H. (1985). *J. Organomet. Chem.* **295**, 21–28.
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Siemens (1996). *SMART and SAINT*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

supplementary materials

Acta Cryst. (2008). E64, m1303 [doi:10.1107/S1600536808029656]

Triphenylbis(2,4,5-trifluoro-3-methoxybenzoato)antimony(V)

L. Wen, H. Yin, L. Quan and D. Wang

Comment

In recent years organoantimony(V) derivatives have attracted considerable attention due to their significant antimicrobial properties as well as antitumor activities. We have therefore synthesized the title compound, and present its crystal structure here.

The molecular structure of the compound is shown in Fig.1 The Sb atom, which lies on an inversion centre, assumes a distorted trigonal bipyramidal coordination geometry, provided by three phenyl groups at the equatorial positions and two carboxylate groups at the axial positions. The Sb—O bond lengths in organoantimony compounds are extremely variable, ranging from 1.935 Å in triphenylstibine oxide (Ferguson *et al.* 1987) to 2.506 Å in tetraphenylstibonium benzenesulphonate hydrate (Ruether *et al.* 1985). The Sb1—O1 distance (2.132 (2) Å) in the title compound lies within this range. The Sb—C bond distances (Sb1—C9 = 2.122 (3) Å; Sb1—C15 = 2.118 (5) Å) fall in the normal range for Sb—C(phenyl) bonds (2.10–2.13 Å). The molecular conformation is stabilized by C—H···O hydrogen bonds. In the crystal packing, molecules are linked by intermolecular C—H···O hydrogen bonds (Fig.2, Table 1.), into layers parallel to the (-2 0 1) plane.

Experimental

The reaction was carried out under nitrogen atmosphere. 3-Methoxy-2,4,5-trifluorobenzoic acid (2 mmol) and sodium ethoxide (2.4 mmol) were added to a stirred solution of methanol (30 ml) in a Schlenk flask and stirred for 0.5 h. Triphenylantimony dichloride (1 mmol) was then added to the reactor and the reaction mixture was stirred for 12 h at room temperature. The resulting clear solution was evaporated under vacuum. The product was crystallized from a dichloromethane/methanol (1:1 v/v) solution to yield colourless blocks of the title compound (yield 86%. m.p. 458 K). Anal. Calcd (%) for C₃₄H₂₃O₆SbF₆ (Mr = 763.27): C, 53.50; H, 3.04; F, 14.93; Sb, 15.95. Found (%): C, 53.55; H, 3.07; F, 14.89; Sb, 15.99

Refinement

The H atoms were positioned geometrically, with methyl C—H distances of 0.96 Å and aromatic C—H distances of 0.93 Å, and refined as riding on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C}, \text{O})$ or $1.5 U_{\text{eq}}(\text{C})$ for the methyl group.

Figures




Fig. 1. The molecular structure of the compound, showing 50% probability displacement ellipsoids. H atoms have been omitted for clarity. Symmetry code: (A) = 1 - x, y, 3/2 - z.




Fig. 2. View of the two-dimensional layer structure in the title compound. Intermolecular hydrogen bonds are shown as dashed lines. H atoms are omitted.

Triphenylbis(2,4,5-trifluoro-3-methoxybenzoato)antimony(V)

Crystal data

[Sb(C₆H₅)₃(C₈H₄F₃O₃)₂]

$M_r = 763.27$

Monoclinic, *C2/c*

Hall symbol: -C 2yc

$a = 12.7970$ (14) Å

$b = 22.890$ (2) Å

$c = 12.5131$ (10) Å

$\beta = 120.107$ (2)°

$V = 3170.9$ (5) Å³

$Z = 4$

$F_{000} = 1520$

$D_x = 1.599$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 3589 reflections

$\theta = 2.6$ – 23.9 °

$\mu = 0.95$ mm⁻¹

$T = 293$ (2) K

Block, colourless

$0.50 \times 0.40 \times 0.35$ mm

Data collection

Bruker SMART area-detector
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293$ (2) K

φ and ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)

$T_{\min} = 0.635$, $T_{\max} = 0.718$

7869 measured reflections

2791 independent reflections

2338 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.036$

$\theta_{\max} = 25.0$ °

$\theta_{\min} = 2.0$ °

$h = -10 \rightarrow 15$

$k = -27 \rightarrow 23$

$l = -14 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.031$

$wR(F^2) = 0.093$

$S = 1.01$

2791 reflections

215 parameters

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.053P)^2 + 2.3063P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.98$ e Å⁻³

$\Delta\rho_{\min} = -0.37$ e Å⁻³

Primary atom site location: structure-invariant direct methods Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Sb1	0.5000	0.601220 (13)	0.7500	0.04156 (14)
F1	0.2452 (3)	0.54248 (11)	0.8929 (3)	0.0850 (8)
F2	-0.0110 (3)	0.65440 (19)	0.9796 (3)	0.1229 (13)
F3	0.0513 (3)	0.75229 (15)	0.9055 (3)	0.1104 (11)
O1	0.3684 (2)	0.59621 (10)	0.8065 (2)	0.0501 (6)
O2	0.3552 (2)	0.69304 (12)	0.7893 (3)	0.0614 (7)
O3	0.0880 (4)	0.54671 (19)	0.9730 (4)	0.1077 (13)
C1	0.3247 (3)	0.64637 (17)	0.8139 (3)	0.0501 (9)
C2	0.2343 (3)	0.64523 (17)	0.8571 (3)	0.0515 (9)
C3	0.2004 (4)	0.59594 (19)	0.8951 (4)	0.0608 (11)
C4	0.1193 (4)	0.5976 (2)	0.9386 (5)	0.0745 (14)
C5	0.0708 (4)	0.6507 (3)	0.9410 (5)	0.0808 (15)
C6	0.1022 (4)	0.7004 (3)	0.9027 (5)	0.0789 (14)
C7	0.1836 (3)	0.6985 (2)	0.8621 (4)	0.0612 (10)
H7	0.2052	0.7327	0.8380	0.073*
C8	0.1418 (6)	0.5380 (3)	1.1036 (6)	0.121 (2)
H8A	0.2277	0.5425	1.1423	0.181*
H8B	0.1233	0.4994	1.1192	0.181*
H8C	0.1102	0.5663	1.1367	0.181*
C9	0.6319 (3)	0.63273 (16)	0.9264 (3)	0.0455 (8)
C10	0.6261 (4)	0.68703 (18)	0.9726 (4)	0.0609 (10)
H10	0.5623	0.7123	0.9257	0.073*
C11	0.7170 (4)	0.7030 (2)	1.0897 (4)	0.0752 (13)
H11	0.7134	0.7390	1.1222	0.090*
C12	0.8122 (5)	0.6661 (3)	1.1582 (4)	0.0837 (15)
H12	0.8736	0.6775	1.2360	0.100*
C13	0.8168 (4)	0.6123 (3)	1.1118 (5)	0.0858 (15)
H13	0.8805	0.5870	1.1589	0.103*
C14	0.7263 (4)	0.59551 (19)	0.9947 (4)	0.0675 (12)
H14	0.7298	0.5593	0.9629	0.081*
C15	0.5000	0.5087 (2)	0.7500	0.0441 (11)
C16	0.4754 (4)	0.47799 (17)	0.8308 (4)	0.0601 (10)
H16	0.4585	0.4981	0.8849	0.072*
C17	0.4761 (5)	0.4175 (2)	0.8306 (5)	0.0791 (14)
H17	0.4603	0.3972	0.8853	0.095*
C18	0.5000	0.3873 (3)	0.7500	0.085 (2)
H18	0.5000	0.3466	0.7500	0.102*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sb1	0.0500 (2)	0.0384 (2)	0.0411 (2)	0.000	0.02652 (16)	0.000
F1	0.107 (2)	0.0643 (16)	0.114 (2)	-0.0008 (14)	0.0786 (19)	0.0110 (15)

supplementary materials

F2	0.092 (2)	0.195 (4)	0.124 (3)	0.005 (2)	0.086 (2)	-0.008 (3)
F3	0.105 (2)	0.126 (3)	0.121 (3)	0.0416 (19)	0.072 (2)	-0.008 (2)
O1	0.0571 (15)	0.0497 (15)	0.0566 (16)	0.0085 (11)	0.0384 (13)	0.0076 (12)
O2	0.0731 (17)	0.0541 (16)	0.0691 (19)	0.0055 (14)	0.0447 (15)	0.0113 (14)
O3	0.117 (3)	0.136 (3)	0.097 (3)	-0.050 (3)	0.074 (2)	-0.004 (3)
C1	0.048 (2)	0.061 (2)	0.042 (2)	0.0083 (17)	0.0229 (17)	0.0064 (17)
C2	0.0451 (19)	0.065 (2)	0.047 (2)	0.0059 (17)	0.0252 (17)	0.0075 (18)
C3	0.058 (2)	0.073 (3)	0.058 (3)	0.001 (2)	0.034 (2)	0.000 (2)
C4	0.064 (3)	0.105 (4)	0.066 (3)	-0.019 (3)	0.041 (2)	-0.001 (3)
C5	0.060 (3)	0.128 (5)	0.070 (3)	0.004 (3)	0.044 (2)	-0.005 (3)
C6	0.064 (3)	0.105 (4)	0.073 (3)	0.019 (3)	0.038 (2)	-0.010 (3)
C7	0.057 (2)	0.071 (3)	0.059 (2)	0.007 (2)	0.031 (2)	0.004 (2)
C8	0.121 (5)	0.135 (5)	0.110 (5)	-0.011 (4)	0.061 (4)	0.011 (5)
C9	0.052 (2)	0.047 (2)	0.041 (2)	-0.0099 (16)	0.0265 (16)	-0.0040 (16)
C10	0.072 (3)	0.055 (2)	0.060 (3)	-0.012 (2)	0.036 (2)	-0.010 (2)
C11	0.088 (3)	0.080 (3)	0.059 (3)	-0.033 (3)	0.039 (3)	-0.023 (2)
C12	0.077 (3)	0.116 (4)	0.051 (3)	-0.039 (3)	0.027 (2)	-0.027 (3)
C13	0.066 (3)	0.113 (4)	0.058 (3)	0.007 (3)	0.016 (2)	0.006 (3)
C14	0.062 (3)	0.077 (3)	0.052 (3)	0.003 (2)	0.020 (2)	-0.008 (2)
C15	0.048 (3)	0.043 (3)	0.040 (3)	0.000	0.022 (2)	0.000
C16	0.082 (3)	0.052 (2)	0.056 (2)	0.000 (2)	0.042 (2)	0.0034 (19)
C17	0.108 (4)	0.054 (3)	0.085 (4)	-0.005 (3)	0.056 (3)	0.013 (2)
C18	0.116 (6)	0.040 (3)	0.102 (6)	0.000	0.056 (5)	0.000

Geometric parameters (Å, °)

Sb1—C15	2.118 (5)	C8—H8B	0.9600
Sb1—C9	2.122 (3)	C8—H8C	0.9600
Sb1—C9 ⁱ	2.122 (3)	C9—C14	1.371 (5)
Sb1—O1 ⁱ	2.132 (2)	C9—C10	1.388 (5)
Sb1—O1	2.132 (2)	C10—C11	1.387 (6)
F1—C3	1.358 (5)	C10—H10	0.9300
F2—C5	1.358 (5)	C11—C12	1.372 (7)
F3—C6	1.364 (6)	C11—H11	0.9300
O1—C1	1.301 (4)	C12—C13	1.376 (7)
O2—C1	1.228 (4)	C12—H12	0.9300
O3—C4	1.370 (6)	C13—C14	1.391 (6)
O3—C8	1.434 (7)	C13—H13	0.9300
C1—C2	1.504 (5)	C14—H14	0.9300
C2—C3	1.376 (5)	C15—C16 ⁱ	1.390 (4)
C2—C7	1.398 (5)	C15—C16	1.390 (5)
C3—C4	1.393 (6)	C16—C17	1.384 (6)
C4—C5	1.371 (7)	C16—H16	0.9300
C5—C6	1.370 (8)	C17—C18	1.378 (6)
C6—C7	1.369 (6)	C17—H17	0.9300
C7—H7	0.9300	C18—C17 ⁱ	1.378 (6)
C8—H8A	0.9600	C18—H18	0.9300
C15—Sb1—C9	109.87 (10)	H8A—C8—H8B	109.5

C15—Sb1—C9 ⁱ	109.87 (10)	O3—C8—H8C	109.5
C9—Sb1—C9 ⁱ	140.3 (2)	H8A—C8—H8C	109.5
C15—Sb1—O1 ⁱ	86.91 (6)	H8B—C8—H8C	109.5
C9—Sb1—O1 ⁱ	90.84 (12)	C14—C9—C10	120.7 (4)
C9 ⁱ —Sb1—O1 ⁱ	91.25 (12)	C14—C9—Sb1	115.4 (3)
C15—Sb1—O1	86.91 (6)	C10—C9—Sb1	123.8 (3)
C9—Sb1—O1	91.25 (12)	C11—C10—C9	118.9 (4)
C9 ⁱ —Sb1—O1	90.84 (12)	C11—C10—H10	120.5
O1 ⁱ —Sb1—O1	173.83 (12)	C9—C10—H10	120.5
C1—O1—Sb1	114.6 (2)	C12—C11—C10	120.7 (4)
C4—O3—C8	115.3 (5)	C12—C11—H11	119.7
O2—C1—O1	123.2 (3)	C10—C11—H11	119.7
O2—C1—C2	120.3 (3)	C11—C12—C13	119.9 (4)
O1—C1—C2	116.5 (3)	C11—C12—H12	120.0
C3—C2—C7	117.8 (3)	C13—C12—H12	120.0
C3—C2—C1	124.9 (4)	C12—C13—C14	120.2 (5)
C7—C2—C1	117.3 (3)	C12—C13—H13	119.9
F1—C3—C2	121.5 (3)	C14—C13—H13	119.9
F1—C3—C4	116.0 (4)	C9—C14—C13	119.5 (4)
C2—C3—C4	122.5 (4)	C9—C14—H14	120.2
O3—C4—C5	122.7 (5)	C13—C14—H14	120.2
O3—C4—C3	119.5 (5)	C16 ⁱ —C15—C16	119.3 (5)
C5—C4—C3	117.8 (4)	C16 ⁱ —C15—Sb1	120.3 (2)
F2—C5—C6	119.1 (5)	C16—C15—Sb1	120.3 (2)
F2—C5—C4	120.0 (5)	C17—C16—C15	120.0 (4)
C6—C5—C4	120.9 (4)	C17—C16—H16	120.0
F3—C6—C7	119.9 (5)	C15—C16—H16	120.0
F3—C6—C5	119.1 (4)	C18—C17—C16	120.5 (5)
C7—C6—C5	121.0 (5)	C18—C17—H17	119.7
C6—C7—C2	120.0 (4)	C16—C17—H17	119.7
C6—C7—H7	120.0	C17—C18—C17 ⁱ	119.6 (6)
C2—C7—H7	120.0	C17—C18—H18	120.2
O3—C8—H8A	109.5	C17 ⁱ —C18—H18	120.2
O3—C8—H8B	109.5		
C15—Sb1—O1—C1	-175.3 (2)	C1—C2—C7—C6	-178.9 (4)
C9—Sb1—O1—C1	74.9 (3)	C15—Sb1—C9—C14	25.4 (3)
C9 ⁱ —Sb1—O1—C1	-65.5 (3)	C9 ⁱ —Sb1—C9—C14	-154.6 (3)
Sb1—O1—C1—O2	0.8 (5)	O1 ⁱ —Sb1—C9—C14	-61.7 (3)
Sb1—O1—C1—C2	-178.2 (2)	O1—Sb1—C9—C14	112.5 (3)
O2—C1—C2—C3	-175.3 (4)	C15—Sb1—C9—C10	-154.6 (3)
O1—C1—C2—C3	3.8 (6)	C9 ⁱ —Sb1—C9—C10	25.4 (3)
O2—C1—C2—C7	2.9 (5)	O1 ⁱ —Sb1—C9—C10	118.3 (3)
O1—C1—C2—C7	-178.1 (3)	O1—Sb1—C9—C10	-67.5 (3)
C7—C2—C3—F1	179.9 (4)	C14—C9—C10—C11	-0.5 (6)
C1—C2—C3—F1	-2.0 (6)	Sb1—C9—C10—C11	179.5 (3)
C7—C2—C3—C4	-0.5 (6)	C9—C10—C11—C12	1.0 (7)

supplementary materials

C1—C2—C3—C4	177.6 (4)	C10—C11—C12—C13	-1.4 (7)
C8—O3—C4—C5	-77.2 (7)	C11—C12—C13—C14	1.2 (8)
C8—O3—C4—C3	105.1 (6)	C10—C9—C14—C13	0.4 (7)
F1—C3—C4—O3	-1.5 (7)	Sb1—C9—C14—C13	-179.6 (4)
C2—C3—C4—O3	178.9 (4)	C12—C13—C14—C9	-0.7 (8)
F1—C3—C4—C5	-179.3 (4)	C9—Sb1—C15—C16 ⁱ	-119.9 (2)
C2—C3—C4—C5	1.1 (7)	C9 ⁱ —Sb1—C15—C16 ⁱ	60.1 (2)
O3—C4—C5—F2	0.6 (8)	O1 ⁱ —Sb1—C15—C16 ⁱ	-30.1 (2)
C3—C4—C5—F2	178.3 (4)	O1—Sb1—C15—C16 ⁱ	149.9 (2)
O3—C4—C5—C6	-178.2 (5)	C9—Sb1—C15—C16	60.1 (2)
C3—C4—C5—C6	-0.5 (8)	C9 ⁱ —Sb1—C15—C16	-119.9 (2)
F2—C5—C6—F3	0.6 (7)	O1 ⁱ —Sb1—C15—C16	149.9 (2)
C4—C5—C6—F3	179.5 (5)	O1—Sb1—C15—C16	-30.1 (2)
F2—C5—C6—C7	-179.5 (4)	C16 ⁱ —C15—C16—C17	0.3 (3)
C4—C5—C6—C7	-0.7 (8)	Sb1—C15—C16—C17	-179.7 (3)
F3—C6—C7—C2	-178.9 (4)	C15—C16—C17—C18	-0.6 (7)
C5—C6—C7—C2	1.3 (7)	C16—C17—C18—C17 ⁱ	0.3 (3)
C3—C2—C7—C6	-0.7 (6)		

Symmetry codes: (i) $-x+1, y, -z+3/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C10—H10 \cdots O2	0.93	2.36	3.053 (5)	131
C16—H16 \cdots O1	0.93	2.49	2.979 (5)	113
C8—H8B \cdots O3 ⁱⁱ	0.96	2.57	3.240 (7)	127
C11—H11 \cdots O2 ⁱⁱⁱ	0.93	2.51	3.255 (5)	138

Symmetry codes: (ii) $-x, -y+1, -z+2$; (iii) $x+1/2, -y+3/2, z+1/2$.

Fig. 2

