

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# 1-(2,4-Dichlorophenyl)-4,4-dimethylpent-1-en-3-one

Lin Xia and Ai-Xi Hu\*


College of Chemistry and Chemical Engineering, Hunan University, 410082 Changsha, People's Republic of China  
Correspondence e-mail: axhu0731@yahoo.com.cn

Received 28 August 2008; accepted 17 September 2008

Key indicators: single-crystal X-ray study;  $T = 173$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.032;  $wR$  factor = 0.085; data-to-parameter ratio = 15.2.

In the title compound,  $\text{C}_{13}\text{H}_{14}\text{Cl}_2\text{O}$ , the carbonyl and ethenyl groups are coplanar with the aromatic ring. There are four molecules in the asymmetric unit and all atoms in the molecule lie on mirror planes. The molecules are packed in an offset face-to-face arrangement showing  $\pi$ - $\pi$  stacking interactions involving the benzene rings [centroid-centroid distance = 3.564 (2) Å].

## Related literature

 For related compounds, see: Wang *et al.* (2008).


## Experimental

## Crystal data

$\text{C}_{13}\text{H}_{14}\text{Cl}_2\text{O}$   
 $M_r = 257.14$   
Orthorhombic,  $Pnma$   
 $a = 11.2553$  (7) Å  
 $b = 7.0458$  (4) Å  
 $c = 15.4969$  (9) Å  
 $V = 1228.94$  (13) Å<sup>3</sup>  
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.50$  mm<sup>-1</sup>  
 $T = 173$  (2) K  
 $0.47 \times 0.39 \times 0.15$  mm

## Data collection

Bruker SMART 1000 CCD diffractometer  
Absorption correction: multi-scan (SADABS; Sheldrick, 2004)  
 $T_{\min} = 0.798$ ,  $T_{\max} = 0.928$   
6162 measured reflections  
1444 independent reflections  
1227 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.021$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$   
 $wR(F^2) = 0.085$   
 $S = 1.07$   
1444 reflections  
95 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.28$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.36$  e Å<sup>-3</sup>

Data collection: SMART (Bruker, 2001); cell refinement: SAINT-Plus (Bruker, 2003); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL.

The authors express their thanks to the National Key Technology R&D Program (No. 2006BAE01A01-4)

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SG2260).

## References

- Bruker (2001). SMART. Bruker AXS Inc., Madison, Wisconsin, USA.  
Bruker (2003). SAINT-Plus. Bruker AXS Inc., Madison, Wisconsin, USA.  
Sheldrick, G. M. (2004). SADABS. University of Göttingen, Germany.  
Sheldrick, G. M. (2008). Acta Cryst. **A64**, 112–122.  
Wang, Y., Hu, A.-X., Cao, G., Li, G.-X., Zhang, J.-Y., Xia, L., Ou, X.-M. & Xu, J.-B. (2008). Chin. J. Org. Chem. **28**, 443–448.

**supplementary materials**

*Acta Cryst.* (2008). E64, o1983 [ doi:10.1107/S1600536808029838 ]

## 1-(2,4-Dichlorophenyl)-4,4-dimethylpent-1-en-3-one

L. Xia and A.-X. Hu

### Comment

Phenyl-4,4-dimethylpentan-3-one derivatives are important intermediates in medicinal industry (Wang *et al.*, 2006). Herein we report the synthesis and crystal structure of 1-(2,4-dichlorophen-yl)-4,4-dimethylpentan-3-one.

### Experimental

3,3-dimethylbutan-2-one(0.0105 mol) was added dropwise into a solution of 2,4-dichlorobenzaldehyde (0.01 mol) and 60 ml ethanol. Then 0.1 g 50% NaOH solution as catalyst was added and the reaction was stirred at 333 K for 5 h . The solvent was evaporated to about half volume, then cooled to 277 K and the precipitate formed.This was filtered and dried to give the desired product (yield: 94.7%). Crystals suitable for X-ray structure determination were obtained by slow evaporation of an ethanol solution at room temperature.

### Refinement

The methyl H atoms were positioned geometrically (C—H=0.98 Å) and torsion angles refined to fit the electron density [ $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ ]. Other H atoms were placed in calculated position (methylene C—H = 0.95Å and aromatic C—H=0.95 Å) and refined as riding [ $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ ].

### Figures




Fig. 1. Molecular structure showing 30% probability displacement ellipsoids.H atoms are omitted for clarity.




Fig. 2. Packing diagram showing  $\pi$ - $\pi$  stacking interactions.

## 1-(2,4-Dichlorophenyl)-4,4-dimethylpent-1-en-3-one

### Crystal data

$\text{C}_{13}\text{H}_{14}\text{Cl}_2\text{O}$

$M_r = 257.14$

Orthorhombic, *Pnma*

Hall symbol: -P 2ac 2n

$D_x = 1.390 \text{ Mg m}^{-3}$

Melting point: 385 K

Mo  $K\alpha$  radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 3552 reflections

# supplementary materials

---

$a = 11.2553 (7) \text{ \AA}$	$\theta = 2.2\text{--}27.0^\circ$
$b = 7.0458 (4) \text{ \AA}$	$\mu = 0.50 \text{ mm}^{-1}$
$c = 15.4969 (9) \text{ \AA}$	$T = 173 (2) \text{ K}$
$V = 1228.94 (13) \text{ \AA}^3$	Block, colorless
$Z = 4$	$0.47 \times 0.39 \times 0.15 \text{ mm}$
$F_{000} = 536$	

## Data collection

Bruker SMART 1000 CCD diffractometer	1444 independent reflections
Radiation source: fine-focus sealed tube	1227 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.021$
$T = 173(2) \text{ K}$	$\theta_{\text{max}} = 27.0^\circ$
$\omega$ scans	$\theta_{\text{min}} = 2.2^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 2004)	$h = -12 \rightarrow 14$
$T_{\text{min}} = 0.798, T_{\text{max}} = 0.928$	$k = -8 \rightarrow 9$
6162 measured reflections	$l = -19 \rightarrow 14$

## Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.032$	H-atom parameters constrained
$wR(F^2) = 0.085$	$w = 1/[\sigma^2(F_o^2) + (0.0391P)^2 + 0.6612P]$
$S = 1.07$	where $P = (F_o^2 + 2F_c^2)/3$
1444 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
95 parameters	$\Delta\rho_{\text{max}} = 0.28 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.36 \text{ e \AA}^{-3}$
	Extinction correction: none

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C11	0.73794 (5)	0.2500	0.45333 (4)	0.03276 (17)	
C12	0.34147 (5)	0.2500	0.26347 (3)	0.03138 (17)	
C1	0.57275 (19)	0.2500	0.61286 (13)	0.0239 (4)	
H1	0.6571	0.2500	0.6149	0.029*	
C2	0.51505 (19)	0.2500	0.68766 (13)	0.0265 (5)	
H2	0.4307	0.2500	0.6869	0.032*	
C3	0.57686 (18)	0.2500	0.77240 (13)	0.0225 (4)	
C4	0.49783 (18)	0.2500	0.85260 (12)	0.0227 (4)	
C5	0.5740 (2)	0.2500	0.93380 (14)	0.0314 (5)	
H5A	0.6163	0.1518	0.9347	0.047*	0.50
H5B	0.5225	0.2500	0.9848	0.047*	
H5C	0.6252	0.3625	0.9340	0.047*	0.50
C6	0.41949 (13)	0.0709 (2)	0.85096 (10)	0.0285 (3)	
H6A	0.3670	0.0751	0.8005	0.043*	
H6B	0.3714	0.0656	0.9036	0.043*	
H6C	0.4701	-0.0421	0.8477	0.043*	
C7	0.51674 (18)	0.2500	0.52726 (13)	0.0224 (4)	
C8	0.58317 (17)	0.2500	0.45064 (13)	0.0217 (4)	
C9	0.53111 (19)	0.2500	0.36945 (13)	0.0244 (4)	
H9	0.5785	0.2500	0.3187	0.029*	
C10	0.40892 (19)	0.2500	0.36426 (13)	0.0241 (4)	
C11	0.33861 (18)	0.2500	0.43745 (15)	0.0267 (5)	
H11	0.2544	0.2500	0.4329	0.032*	
C12	0.39335 (19)	0.2500	0.51767 (14)	0.0265 (5)	
H12	0.3451	0.2500	0.5680	0.032*	
O1	0.68512 (13)	0.2500	0.77742 (10)	0.0323 (4)	

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C11	0.0204 (3)	0.0528 (4)	0.0251 (3)	0.000	0.00297 (19)	0.000
C12	0.0291 (3)	0.0413 (3)	0.0237 (3)	0.000	-0.0039 (2)	0.000
C1	0.0243 (10)	0.0245 (10)	0.0228 (10)	0.000	0.0018 (8)	0.000
C2	0.0232 (10)	0.0337 (12)	0.0225 (10)	0.000	-0.0004 (8)	0.000
C3	0.0221 (10)	0.0236 (10)	0.0218 (10)	0.000	0.0009 (8)	0.000
C4	0.0202 (9)	0.0284 (11)	0.0195 (9)	0.000	0.0007 (8)	0.000
C5	0.0256 (11)	0.0463 (14)	0.0223 (10)	0.000	-0.0029 (9)	0.000
C6	0.0269 (7)	0.0303 (8)	0.0285 (8)	-0.0032 (6)	0.0043 (6)	0.0008 (6)
C7	0.0238 (10)	0.0204 (10)	0.0229 (10)	0.000	0.0011 (8)	0.000
C8	0.0190 (9)	0.0209 (10)	0.0252 (10)	0.000	0.0019 (8)	0.000
C9	0.0276 (11)	0.0240 (10)	0.0217 (10)	0.000	0.0039 (8)	0.000
C10	0.0286 (11)	0.0220 (10)	0.0217 (10)	0.000	-0.0019 (8)	0.000
C11	0.0205 (10)	0.0311 (12)	0.0285 (11)	0.000	0.0005 (8)	0.000
C12	0.0256 (11)	0.0310 (11)	0.0228 (10)	0.000	0.0055 (8)	0.000

# supplementary materials

O1                    0.0200 (7)            0.0468 (10)            0.0301 (8)            0.000                    0.0026 (6)            0.000

## *Geometric parameters (Å, °)*

C11—C8	1.742 (2)	C5—H5C	0.9800
C12—C10	1.737 (2)	C6—H6A	0.9800
C1—C2	1.329 (3)	C6—H6B	0.9800
C1—C7	1.469 (3)	C6—H6C	0.9800
C1—H1	0.9500	C7—C12	1.397 (3)
C2—C3	1.486 (3)	C7—C8	1.403 (3)
C2—H2	0.9500	C8—C9	1.388 (3)
C3—O1	1.221 (2)	C9—C10	1.378 (3)
C3—C4	1.528 (3)	C9—H9	0.9500
C4—C5	1.522 (3)	C10—C11	1.383 (3)
C4—C6	1.5398 (19)	C11—C12	1.387 (3)
C4—C6 <sup>i</sup>	1.5398 (19)	C11—H11	0.9500
C5—H5A	0.8400	C12—H12	0.9500
C5—H5B	0.9799		
C2—C1—C7	125.3 (2)	H6A—C6—H6B	109.5
C2—C1—H1	117.3	C4—C6—H6C	109.5
C7—C1—H1	117.3	H6A—C6—H6C	109.5
C1—C2—C3	122.83 (19)	H6B—C6—H6C	109.5
C1—C2—H2	118.6	C12—C7—C8	116.09 (18)
C3—C2—H2	118.6	C12—C7—C1	121.53 (19)
O1—C3—C2	121.57 (19)	C8—C7—C1	122.39 (18)
O1—C3—C4	121.93 (18)	C9—C8—C7	122.84 (19)
C2—C3—C4	116.50 (17)	C9—C8—C11	116.34 (15)
C5—C4—C3	110.15 (17)	C7—C8—C11	120.82 (15)
C5—C4—C6	109.64 (11)	C10—C9—C8	118.32 (19)
C3—C4—C6	108.65 (11)	C10—C9—H9	120.8
C5—C4—C6 <sup>i</sup>	109.64 (11)	C8—C9—H9	120.8
C3—C4—C6 <sup>i</sup>	108.65 (11)	C9—C10—C11	121.55 (19)
C6—C4—C6 <sup>i</sup>	110.09 (17)	C9—C10—C12	119.27 (16)
C4—C5—H5A	109.5	C11—C10—C12	119.17 (16)
C4—C5—H5B	109.5	C10—C11—C12	118.73 (19)
H5A—C5—H5B	108.7	C10—C11—H11	120.6
C4—C5—H5C	109.4	C12—C11—H11	120.6
H5A—C5—H5C	109.5	C11—C12—C7	122.5 (2)
H5B—C5—H5C	110.2	C11—C12—H12	118.8
C4—C6—H6A	109.5	C7—C12—H12	118.8
C4—C6—H6B	109.5		
C7—C1—C2—C3	180.0	C1—C7—C8—C9	180.0
C1—C2—C3—O1	0.0	C12—C7—C8—C11	180.0
C1—C2—C3—C4	180.0	C1—C7—C8—C11	0.0
O1—C3—C4—C5	0.0	C7—C8—C9—C10	0.0
C2—C3—C4—C5	180.0	C11—C8—C9—C10	180.0
O1—C3—C4—C6	-120.11 (11)	C8—C9—C10—C11	0.0
C2—C3—C4—C6	59.89 (11)	C8—C9—C10—C12	180.0

## supplementary materials

---

O1—C3—C4—C6 <sup>i</sup>	120.11 (11)	C9—C10—C11—C12	0.0
C2—C3—C4—C6 <sup>i</sup>	-59.89 (11)	C12—C10—C11—C12	180.0
C2—C1—C7—C12	0.0	C10—C11—C12—C7	0.0
C2—C1—C7—C8	180.0	C8—C7—C12—C11	0.0
C12—C7—C8—C9	0.0	C1—C7—C12—C11	180.0

Symmetry codes: (i)  $x, -y+1/2, z$ .

Fig. 1




Fig. 2

