

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-Formyl-3-hydroxy-9,10-anthroquinone

Nor Hadiani Ismail,^a Che Puteh Osman,^a Khalijah Awang,^b Sri Nurestri Abdul Malek^c and Seik Weng Ng^{b*}

^aFaculty of Applied Sciences, Universiti Teknologi MARA, 40450 Shah Alam, Malaysia, ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^cInstitute of Biological Sciences, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 23 September 2008; accepted 6 October 2008

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.087; wR factor = 0.343; data-to-parameter ratio = 14.0.

The molecule of the title compound, $\text{C}_{15}\text{H}_8\text{O}_4$, is approximately planar. An intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond is observed between the hydroxy and formyl groups. The crystal used was a nonmerohedral twin, with a minor twin component of 15.9%.

Related literature

For antileishmanial and antiplasmodial activities, see: Sittie *et al.* (1999). For the treatment of twinned diffraction data, see: Spek (2003).



Experimental

Crystal data

$\text{C}_{15}\text{H}_8\text{O}_4$
 $M_r = 252.21$
 Triclinic, $P\bar{1}$
 $a = 6.9194$ (2) Å
 $b = 8.0650$ (2) Å
 $c = 10.7601$ (3) Å
 $\alpha = 86.250$ (2)°
 $\beta = 83.214$ (2)°
 $\gamma = 64.692$ (2)°
 $V = 538.96$ (3) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.11$ mm⁻¹
 $T = 100$ (2) K
 $0.22 \times 0.04 \times 0.04$ mm

Data collection

Bruker SMART APEXII area-detector diffractometer
 Absorption correction: none
 4946 measured reflections
 2419 independent reflections
 1880 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.087$
 $wR(F^2) = 0.343$
 $S = 1.11$
 2419 reflections
 173 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.49$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.44$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O2}-\text{H2}\cdots\text{O1}$	0.84	2.00	2.635 (5)	132

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *pubCIF* (Westrip, 2008).

The authors thank the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2681).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Sittie, A. A., Lemnich, E., Olsen, C. E., Hviid, L., Kharazmi, A., Nkrumah, F. K. & Christensen, S. B. (1999). *Planta Med.* **65**, 259–261.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
 Westrip, S. P. (2008). *pubCIF*. In preparation.

supplementary materials

Acta Cryst. (2008). E64, o2164 [doi:10.1107/S1600536808032224]

2-Formyl-3-hydroxy-9,10-anthroquinone

N. H. Ismail, C. P. Osman, K. Awang, S. N. Abdul Malek and S. W. Ng

Experimental

Rennellia elliptica Korth from the Rubiaceae family was collected from Kuala Keniam, Pahang, Malaysia. The root was chopped into small pieces and dried. The dried sample (1 kg) was ground and then extracted successively with hexane, dichloromethane and methanol. The dichloromethane extract was concentrated *in vacuo* to give 27 g crude extract. The crude extract was fractionated by column chromatography. The column (60 cm \times 5 cm) was packed with acid-washed silica gel and eluted with hexane, dichloromethane and methanol. Nine fractions were obtained, and 3-hydroxy-2-formyl-9,10-anthraquinone (41.5 mg) was isolated from the third fraction (hexane:dichloromethane, 30:70) by slow evaporation of the solvent mixture. The yellow crystals obtained were washed with acetone.

Refinement

Carbon- and oxygen-bound H-atoms were placed in calculated positions (C—H = 0.95 Å and O—H = 0.84 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to $1.2\text{--}1.5U_{\text{eq}}(\text{C}, \text{O})$. The crystal studied was a non-merohedral twin. The *TwinRotMat* in *PLATON* (Spek, 2003) gave the twin law as $(-1\ 0\ 0, 0\ -1\ 0, -0.343\ -0.049\ 1)$, whose inclusion in the refinement lowered the R index from 11.3 to 8.7%. The twin component refined to 18.9%. The refinement is deemed satisfactory although the wR_2 value for all reflections is somewhat high. The structure has a long C5—C14 bond; as the anisotropic displacement parameters are normal, the likely reason is localization of the double bonds in the ring. On the other hand, the C13—C14 bond is somewhat short.

Figures




Fig. 1. The molecular structure of the title compound, showing 70% probability displacement ellipsoids. H atoms are drawn as spheres of arbitrary radii.

2-Formyl-3-hydroxy-9,10-anthroquinone

Crystal data

$\text{C}_{15}\text{H}_8\text{O}_4$

$M_r = 252.21$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 6.9194\ (2)\ \text{\AA}$

$b = 8.0650\ (2)\ \text{\AA}$

$Z = 2$

$F_{000} = 260$

$D_x = 1.554\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 1888 reflections

$\theta = 3.3\text{--}28.3^\circ$

supplementary materials

$c = 10.7601 (3) \text{ \AA}$
 $\alpha = 86.250 (2)^\circ$
 $\beta = 83.214 (2)^\circ$
 $\gamma = 64.692 (2)^\circ$
 $V = 538.96 (3) \text{ \AA}^3$

$\mu = 0.11 \text{ mm}^{-1}$
 $T = 100 (2) \text{ K}$
Block, yellow
 $0.22 \times 0.04 \times 0.04 \text{ mm}$

Data collection

Bruker SMART APEXII area-detector diffractometer
Radiation source: fine-focus sealed tube
Monochromator: graphite
 $T = 100(2) \text{ K}$
 ω scans
Absorption correction: none
4946 measured reflections
2419 independent reflections

1880 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$
 $\theta_{\text{max}} = 27.5^\circ$
 $\theta_{\text{min}} = 2.8^\circ$
 $h = -8 \rightarrow 8$
 $k = -10 \rightarrow 10$
 $l = -13 \rightarrow 13$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.087$
 $wR(F^2) = 0.343$
 $S = 1.11$
2419 reflections
173 parameters
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.1778P)^2 + 2.2381P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.49 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.44 \text{ e \AA}^{-3}$
Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.3155 (6)	0.6486 (5)	0.0888 (3)	0.0251 (8)
O2	0.2501 (6)	0.3493 (5)	0.0941 (3)	0.0242 (8)
H2	0.2611	0.4345	0.0490	0.029*
O3	0.2366 (5)	-0.0898 (4)	0.4383 (3)	0.0153 (7)
O4	0.2779 (5)	0.4648 (4)	0.6688 (3)	0.0172 (7)
C1	0.3060 (7)	0.6395 (6)	0.2041 (4)	0.0178 (9)
H1	0.3171	0.7336	0.2474	0.021*
C2	0.2786 (7)	0.4901 (6)	0.2772 (4)	0.0143 (9)
C3	0.2544 (7)	0.3500 (6)	0.2187 (4)	0.0161 (9)
C4	0.2356 (7)	0.2059 (6)	0.2894 (4)	0.0152 (9)
H4	0.2187	0.1116	0.2500	0.018*
C5	0.2419 (6)	0.2016 (5)	0.4183 (4)	0.0123 (8)

C6	0.2335 (6)	0.0406 (5)	0.4916 (4)	0.0112 (8)
C7	0.2285 (6)	0.0417 (5)	0.6299 (4)	0.0112 (8)
C8	0.2117 (7)	-0.1043 (6)	0.7003 (4)	0.0139 (8)
H8	0.1986	-0.1999	0.6601	0.017*
C9	0.2144 (7)	-0.1091 (6)	0.8286 (4)	0.0179 (9)
H9	0.2031	-0.2083	0.8766	0.021*
C10	0.2336 (7)	0.0310 (6)	0.8883 (4)	0.0188 (9)
H10	0.2353	0.0269	0.9767	0.023*
C11	0.2503 (7)	0.1762 (6)	0.8187 (4)	0.0173 (9)
H11	0.2645	0.2709	0.8594	0.021*
C12	0.2462 (6)	0.1835 (5)	0.6895 (4)	0.0113 (8)
C13	0.2641 (6)	0.3403 (6)	0.6165 (4)	0.0126 (8)
C14	0.2621 (6)	0.3426 (5)	0.4787 (4)	0.0120 (8)
C15	0.2815 (7)	0.4851 (6)	0.4072 (4)	0.0133 (8)
H15	0.2968	0.5801	0.4468	0.016*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.038 (2)	0.0258 (18)	0.0125 (16)	-0.0155 (16)	-0.0033 (13)	0.0054 (13)
O2	0.046 (2)	0.0283 (18)	0.0063 (15)	-0.0226 (17)	-0.0062 (13)	0.0020 (12)
O3	0.0194 (15)	0.0131 (14)	0.0147 (15)	-0.0079 (12)	-0.0028 (11)	-0.0009 (11)
O4	0.0228 (16)	0.0148 (15)	0.0165 (15)	-0.0103 (13)	-0.0003 (12)	-0.0039 (11)
C1	0.021 (2)	0.017 (2)	0.016 (2)	-0.0078 (17)	-0.0040 (16)	0.0035 (16)
C2	0.0146 (19)	0.0149 (19)	0.0119 (19)	-0.0053 (16)	-0.0011 (14)	0.0030 (15)
C3	0.017 (2)	0.021 (2)	0.0103 (19)	-0.0080 (17)	-0.0030 (15)	0.0010 (15)
C4	0.016 (2)	0.016 (2)	0.014 (2)	-0.0074 (16)	-0.0007 (15)	-0.0025 (15)
C5	0.0106 (18)	0.0107 (18)	0.0135 (19)	-0.0032 (14)	0.0010 (14)	0.0003 (14)
C6	0.0099 (17)	0.0105 (18)	0.0121 (19)	-0.0036 (14)	-0.0001 (14)	-0.0004 (14)
C7	0.0109 (18)	0.0114 (18)	0.0103 (18)	-0.0038 (14)	-0.0025 (13)	0.0011 (14)
C8	0.0145 (19)	0.0118 (18)	0.015 (2)	-0.0053 (15)	-0.0023 (15)	0.0013 (14)
C9	0.019 (2)	0.017 (2)	0.017 (2)	-0.0081 (17)	-0.0012 (16)	0.0055 (16)
C10	0.024 (2)	0.022 (2)	0.0110 (19)	-0.0100 (18)	-0.0009 (16)	0.0026 (16)
C11	0.021 (2)	0.018 (2)	0.013 (2)	-0.0084 (17)	-0.0008 (16)	-0.0009 (15)
C12	0.0108 (18)	0.0112 (18)	0.0109 (18)	-0.0039 (14)	-0.0002 (14)	-0.0005 (14)
C13	0.0112 (18)	0.0125 (19)	0.0139 (19)	-0.0046 (15)	-0.0015 (14)	-0.0010 (14)
C14	0.0116 (18)	0.0112 (18)	0.0127 (19)	-0.0044 (14)	0.0001 (14)	-0.0002 (14)
C15	0.0138 (19)	0.0112 (18)	0.014 (2)	-0.0049 (15)	0.0006 (14)	-0.0017 (14)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.234 (5)	C7—C8	1.398 (5)
O2—C3	1.345 (5)	C7—C12	1.403 (5)
O2—H2	0.84	C8—C9	1.380 (6)
O3—C6	1.222 (5)	C8—H8	0.95
O4—C13	1.226 (5)	C9—C10	1.397 (6)
C1—C2	1.464 (6)	C9—H9	0.95
C1—H1	0.95	C10—C11	1.388 (6)
C2—C15	1.400 (6)	C10—H10	0.95

supplementary materials

C2—C3	1.407 (6)	C11—C12	1.391 (6)
C3—C4	1.391 (6)	C11—H11	0.95
C4—C5	1.390 (6)	C12—C13	1.487 (5)
C4—H4	0.95	C13—C14	1.483 (6)
C5—C14	1.410 (6)	C14—C15	1.387 (6)
C5—C6	1.494 (5)	C15—H15	0.95
C6—C7	1.485 (5)		
C3—O2—H2	120.0	C9—C8—H8	120.1
O1—C1—C2	122.8 (4)	C7—C8—H8	120.1
O1—C1—H1	118.6	C8—C9—C10	120.4 (4)
C2—C1—H1	118.6	C8—C9—H9	119.8
C15—C2—C3	119.7 (4)	C10—C9—H9	119.8
C15—C2—C1	119.1 (4)	C11—C10—C9	120.0 (4)
C3—C2—C1	121.2 (4)	C11—C10—H10	120.0
O2—C3—C4	117.9 (4)	C9—C10—H10	120.0
O2—C3—C2	121.8 (4)	C10—C11—C12	120.2 (4)
C4—C3—C2	120.3 (4)	C10—C11—H11	119.9
C5—C4—C3	119.2 (4)	C12—C11—H11	119.9
C5—C4—H4	120.4	C11—C12—C7	119.6 (4)
C3—C4—H4	120.4	C11—C12—C13	119.4 (4)
C4—C5—C14	121.2 (4)	C7—C12—C13	121.1 (3)
C4—C5—C6	118.4 (4)	O4—C13—C14	121.0 (4)
C14—C5—C6	120.3 (4)	O4—C13—C12	121.0 (4)
O3—C6—C7	121.3 (4)	C14—C13—C12	118.0 (3)
O3—C6—C5	120.5 (4)	C15—C14—C5	119.0 (4)
C7—C6—C5	118.1 (3)	C15—C14—C13	119.7 (4)
C8—C7—C12	120.1 (4)	C5—C14—C13	121.3 (4)
C8—C7—C6	118.9 (4)	C14—C15—C2	120.5 (4)
C12—C7—C6	121.0 (3)	C14—C15—H15	119.8
C9—C8—C7	119.7 (4)	C2—C15—H15	119.8
O1—C1—C2—C15	176.5 (4)	C10—C11—C12—C7	-1.0 (6)
O1—C1—C2—C3	-2.1 (7)	C10—C11—C12—C13	180.0 (4)
C15—C2—C3—O2	179.8 (4)	C8—C7—C12—C11	1.0 (6)
C1—C2—C3—O2	-1.7 (7)	C6—C7—C12—C11	-177.2 (4)
C15—C2—C3—C4	-0.7 (7)	C8—C7—C12—C13	-180.0 (3)
C1—C2—C3—C4	177.9 (4)	C6—C7—C12—C13	1.8 (6)
O2—C3—C4—C5	179.3 (4)	C11—C12—C13—O4	-1.5 (6)
C2—C3—C4—C5	-0.3 (7)	C7—C12—C13—O4	179.4 (4)
C3—C4—C5—C14	1.4 (6)	C11—C12—C13—C14	179.2 (4)
C3—C4—C5—C6	-176.7 (4)	C7—C12—C13—C14	0.2 (6)
C4—C5—C6—O3	5.1 (6)	C4—C5—C14—C15	-1.6 (6)
C14—C5—C6—O3	-173.0 (4)	C6—C5—C14—C15	176.4 (3)
C4—C5—C6—C7	-176.8 (4)	C4—C5—C14—C13	178.7 (4)
C14—C5—C6—C7	5.1 (6)	C6—C5—C14—C13	-3.2 (6)
O3—C6—C7—C8	-4.6 (6)	O4—C13—C14—C15	1.6 (6)
C5—C6—C7—C8	177.4 (4)	C12—C13—C14—C15	-179.1 (4)
O3—C6—C7—C12	173.7 (4)	O4—C13—C14—C5	-178.7 (4)
C5—C6—C7—C12	-4.4 (6)	C12—C13—C14—C5	0.6 (6)

C12—C7—C8—C9	-0.5 (6)	C5—C14—C15—C2	0.7 (6)
C6—C7—C8—C9	177.7 (4)	C13—C14—C15—C2	-179.6 (4)
C7—C8—C9—C10	0.0 (7)	C3—C2—C15—C14	0.4 (6)
C8—C9—C10—C11	0.0 (7)	C1—C2—C15—C14	-178.1 (4)
C9—C10—C11—C12	0.5 (7)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O2—H2...O1	0.84	2.00	2.635 (5)	132

Fig. 1

