

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N,N'-Bis(2-methoxyphenyl)biphenyl-2,2'-dicarboxamide

Gui-Yu Wang, Di Li, Da-Bin Qin,* Jie-Wei Luo and Li-Hui Guo

School of Chemistry and Chemical Engineering, China West Normal University, Nanchong 637002, People's Republic of China
Correspondence e-mail: qindabincwnu@yahoo.com.cn


Received 30 June 2008; accepted 7 October 2008

Key indicators: single-crystal X-ray study; $T = 113$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.039; wR factor = 0.093; data-to-parameter ratio = 6.4.

In the title compound, $\text{C}_{28}\text{H}_{24}\text{N}_2\text{O}_4$, the dihedral angle between the two rings of the biphenyl unit is 75.34 (9)°. The outer aromatic rings form dihedral angles of 66.96 (1) and 85.69 (8)° with the rings to which they are attached. The molecular structure is stabilized by intramolecular $\text{C}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds. In the crystal structure, intermolecular $\text{N}-\text{H}\cdots\text{O}$ interactions are observed.

Related literature

For the synthesis, see: Gao & Gao (2002). For related structures, see: Wang & Han (2004); Wang & Jiang (2004); Huang & Yang (2008).



Experimental

Crystal data

$\text{C}_{28}\text{H}_{24}\text{N}_2\text{O}_4$
 $M_r = 452.49$
Monoclinic, Cc
 $a = 18.184$ (4) Å

$b = 16.304$ (3) Å
 $c = 7.9998$ (16) Å
 $\beta = 108.90$ (3)°
 $V = 2243.9$ (8) Å³

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹

$T = 113$ (2) K
 $0.16 \times 0.14 \times 0.10$ mm

Data collection

Rigaku Saturn CCD diffractometer
Absorption correction: multi-scan
(*CrystalStructure*; Rigaku/MS, 2004)
 $T_{\min} = 0.986$, $T_{\max} = 0.991$

6422 measured reflections
1991 independent reflections
1858 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.054$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.093$
 $S = 1.06$
1991 reflections
309 parameters

2 restraints
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.17$ e Å⁻³
 $\Delta\rho_{\min} = -0.20$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1N}\cdots\text{O3}^i$	0.86	2.02	2.833 (3)	157
$\text{N2}-\text{H2N}\cdots\text{O2}$	0.86	2.24	3.081 (4)	167
$\text{N2}-\text{H2N}\cdots\text{O4}$	0.86	2.24	2.612 (3)	106
$\text{C22}-\text{H22}\cdots\text{O3}$	0.93	2.30	2.885 (4)	120

Symmetry code: (i) $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *RAPID-AUTO* (Rigaku/MS, 2004); cell refinement: *RAPID-AUTO*; data reduction: *RAPID-AUTO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the Scientific Research Fund Projects of China West Normal University (grant No. 06B003) and the Youth Fund Projects of Sichuan Education Department (grant No. 2006B039).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GW2044).

References

- Gao, M. Z. & Gao, J. (2002). *Tetrahedron Lett.* **43**, 5001–5003.
Huang, W.-W. & Yang, S.-P. (2008). *Acta Cryst.* **E64**, m525–m526.
Rigaku/MS (2004). *RAPID-AUTO* and *CrystalStructure*. Rigaku/MS Inc., The Woodlands, Texas, USA.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Wang, R. H. & Han, L. (2004). *J. Mol. Struct.* **694**, 79–83.
Wang, R. H. & Jiang, F. L. (2004). *J. Mol. Struct.* **699**, 79–84.

supplementary materials

Acta Cryst. (2008). E64, o2104 [doi:10.1107/S1600536808032352]

N,N'-Bis(2-methoxyphenyl)biphenyl-2,2'-dicarboxamide

G.-Y. Wang, D. Li, D.-B. Qin, J.-W. Luo and L.-H. Guo

Comment

The distortion of diphenyl spacer about central bond not only endows dpa a peculiar characterization to link metal ions or metal clusters into macrocycles or helical chains, but also makes diphenic acid (H2dpa) can deprotonate partially forming hydrogen bonds of carboxylic groups to meet both geometric and energetic requirements. We here report the crystal structure of the title compound.

The C8—C13 and C14—C19 planes form the dihedral angle of 75.34 (9)°, and C1—C6 ring are nearly perpendicular to C14—C19 ring, with a dihedral angle of 85.69 (8)°. The molecular structure is stabilized by C—H···O and N—H···O intramolecular hydrogen bonds. In addition, weak C—H···O intermolecular hydrogen bonds are observed.

Experimental

The title compound was prepared according to the reported procedure of *M. Z. Gao & Gao (2002)*. Colourless single crystals suitable for X-ray diffraction were obtained by recrystallization from dimethyl sulfoxide.

Refinement

H atoms were placed in calculated positions with C—H = 0.93 Å, and N—H = 0.86 Å, and refined in riding mode with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$.

Figures




Fig. 1. The molecular structure of the title compound, showing 30% probability displacement ellipsoids and the atomic numbering.

N,N'-Bis(2-methoxyphenyl)biphenyl-2,2'-dicarboxamide

Crystal data

C₂₈H₂₄N₂O₄

$F_{000} = 952$

supplementary materials

$M_r = 452.49$

Monoclinic, Cc

$a = 18.184 (4) \text{ \AA}$

$b = 16.304 (3) \text{ \AA}$

$c = 7.9998 (16) \text{ \AA}$

$\beta = 108.90 (3)^\circ$

$V = 2243.9 (8) \text{ \AA}^3$

$Z = 4$

$D_x = 1.339 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 3190 reflections

$\theta = 1.7\text{--}27.9^\circ$

$\mu = 0.09 \text{ mm}^{-1}$

$T = 113 (2) \text{ K}$

Block, colourless

$0.16 \times 0.14 \times 0.10 \text{ mm}$

Data collection

Rigaku Saturn
diffractometer

Radiation source: rotating anode

Monochromator: confocal

$T = 113(2) \text{ K}$

ω scans

Absorption correction: multi-scan
(CrystalStructure; Rigaku/MSC, 2004)

$T_{\min} = 0.986$, $T_{\max} = 0.991$

6422 measured reflections

1991 independent reflections

1858 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.054$

$\theta_{\max} = 25.0^\circ$

$\theta_{\min} = 1.7^\circ$

$h = -21 \rightarrow 20$

$k = -17 \rightarrow 19$

$l = -9 \rightarrow 9$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.039$

$wR(F^2) = 0.093$

$S = 1.06$

1991 reflections

309 parameters

2 restraints

Primary atom site location: structure-invariant direct
methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring
sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0537P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.007$

$\Delta\rho_{\max} = 0.17 \text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.20 \text{ e \AA}^{-3}$

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -

factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. *R*-factors based on F^2 are statistically about twice as large as those based on *F*, and *R*- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	-0.01240 (15)	0.26417 (14)	0.9137 (3)	0.0338 (6)
O2	0.21854 (13)	0.17729 (12)	1.0081 (2)	0.0234 (5)
O3	0.45976 (12)	0.32159 (13)	1.0239 (2)	0.0239 (5)
O4	0.31213 (13)	0.07137 (13)	0.8380 (3)	0.0273 (5)
N1	0.09211 (15)	0.17216 (15)	0.8317 (3)	0.0222 (6)
H1N	0.0615	0.1801	0.7258	0.027*
N2	0.37809 (15)	0.21050 (14)	0.9645 (3)	0.0216 (6)
H2N	0.3350	0.1932	0.9743	0.026*
C1	0.00360 (19)	0.19053 (18)	0.9988 (4)	0.0229 (7)
C2	-0.0328 (2)	0.16090 (19)	1.1154 (4)	0.0277 (7)
H2	-0.0694	0.1928	1.1441	0.033*
C3	-0.0139 (2)	0.0831 (2)	1.1889 (4)	0.0298 (8)
H3	-0.0378	0.0632	1.2674	0.036*
C4	0.0399 (2)	0.0355 (2)	1.1457 (4)	0.0305 (8)
H4	0.0512	-0.0170	1.1926	0.037*
C5	0.07701 (19)	0.06580 (18)	1.0327 (3)	0.0258 (7)
H5	0.1148	0.0345	1.0072	0.031*
C6	0.05817 (18)	0.14248 (18)	0.9576 (3)	0.0210 (7)
C7	0.16777 (18)	0.18877 (17)	0.8643 (3)	0.0188 (6)
C8	0.18756 (18)	0.22120 (18)	0.7085 (3)	0.0201 (7)
C9	0.16253 (18)	0.17982 (18)	0.5469 (4)	0.0219 (7)
H9	0.1304	0.1342	0.5336	0.026*
C10	0.18522 (19)	0.20637 (19)	0.4066 (3)	0.0251 (7)
H10	0.1695	0.1779	0.3000	0.030*
C11	0.2317 (2)	0.27591 (19)	0.4260 (4)	0.0275 (7)
H11	0.2471	0.2940	0.3321	0.033*
C12	0.25501 (19)	0.31821 (19)	0.5843 (4)	0.0252 (7)
H12	0.2853	0.3651	0.5956	0.030*
C13	0.23333 (17)	0.29096 (18)	0.7281 (3)	0.0190 (6)
C14	0.25474 (18)	0.34128 (17)	0.8950 (3)	0.0204 (6)
C15	0.1978 (2)	0.39160 (18)	0.9223 (4)	0.0270 (7)
H15	0.1475	0.3901	0.8428	0.032*
C16	0.2152 (2)	0.4441 (2)	1.0670 (4)	0.0307 (8)
H16	0.1763	0.4766	1.0847	0.037*
C17	0.2897 (2)	0.44812 (19)	1.1841 (4)	0.0290 (8)
H17	0.3015	0.4837	1.2800	0.035*
C18	0.34668 (19)	0.39879 (18)	1.1576 (4)	0.0244 (7)
H18	0.3972	0.4021	1.2355	0.029*
C19	0.32979 (19)	0.34399 (18)	1.0156 (3)	0.0211 (6)
C20	0.39544 (18)	0.29130 (17)	1.0011 (3)	0.0210 (7)
C21	0.42452 (18)	0.15199 (18)	0.9115 (3)	0.0212 (7)
C22	0.5013 (2)	0.1627 (2)	0.9232 (4)	0.0317 (8)

supplementary materials

H22	0.5276	0.2099	0.9756	0.038*
C23	0.5398 (2)	0.1035 (2)	0.8573 (5)	0.0407 (9)
H23	0.5918	0.1111	0.8674	0.049*
C24	0.5015 (2)	0.0338 (2)	0.7773 (4)	0.0354 (8)
H24	0.5271	-0.0048	0.7305	0.042*
C25	0.4250 (2)	0.0217 (2)	0.7669 (4)	0.0275 (8)
H25	0.3988	-0.0252	0.7127	0.033*
C26	0.38714 (18)	0.07931 (18)	0.8372 (3)	0.0225 (7)
C28	0.2734 (2)	-0.0051 (2)	0.7829 (5)	0.0366 (9)
H28A	0.2719	-0.0168	0.6642	0.055*
H28B	0.2214	-0.0019	0.7873	0.055*
H28C	0.3011	-0.0480	0.8601	0.055*
C27	-0.0779 (3)	0.3085 (2)	0.9290 (5)	0.0458 (10)
H27A	-0.1234	0.2746	0.8892	0.069*
H27B	-0.0856	0.3571	0.8578	0.069*
H27C	-0.0684	0.3234	1.0502	0.069*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0436 (16)	0.0284 (13)	0.0359 (11)	0.0087 (11)	0.0218 (11)	0.0078 (9)
O2	0.0200 (12)	0.0303 (12)	0.0181 (10)	-0.0008 (9)	0.0038 (9)	0.0020 (8)
O3	0.0182 (13)	0.0273 (12)	0.0252 (10)	-0.0060 (9)	0.0054 (10)	-0.0002 (8)
O4	0.0194 (12)	0.0223 (12)	0.0408 (11)	-0.0041 (9)	0.0104 (10)	-0.0049 (9)
N1	0.0164 (14)	0.0333 (15)	0.0163 (11)	-0.0017 (11)	0.0045 (11)	0.0036 (9)
N2	0.0172 (14)	0.0205 (14)	0.0279 (13)	0.0000 (10)	0.0084 (11)	-0.0017 (10)
C1	0.0267 (18)	0.0191 (15)	0.0228 (14)	-0.0017 (13)	0.0076 (13)	0.0001 (11)
C2	0.030 (2)	0.0306 (19)	0.0255 (14)	0.0003 (14)	0.0135 (15)	-0.0028 (12)
C3	0.039 (2)	0.0303 (19)	0.0251 (14)	-0.0089 (15)	0.0176 (16)	0.0009 (12)
C4	0.042 (2)	0.0234 (18)	0.0272 (16)	0.0002 (14)	0.0133 (16)	0.0033 (12)
C5	0.030 (2)	0.0248 (17)	0.0225 (14)	0.0023 (14)	0.0082 (14)	-0.0017 (12)
C6	0.0203 (17)	0.0250 (17)	0.0163 (12)	-0.0035 (12)	0.0042 (13)	-0.0008 (11)
C7	0.0183 (17)	0.0183 (15)	0.0201 (14)	0.0011 (12)	0.0066 (14)	-0.0023 (11)
C8	0.0134 (16)	0.0294 (18)	0.0170 (12)	0.0040 (12)	0.0045 (12)	0.0016 (11)
C9	0.0174 (17)	0.0245 (17)	0.0219 (14)	-0.0004 (13)	0.0036 (13)	-0.0011 (11)
C10	0.0213 (19)	0.0347 (18)	0.0182 (13)	0.0025 (14)	0.0050 (13)	-0.0018 (11)
C11	0.028 (2)	0.0348 (19)	0.0219 (14)	0.0019 (14)	0.0117 (14)	0.0081 (12)
C12	0.0236 (19)	0.0249 (17)	0.0278 (15)	-0.0002 (13)	0.0094 (14)	0.0035 (12)
C13	0.0123 (16)	0.0232 (16)	0.0209 (13)	0.0025 (11)	0.0045 (12)	0.0027 (11)
C14	0.0205 (17)	0.0175 (15)	0.0238 (14)	-0.0011 (12)	0.0079 (13)	0.0020 (11)
C15	0.0231 (18)	0.0254 (17)	0.0305 (15)	0.0006 (14)	0.0057 (14)	0.0010 (12)
C16	0.032 (2)	0.0278 (19)	0.0365 (17)	0.0041 (14)	0.0167 (17)	-0.0022 (13)
C17	0.037 (2)	0.0222 (17)	0.0277 (15)	-0.0017 (14)	0.0105 (15)	-0.0026 (12)
C18	0.0267 (19)	0.0206 (17)	0.0239 (14)	-0.0020 (13)	0.0054 (13)	-0.0010 (11)
C19	0.0221 (17)	0.0192 (16)	0.0234 (14)	0.0007 (12)	0.0091 (14)	0.0030 (11)
C20	0.0223 (19)	0.0250 (17)	0.0143 (13)	-0.0018 (13)	0.0041 (13)	0.0012 (10)
C21	0.0207 (18)	0.0212 (16)	0.0222 (14)	0.0009 (12)	0.0076 (14)	-0.0005 (11)
C22	0.0246 (19)	0.0289 (18)	0.0423 (17)	-0.0024 (14)	0.0118 (16)	-0.0068 (14)

C23	0.0207 (19)	0.042 (2)	0.063 (2)	-0.0016 (15)	0.0182 (18)	-0.0110 (17)
C24	0.034 (2)	0.029 (2)	0.0465 (19)	0.0069 (15)	0.0177 (17)	-0.0012 (14)
C25	0.031 (2)	0.0201 (17)	0.0321 (16)	-0.0004 (13)	0.0110 (15)	-0.0015 (12)
C26	0.0204 (19)	0.0219 (16)	0.0241 (14)	-0.0011 (12)	0.0059 (13)	0.0019 (11)
C28	0.033 (2)	0.031 (2)	0.0477 (19)	-0.0132 (15)	0.0152 (17)	-0.0088 (14)
C27	0.062 (3)	0.037 (2)	0.049 (2)	0.0233 (19)	0.032 (2)	0.0113 (16)

Geometric parameters (Å, °)

O1—C1	1.364 (4)	C12—C13	1.403 (4)
O1—C27	1.431 (4)	C12—H12	0.9300
O2—C7	1.234 (3)	C13—C14	1.507 (4)
O3—C20	1.228 (4)	C14—C19	1.394 (4)
O4—C26	1.372 (4)	C14—C15	1.393 (4)
O4—C28	1.429 (4)	C15—C16	1.390 (4)
N1—C7	1.342 (4)	C15—H15	0.9300
N1—C6	1.425 (4)	C16—C17	1.377 (5)
N1—H1N	0.8600	C16—H16	0.9300
N2—C20	1.364 (4)	C17—C18	1.381 (5)
N2—C21	1.426 (4)	C17—H17	0.9300
N2—H2N	0.8600	C18—C19	1.398 (4)
C1—C6	1.385 (5)	C18—H18	0.9300
C1—C2	1.393 (4)	C19—C20	1.506 (4)
C2—C3	1.393 (5)	C21—C22	1.380 (5)
C2—H2	0.9300	C21—C26	1.399 (4)
C3—C4	1.377 (5)	C22—C23	1.393 (5)
C3—H3	0.9300	C22—H22	0.9300
C4—C5	1.383 (5)	C23—C24	1.377 (5)
C4—H4	0.9300	C23—H23	0.9300
C5—C6	1.381 (4)	C24—C25	1.381 (5)
C5—H5	0.9300	C24—H24	0.9300
C7—C8	1.501 (4)	C25—C26	1.387 (4)
C8—C13	1.388 (4)	C25—H25	0.9300
C8—C9	1.397 (4)	C28—H28A	0.9600
C9—C10	1.385 (4)	C28—H28B	0.9600
C9—H9	0.9300	C28—H28C	0.9600
C10—C11	1.392 (5)	C27—H27A	0.9600
C10—H10	0.9300	C27—H27B	0.9600
C11—C12	1.383 (4)	C27—H27C	0.9600
C11—H11	0.9300		
C1—O1—C27	116.8 (3)	C15—C14—C13	117.9 (3)
C26—O4—C28	118.1 (2)	C16—C15—C14	120.8 (3)
C7—N1—C6	125.6 (2)	C16—C15—H15	119.6
C7—N1—H1N	117.2	C14—C15—H15	119.6
C6—N1—H1N	117.2	C17—C16—C15	120.3 (3)
C20—N2—C21	126.3 (3)	C17—C16—H16	119.9
C20—N2—H2N	116.9	C15—C16—H16	119.9
C21—N2—H2N	116.9	C16—C17—C18	119.3 (3)
O1—C1—C6	115.6 (3)	C16—C17—H17	120.3

supplementary materials

O1—C1—C2	124.6 (3)	C18—C17—H17	120.3
C6—C1—C2	119.7 (3)	C17—C18—C19	121.2 (3)
C1—C2—C3	119.5 (3)	C17—C18—H18	119.4
C1—C2—H2	120.2	C19—C18—H18	119.4
C3—C2—H2	120.2	C14—C19—C18	119.4 (3)
C4—C3—C2	120.3 (3)	C14—C19—C20	123.5 (3)
C4—C3—H3	119.8	C18—C19—C20	117.1 (3)
C2—C3—H3	119.8	O3—C20—N2	124.3 (3)
C3—C4—C5	120.0 (3)	O3—C20—C19	120.0 (3)
C3—C4—H4	120.0	N2—C20—C19	115.7 (3)
C5—C4—H4	120.0	C22—C21—C26	118.6 (3)
C6—C5—C4	120.2 (3)	C22—C21—N2	125.3 (3)
C6—C5—H5	119.9	C26—C21—N2	116.1 (3)
C4—C5—H5	119.9	C21—C22—C23	120.5 (3)
C5—C6—C1	120.3 (3)	C21—C22—H22	119.7
C5—C6—N1	120.8 (3)	C23—C22—H22	119.7
C1—C6—N1	118.9 (3)	C24—C23—C22	120.5 (3)
O2—C7—N1	124.1 (3)	C24—C23—H23	119.7
O2—C7—C8	121.3 (3)	C22—C23—H23	119.7
N1—C7—C8	114.6 (2)	C23—C24—C25	119.6 (3)
C13—C8—C9	120.3 (3)	C23—C24—H24	120.2
C13—C8—C7	119.4 (2)	C25—C24—H24	120.2
C9—C8—C7	120.3 (3)	C24—C25—C26	120.1 (3)
C10—C9—C8	120.4 (3)	C24—C25—H25	120.0
C10—C9—H9	119.8	C26—C25—H25	120.0
C8—C9—H9	119.8	O4—C26—C25	124.3 (3)
C9—C10—C11	119.6 (3)	O4—C26—C21	115.1 (3)
C9—C10—H10	120.2	C25—C26—C21	120.6 (3)
C11—C10—H10	120.2	O4—C28—H28A	109.5
C12—C11—C10	120.2 (3)	O4—C28—H28B	109.5
C12—C11—H11	119.9	H28A—C28—H28B	109.5
C10—C11—H11	119.9	O4—C28—H28C	109.5
C11—C12—C13	120.6 (3)	H28A—C28—H28C	109.5
C11—C12—H12	119.7	H28B—C28—H28C	109.5
C13—C12—H12	119.7	O1—C27—H27A	109.5
C8—C13—C12	119.0 (3)	O1—C27—H27B	109.5
C8—C13—C14	121.3 (2)	H27A—C27—H27B	109.5
C12—C13—C14	119.6 (3)	O1—C27—H27C	109.5
C19—C14—C15	118.9 (3)	H27A—C27—H27C	109.5
C19—C14—C13	123.0 (3)	H27B—C27—H27C	109.5
C27—O1—C1—C6	169.2 (3)	C8—C13—C14—C15	-75.3 (4)
C27—O1—C1—C2	-8.3 (4)	C12—C13—C14—C15	99.9 (3)
O1—C1—C2—C3	177.3 (3)	C19—C14—C15—C16	0.3 (4)
C6—C1—C2—C3	-0.2 (4)	C13—C14—C15—C16	-175.3 (3)
C1—C2—C3—C4	-0.4 (5)	C14—C15—C16—C17	1.2 (5)
C2—C3—C4—C5	1.8 (5)	C15—C16—C17—C18	-0.8 (5)
C3—C4—C5—C6	-2.5 (4)	C16—C17—C18—C19	-1.0 (5)
C4—C5—C6—C1	1.8 (4)	C15—C14—C19—C18	-2.0 (4)
C4—C5—C6—N1	-175.7 (3)	C13—C14—C19—C18	173.3 (3)

O1—C1—C6—C5	-178.2 (3)	C15—C14—C19—C20	178.8 (3)
C2—C1—C6—C5	-0.5 (4)	C13—C14—C19—C20	-5.9 (4)
O1—C1—C6—N1	-0.6 (4)	C17—C18—C19—C14	2.4 (4)
C2—C1—C6—N1	177.1 (3)	C17—C18—C19—C20	-178.4 (3)
C7—N1—C6—C5	-65.1 (4)	C21—N2—C20—O3	-13.6 (4)
C7—N1—C6—C1	117.3 (3)	C21—N2—C20—C19	167.3 (2)
C6—N1—C7—O2	2.9 (5)	C14—C19—C20—O3	135.1 (3)
C6—N1—C7—C8	-178.7 (3)	C18—C19—C20—O3	-44.0 (4)
O2—C7—C8—C13	-50.2 (4)	C14—C19—C20—N2	-45.7 (4)
N1—C7—C8—C13	131.3 (3)	C18—C19—C20—N2	135.1 (3)
O2—C7—C8—C9	127.4 (3)	C20—N2—C21—C22	14.1 (4)
N1—C7—C8—C9	-51.1 (4)	C20—N2—C21—C26	-163.9 (2)
C13—C8—C9—C10	2.1 (5)	C26—C21—C22—C23	2.0 (5)
C7—C8—C9—C10	-175.5 (3)	N2—C21—C22—C23	-175.9 (3)
C8—C9—C10—C11	-1.6 (5)	C21—C22—C23—C24	0.9 (5)
C9—C10—C11—C12	0.0 (5)	C22—C23—C24—C25	-1.8 (6)
C10—C11—C12—C13	1.1 (5)	C23—C24—C25—C26	-0.2 (5)
C9—C8—C13—C12	-1.1 (4)	C28—O4—C26—C25	8.0 (4)
C7—C8—C13—C12	176.6 (3)	C28—O4—C26—C21	-173.0 (3)
C9—C8—C13—C14	174.1 (3)	C24—C25—C26—O4	-177.9 (3)
C7—C8—C13—C14	-8.2 (4)	C24—C25—C26—C21	3.1 (4)
C11—C12—C13—C8	-0.5 (5)	C22—C21—C26—O4	176.9 (3)
C11—C12—C13—C14	-175.8 (3)	N2—C21—C26—O4	-4.9 (3)
C8—C13—C14—C19	109.4 (3)	C22—C21—C26—C25	-4.0 (4)
C12—C13—C14—C19	-75.4 (4)	N2—C21—C26—C25	174.2 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1N \cdots O3 ⁱ	0.86	2.02	2.833 (3)	157
N2—H2N \cdots O2	0.86	2.24	3.081 (4)	167
N2—H2N \cdots O4	0.86	2.24	2.612 (3)	106
C22—H22 \cdots O3	0.93	2.30	2.885 (4)	120

Symmetry codes: (i) $x-1/2, -y+1/2, z-1/2$.

Fig. 1

