

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2,2,10-Trimethyl-2,3-dihydropyrano-[2,3-a]carbazol-4(11H)-one

 Makuteswaran Sridharan,^a Karnam J. Rajendra Prasad,^a Aimable Ngendahimana^b and Matthias Zeller^{b*}
^aDepartment of Chemistry, Bharathiar University, Coimbatore 641 046, Tamil Nadu, India, and ^bDepartment of Chemistry, Youngstown State University, One University Plaza, Youngstown, OH 44555, USA

Correspondence e-mail: mzeller@cc.yosu.edu


Received 22 September 2008; accepted 16 October 2008

 Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.040; wR factor = 0.109; data-to-parameter ratio = 18.3.

The title compound, $\text{C}_{18}\text{H}_{17}\text{NO}_2$, was prepared from 1-hydroxy-8-methylcarbazole and 3,3-dimethylacrylic acid with trifluoroacetic acid as the cyclization catalyst. Due to the $-\text{CMe}_2-$ group, the molecule is not quite planar. The packing is dominated by the strong $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds and some weaker $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\pi$ interactions. $\pi-\pi$ Stacking interactions [centroid-centroid separation = $3.806(2)$ Å] join neighboring molecules into loosely connected inversion dimers.

Related literature

Knölker & Reddy (2002) report on the isolation of pyranocarbazoles from various plant species. Sridharan *et al.* (2007) describe the synthesis of compounds related to the title compound. Sridharan, Rajendra Prasad & Zeller (2008) report the structure of the 9-methyl derivative of the title compound. Sridharan, Rajendra Prasad, Ngendahimana *et al.* (2008) report the structure of the 10-*H* derivative of the title compound.



Experimental

Crystal data

$\text{C}_{18}\text{H}_{17}\text{NO}_2$
 $M_r = 279.33$
 Monoclinic, $P2_1/c$
 $a = 12.9740(16)$ Å
 $b = 9.4195(12)$ Å
 $c = 12.8444(16)$ Å
 $\beta = 114.733(2)^\circ$
 $V = 1425.7(3)$ Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 100(2)$ K
 $0.53 \times 0.43 \times 0.19$ mm

Data collection

Bruker SMART APEX CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2007)
 $T_{\min} = 0.886$, $T_{\max} = 0.984$
 13755 measured reflections
 3526 independent reflections
 2941 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.109$
 $S = 1.03$
 3526 reflections
 193 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.31$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.26$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{O2}^{\text{i}}$	0.88	1.99	2.8634 (13)	173
$\text{C15}-\text{H15A}\cdots\text{O1}^{\text{ii}}$	0.99	2.59	3.5411 (15)	161

 Symmetry codes: (i) $x, -y + \frac{3}{2}, z + \frac{1}{2}$; (ii) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: Mercury (Macrae *et al.*, 2006); software used to prepare material for publication: SHELXTL.

The authors acknowledge UGC, New Delhi, India, for the award of a Major Research Project (grant No. F31-122/2005). MS thanks UGC, New Delhi, India, for the award of a research fellowship. The diffractometer was funded by the NSF (grant No. 0087210), the Ohio Board of Regents (grant No. CAP-491) and YSU.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2805).

References

- Bruker (2007). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
 Knölker, H. J. & Reddy, K. R. (2002). *Chem. Rev.* **102**, 4303–4427.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Sridharan, M., Prasad, K. J. R. & Zeller, M. (2007). *Acta Cryst.* **E63**, o4344.
 Sridharan, M., Rajendra Prasad, K. J., Ngendahimana, A. & Zeller, M. (2008). *Acta Cryst.* **E64**, o2156.
 Sridharan, M., Rajendra Prasad, K. J. & Zeller, M. (2008). *Acta Cryst.* **E64**, o2155.

supplementary materials

Acta Cryst. (2008). E64, o2157 [doi:10.1107/S1600536808033862]

2,2,10-Trimethyl-2,3-dihydropyrano[2,3-*a*]carbazol-4(11*H*)-one

M. Sridharan, K. J. R. Prasad, A. Ngendahimana and M. Zeller

Comment

Carbazole alkaloids have been isolated from the taxonomically related higher plants of the genus *Murraya*, *Glycosmis*, and *Clausena* from the family Rutaceae. Among the carbazole alkaloids pyranocarbazole alkaloids play a very important role. In this class girinimbine was the first member of the pyrano[3,2-*a*]carbazole alkaloid family to be isolated from *M. Koenigii* Spreng (Knölker & Reddy, 2002, and references therein). The isolation of these classes of compounds became an active area of study since these compounds possess high levels of biological and pharmacological activity. Hence we attempted to synthesize pyranocarbazoles in a simple and efficient route.

Using trifluoroacetic acid as the acylating agent we had been able to synthesize in high yields a range of pyranocarbazolones and we recently reported (Sridharan *et al.*, 2007) the synthesis and crystallographic behaviour of 2,3-dihydro-2,2,8-trimethylpyrano[2,3-*a*]carbazol-4-(11*H*)-one. As an extension of this research, and to further proof the credibility of trifluoroacetic acid as a good acylating agent, we further extended this synthetic route with a series of substituted 1-hydroxycarbazoles. The components thus synthesized were used as starting synthons to develop routes towards substituted pyranocarbazole derivatives. Herein we report the crystal structures of two of the compounds thus obtained: 2,3-dihydro-2,2,9-trimethylpyrano[2,3-*a*]carbazol-4-(11*H*)-one (Sridharan, Rajendra Prasad & Zeller, 2008), the title compound of the preceding article in this journal) and of the title compound 2,3-dihydro-2,2,10-trimethylpyrano[2,3-*a*]carbazol-4-(11*H*)-one (Figure 1).

The single-crystal structure confirmed the formation of the dihydropyrano-[2,3-*a*]carbazol-4(11*H*)-one framework as shown in Figure 2. Data collection and structure refinement were unproblematic and all structural parameters (bond lengths, angles, *etc*) are in the expected ranges. The molecules crystallize in a monoclinic setting in $P2_1/c$ with four largely planar molecules per unit cell. The plane defined by the sp^2 hybridized carbon atoms, the C1 methyl and C15 methylene carbon atoms, and the N and O atoms has an r.m.s. deviation from planarity of only 0.0754 Å. Of all the ring C atoms only C14 of the pyran C(Me)₂ unit is significantly out of plane with the atoms of the four fused rings, its deviation being 0.534 (1) Å. The pyran ring thus exhibits a half chair conformation.

One of the methyl groups of the C(Me)₂ unit is also located close to the average plane of the molecule (C18 with a deviation of 0.125 (2) Å). The other, C17, is however located 2.039 (2) Å away from this plane and thus makes the molecule as a whole not planar and prevents it from forming extensive π - π stacked entities in the solid state. The packing is thus indeed dominated by strong N—H \cdots O hydrogen bonds (Figure 3, Table 1) and some weaker C—H \cdots O (Table 1, Figure 4) and C—H \cdots π interactions (*e.g.* C18—H18 b \cdots Cg1ⁱⁱ = 2.94 Å with Cg1 being the ring C8 to C13 and ii = -*x*, -1/2 + *y*, 1/2 - *z*). The only significant π \cdots π stacking interaction with a centroid to centroid distance of 3.806 (2) Å is found between the pyrrole ring and the aromatic ring made up of C2 to C7 (Figure 4). Two neighboring molecules related by an inversion center are forming loosely connected dimers *via* two sets of these π - π interactions (symmetry operator 1 - *x*, 2 - *y*, 1 - *z*).

supplementary materials

The structures of the 2,2-dimethyl and the 2,2,10-methyl derivatives of the title compound are described in Sridharan, Rajendra Prasad, Ngendahimana *et al.* (2008) and Sridharan, Rajendra Prasad & Zeller (2008), the two preceding articles in this journal. For a more detailed comparison of structures and packing of the three two derivatives please see in Sridharan, Rajendra Prasad & Zeller (2008).

Experimental

1-hydroxy-8-methylcarbazole (0.001 mol) dissolved in 10 ml of trifluoroacetic acid and was heated with 3,3-dimethylacrylic acid (0.001 mol) at 323 K for 5 h. The reaction was monitored by TLC. After completion of the reaction, the excess trifluoroacetic acid was removed using rotary evaporation. The solid that precipitated out was poured onto ice water, then extracted using ethyl acetate and dried over anhydrous sodium sulfate and filtered. Then the solvent was removed under vacuum and the residue was purified by column chromatography on silica gel using petroleum ether/ethyl acetate (95:5 v/v) as eluant to yield yellow plates of (I) (0.239 g, 86%), m.p. 475–477 K.

Refinement

All hydrogen atoms were added in calculated positions with C—H = 0.99 Å (methylene), 0.95 Å (aromatic) and 0.98 Å (methyl) and N—H = 0.88 Å. They were refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C,N})$ or $1.5U_{\text{eq}}(\text{methyl C})$.

Figures



Fig. 1. Reaction sequence




Fig. 2. View of (I) showing xx% displacement ellipsoids. H atoms are represented in stick mode.




Fig. 3. Packing view of (I) down the *a* axis showing chains built by the N—H···O hydrogen bonds (indicated by blue dashed lines).




Fig. 4. Packing view of (I) showing the secondary C—H... π and C—H...O interactions indicated by green lines. Numbers given are distances in Å. N—H...O hydrogen bonds are omitted for clarity.

2,2,10-Trimethyl-2,3-dihydropyrano[2,3-a]carbazol-4(11H)-one

Crystal data

$C_{18}H_{17}NO_2$	$F_{000} = 592$
$M_r = 279.33$	$D_x = 1.301 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 12.9740 (16) \text{ \AA}$	Cell parameters from 4194 reflections
$b = 9.4195 (12) \text{ \AA}$	$\theta = 2.8\text{--}31.5^\circ$
$c = 12.8444 (16) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$\beta = 114.733 (2)^\circ$	$T = 100 (2) \text{ K}$
$V = 1425.7 (3) \text{ \AA}^3$	Plate, yellow
$Z = 4$	$0.53 \times 0.43 \times 0.19 \text{ mm}$

Data collection

Bruker APEXII CCD diffractometer	3526 independent reflections
Radiation source: fine-focus sealed tube	2941 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.027$
$T = 100(2) \text{ K}$	$\theta_{\text{max}} = 28.3^\circ$
ω scans	$\theta_{\text{min}} = 1.7^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2007)	$h = -17 \rightarrow 17$
$T_{\text{min}} = 0.886$, $T_{\text{max}} = 0.984$	$k = -12 \rightarrow 12$
13755 measured reflections	$l = -17 \rightarrow 17$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.040$	H-atom parameters constrained
$wR(F^2) = 0.109$	$w = 1/[\sigma^2(F_o^2) + (0.0532P)^2 + 0.5237P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
3526 reflections	$(\Delta/\sigma)_{\text{max}} = 0.002$
193 parameters	$\Delta\rho_{\text{max}} = 0.31 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.26 \text{ e \AA}^{-3}$

supplementary materials

Primary atom site location: structure-invariant direct methods Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.30161 (12)	1.23026 (14)	0.61446 (11)	0.0287 (3)
H1A	0.3274	1.1580	0.6749	0.043*
H1B	0.2199	1.2193	0.5680	0.043*
H1C	0.3168	1.3249	0.6493	0.043*
C2	0.36377 (10)	1.21263 (12)	0.53998 (10)	0.0205 (2)
C3	0.44870 (10)	1.30290 (13)	0.54197 (11)	0.0235 (3)
H3	0.4698	1.3806	0.5939	0.028*
C4	0.50503 (11)	1.28432 (13)	0.47042 (11)	0.0244 (3)
H4	0.5636	1.3484	0.4757	0.029*
C5	0.47647 (10)	1.17431 (13)	0.39258 (10)	0.0217 (2)
H5	0.5142	1.1624	0.3437	0.026*
C6	0.39071 (10)	1.08061 (12)	0.38720 (10)	0.0186 (2)
C7	0.33727 (9)	1.09998 (12)	0.46184 (10)	0.0178 (2)
C8	0.25796 (9)	0.90879 (12)	0.35603 (9)	0.0165 (2)
C9	0.33879 (9)	0.95775 (12)	0.31816 (10)	0.0176 (2)
C10	0.35259 (10)	0.88720 (13)	0.22829 (10)	0.0197 (2)
H10	0.4078	0.9184	0.2029	0.024*
C11	0.28477 (10)	0.77237 (13)	0.17803 (10)	0.0199 (2)
H11	0.2932	0.7246	0.1169	0.024*
C12	0.20236 (10)	0.72342 (12)	0.21537 (10)	0.0176 (2)
C13	0.19004 (9)	0.79075 (12)	0.30620 (10)	0.0165 (2)
C14	0.06708 (10)	0.60905 (12)	0.32183 (10)	0.0208 (2)
C15	0.03119 (10)	0.58064 (13)	0.19434 (10)	0.0199 (2)
H15A	0.0054	0.4810	0.1773	0.024*
H15B	-0.0338	0.6428	0.1493	0.024*
C16	0.12561 (10)	0.60623 (12)	0.15718 (10)	0.0186 (2)
C17	0.15681 (12)	0.50413 (14)	0.39605 (11)	0.0288 (3)
H17A	0.2229	0.5095	0.3778	0.043*
H17B	0.1255	0.4078	0.3811	0.043*
H17C	0.1798	0.5275	0.4771	0.043*
C18	-0.03389 (12)	0.61299 (15)	0.35277 (12)	0.0302 (3)

H18A	-0.0085	0.6424	0.4329	0.045*
H18B	-0.0681	0.5183	0.3425	0.045*
H18C	-0.0902	0.6808	0.3029	0.045*
N1	0.25730 (8)	0.99440 (10)	0.44254 (8)	0.0176 (2)
H1	0.2137	0.9838	0.4791	0.021*
O1	0.11424 (7)	0.75262 (9)	0.34888 (7)	0.01967 (19)
O2	0.13335 (7)	0.53564 (10)	0.08062 (7)	0.0245 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0343 (7)	0.0268 (7)	0.0265 (6)	-0.0040 (5)	0.0145 (6)	-0.0059 (5)
C2	0.0223 (6)	0.0185 (5)	0.0189 (5)	0.0003 (4)	0.0068 (5)	0.0011 (4)
C3	0.0256 (6)	0.0190 (6)	0.0220 (6)	-0.0024 (5)	0.0061 (5)	0.0003 (4)
C4	0.0227 (6)	0.0229 (6)	0.0251 (6)	-0.0052 (5)	0.0076 (5)	0.0031 (5)
C5	0.0201 (6)	0.0238 (6)	0.0215 (6)	-0.0018 (5)	0.0090 (5)	0.0040 (5)
C6	0.0187 (5)	0.0185 (5)	0.0182 (5)	0.0004 (4)	0.0074 (4)	0.0025 (4)
C7	0.0173 (5)	0.0168 (5)	0.0185 (5)	0.0008 (4)	0.0068 (4)	0.0031 (4)
C8	0.0177 (5)	0.0170 (5)	0.0160 (5)	0.0018 (4)	0.0083 (4)	0.0024 (4)
C9	0.0175 (5)	0.0179 (5)	0.0182 (5)	0.0002 (4)	0.0083 (4)	0.0032 (4)
C10	0.0192 (5)	0.0229 (6)	0.0209 (6)	-0.0001 (4)	0.0122 (5)	0.0022 (4)
C11	0.0214 (6)	0.0230 (6)	0.0191 (5)	0.0014 (4)	0.0121 (5)	0.0002 (4)
C12	0.0181 (5)	0.0185 (5)	0.0182 (5)	0.0008 (4)	0.0096 (4)	0.0011 (4)
C13	0.0163 (5)	0.0176 (5)	0.0177 (5)	0.0015 (4)	0.0090 (4)	0.0024 (4)
C14	0.0255 (6)	0.0189 (6)	0.0219 (6)	-0.0073 (4)	0.0137 (5)	-0.0034 (4)
C15	0.0200 (5)	0.0218 (6)	0.0200 (5)	-0.0038 (4)	0.0104 (5)	-0.0035 (4)
C16	0.0199 (5)	0.0197 (6)	0.0177 (5)	0.0019 (4)	0.0093 (4)	0.0019 (4)
C17	0.0388 (7)	0.0219 (6)	0.0242 (6)	-0.0039 (5)	0.0117 (6)	0.0022 (5)
C18	0.0349 (7)	0.0347 (7)	0.0312 (7)	-0.0148 (6)	0.0237 (6)	-0.0103 (6)
N1	0.0196 (5)	0.0175 (5)	0.0183 (5)	-0.0013 (4)	0.0106 (4)	-0.0008 (4)
O1	0.0229 (4)	0.0189 (4)	0.0231 (4)	-0.0050 (3)	0.0155 (4)	-0.0034 (3)
O2	0.0277 (5)	0.0264 (5)	0.0239 (4)	-0.0031 (4)	0.0153 (4)	-0.0064 (3)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.4963 (18)	C11—C12	1.4195 (16)
C1—H1A	0.9800	C11—H11	0.9500
C1—H1B	0.9800	C12—C13	1.3945 (16)
C1—H1C	0.9800	C12—C16	1.4649 (16)
C2—C3	1.3836 (17)	C13—O1	1.3594 (13)
C2—C7	1.4010 (16)	C14—O1	1.4650 (13)
C3—C4	1.4040 (19)	C14—C18	1.5202 (17)
C3—H3	0.9500	C14—C17	1.5205 (18)
C4—C5	1.3787 (18)	C14—C15	1.5270 (16)
C4—H4	0.9500	C15—C16	1.5086 (16)
C5—C6	1.3991 (16)	C15—H15A	0.9900
C5—H5	0.9500	C15—H15B	0.9900
C6—C7	1.4103 (16)	C16—O2	1.2254 (14)
C6—C9	1.4420 (16)	C17—H17A	0.9800

supplementary materials

C7—N1	1.3826 (14)	C17—H17B	0.9800
C8—N1	1.3759 (14)	C17—H17C	0.9800
C8—C13	1.3966 (16)	C18—H18A	0.9800
C8—C9	1.4057 (15)	C18—H18B	0.9800
C9—C10	1.4068 (16)	C18—H18C	0.9800
C10—C11	1.3732 (17)	N1—H1	0.8800
C10—H10	0.9500		
C2—C1—H1A	109.5	C13—C12—C16	118.57 (10)
C2—C1—H1B	109.5	C11—C12—C16	121.31 (10)
H1A—C1—H1B	109.5	O1—C13—C12	124.94 (10)
C2—C1—H1C	109.5	O1—C13—C8	116.73 (10)
H1A—C1—H1C	109.5	C12—C13—C8	118.31 (10)
H1B—C1—H1C	109.5	O1—C14—C18	103.62 (9)
C3—C2—C7	115.69 (11)	O1—C14—C17	108.49 (10)
C3—C2—C1	123.92 (11)	C18—C14—C17	111.68 (11)
C7—C2—C1	120.39 (11)	O1—C14—C15	109.02 (9)
C2—C3—C4	122.68 (12)	C18—C14—C15	112.03 (10)
C2—C3—H3	118.7	C17—C14—C15	111.62 (10)
C4—C3—H3	118.7	C16—C15—C14	112.81 (9)
C5—C4—C3	120.85 (11)	C16—C15—H15A	109.0
C5—C4—H4	119.6	C14—C15—H15A	109.0
C3—C4—H4	119.6	C16—C15—H15B	109.0
C4—C5—C6	118.48 (11)	C14—C15—H15B	109.0
C4—C5—H5	120.8	H15A—C15—H15B	107.8
C6—C5—H5	120.8	O2—C16—C12	123.50 (11)
C5—C6—C7	119.46 (11)	O2—C16—C15	121.14 (11)
C5—C6—C9	133.93 (11)	C12—C16—C15	115.31 (10)
C7—C6—C9	106.61 (10)	C14—C17—H17A	109.5
N1—C7—C2	127.88 (11)	C14—C17—H17B	109.5
N1—C7—C6	109.30 (10)	H17A—C17—H17B	109.5
C2—C7—C6	122.81 (11)	C14—C17—H17C	109.5
N1—C8—C13	128.38 (10)	H17A—C17—H17C	109.5
N1—C8—C9	110.17 (10)	H17B—C17—H17C	109.5
C13—C8—C9	121.44 (10)	C14—C18—H18A	109.5
C8—C9—C10	119.89 (11)	C14—C18—H18B	109.5
C8—C9—C6	105.94 (10)	H18A—C18—H18B	109.5
C10—C9—C6	134.16 (11)	C14—C18—H18C	109.5
C11—C10—C9	118.78 (10)	H18A—C18—H18C	109.5
C11—C10—H10	120.6	H18B—C18—H18C	109.5
C9—C10—H10	120.6	C8—N1—C7	107.97 (10)
C10—C11—C12	121.47 (11)	C8—N1—H1	126.0
C10—C11—H11	119.3	C7—N1—H1	126.0
C12—C11—H11	119.3	C13—O1—C14	116.68 (9)
C13—C12—C11	120.07 (11)		
C7—C2—C3—C4	0.25 (18)	C11—C12—C13—O1	-179.88 (10)
C1—C2—C3—C4	-179.51 (12)	C16—C12—C13—O1	2.67 (17)
C2—C3—C4—C5	0.89 (19)	C11—C12—C13—C8	1.94 (17)
C3—C4—C5—C6	-0.60 (18)	C16—C12—C13—C8	-175.51 (10)

C4—C5—C6—C7	-0.79 (17)	N1—C8—C13—O1	-1.17 (17)
C4—C5—C6—C9	179.38 (12)	C9—C8—C13—O1	-179.73 (10)
C3—C2—C7—N1	179.55 (11)	N1—C8—C13—C12	177.16 (11)
C1—C2—C7—N1	-0.67 (19)	C9—C8—C13—C12	-1.39 (16)
C3—C2—C7—C6	-1.70 (17)	O1—C14—C15—C16	53.97 (13)
C1—C2—C7—C6	178.07 (11)	C18—C14—C15—C16	168.06 (10)
C5—C6—C7—N1	-179.03 (10)	C17—C14—C15—C16	-65.85 (13)
C9—C6—C7—N1	0.84 (13)	C13—C12—C16—O2	-176.05 (11)
C5—C6—C7—C2	2.02 (17)	C11—C12—C16—O2	6.53 (18)
C9—C6—C7—C2	-178.11 (10)	C13—C12—C16—C15	6.45 (15)
N1—C8—C9—C10	-178.90 (10)	C11—C12—C16—C15	-170.97 (11)
C13—C8—C9—C10	-0.10 (17)	C14—C15—C16—O2	147.37 (11)
N1—C8—C9—C6	0.40 (12)	C14—C15—C16—C12	-35.06 (14)
C13—C8—C9—C6	179.19 (10)	C13—C8—N1—C7	-178.57 (11)
C5—C6—C9—C8	179.10 (12)	C9—C8—N1—C7	0.12 (12)
C7—C6—C9—C8	-0.74 (12)	C2—C7—N1—C8	178.28 (11)
C5—C6—C9—C10	-1.8 (2)	C6—C7—N1—C8	-0.60 (12)
C7—C6—C9—C10	178.40 (12)	C12—C13—O1—C14	18.95 (16)
C8—C9—C10—C11	1.04 (17)	C8—C13—O1—C14	-162.84 (10)
C6—C9—C10—C11	-178.01 (12)	C18—C14—O1—C13	-165.62 (10)
C9—C10—C11—C12	-0.48 (17)	C17—C14—O1—C13	75.58 (12)
C10—C11—C12—C13	-1.04 (18)	C15—C14—O1—C13	-46.16 (13)
C10—C11—C12—C16	176.35 (11)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1 \cdots O2 ⁱ	0.88	1.99	2.8634 (13)	173
C15—H15A \cdots O1 ⁱⁱ	0.99	2.59	3.5411 (15)	161

Symmetry codes: (i) $x, -y+3/2, z+1/2$; (ii) $-x, y-1/2, -z+1/2$.

Fig. 1




Fig. 2




Fig. 3




Fig. 4

