

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

rac-(3-Hydroxy-2-phenylpropionato- κ O)-triphenyltin(IV)

 Mostafa M. Amini,^a Taraneh Hajiashrafi,^b Ali Nemati Kharat^b and Seik Weng Ng^{c*}
^aDepartment of Chemistry, Shahid Beheshti University, Tehran, Iran, ^bSchool of Chemistry, College of Science, Tehran University, Tehran, Iran, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 7 October 2008; accepted 8 October 2008

 Key indicators: single-crystal X-ray study; $T = 112$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.019; wR factor = 0.054; data-to-parameter ratio = 23.5.

The Sn^{IV} atom in the monomeric title compound, [Sn(C₆H₅)₃(C₉H₉O₃)] exists in a distorted SnC₃O tetrahedral geometry. In the crystal structure, inversion dimers arise from pairs of O—H...O hydrogen bonds.

Related literature

For reviews of organotin carboxylates, see: Tiekink (1991, 1994).



Experimental

Crystal data

 [Sn(C₆H₅)₃(C₉H₉O₃)]

 $M_r = 515.15$

 Triclinic, $P\bar{1}$
 $a = 9.3880$ (2) Å

 $b = 9.4899$ (2) Å

 $c = 14.3399$ (2) Å

 $\alpha = 90.087$ (1)°

 $\beta = 103.664$ (1)°

 $\gamma = 112.758$ (1)°

 $V = 1138.58$ (4) Å³
 $Z = 2$

 Mo $K\alpha$ radiation

 $\mu = 1.15$ mm⁻¹
 $T = 112$ (2) K

 $0.39 \times 0.32 \times 0.31$ mm

Data collection

 Bruker APEXII diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.663$, $T_{\max} = 0.717$

 33531 measured reflections
 6612 independent reflections
 6339 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.019$
 $wR(F^2) = 0.054$
 $S = 1.05$

6612 reflections

281 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.64$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.35$ e Å⁻³
Table 1

Selected bond lengths (Å).

Sn1—O1	2.082 (1)	Sn1—C7	2.122 (1)
Sn1—C1	2.123 (1)	Sn1—C13	2.128 (1)

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O3—H3 ⁱ ...O2 ⁱ	0.84	1.98	2.819 (2)	175

 Symmetry code: (i) $-x + 1, -y + 1, -z + 1$.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2008).

We thank the University of Canterbury, New Zealand, for the diffraction measurements, and the Vice-President's Office for Research Affairs of Shahid Beheshti University and the University of Malaya for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2813).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Tiekink, E. R. T. (1991). *Appl. Organomet. Chem.* **5**, 1–23.
 Tiekink, E. R. T. (1994). *Trends Organomet. Chem.* **1**, 71–116.
 Westrip, S. P. (2008). publCIF. In preparation.

supplementary materials

Acta Cryst. (2008). E64, m1403 [doi:10.1107/S160053680803242X]

***rac*-(3-Hydroxy-2-phenylpropionato- κ O)triphenyltin(IV)**

M. M. Amini, T. Hajjashrafi, A. N. Kharat and S. W. Ng

Comment

The monomeric structure found for triphenyltin 2-hydroxy-3-phenylpropionate (Scheme I, Fig. 1) conforms to expectations based on the presence of bulky substituents (Tiekink, 1991, 1994). In the arbitrarily chosen asymmetric unit, C20 has R configuration, but crystal symmetry generates a racemic mixture. Selected geometrical data are given in Tables 1 and 2.

Experimental

Triphenyltin hydroxide (1.0 g, 2.7 mmol) and *d,l*-tropic acid (0.45 g, 2.7 mmol) were heated in toluene (100 ml) in a Dean–Stark water-separator until all the water had been removed. The solvent was removed under reduced pressure to leave a white solid. The solid was recrystallized from mixture of chloroform, hexane and toluene (2:1:1 *v/v*) to give colorless blocks of (I), m.p. 385–386 K.

IR (KBr, cm^{-1}): 3449 (OH), 1627 (CO, *asym*), 1355 (CO, *sym*), 576, 601 (Sn—C). ^1H NMR (CDCl_3): 3.72–4.08 (m, 3H), 7.28–7.78 (20H, C_6H_5) p.p.m. ^{13}C NMR (CDCl_3): 53.90 (CH), 65.08 (CH_2) 127.5–137.7 (C_6H_5) p.p.m. ^{119}Sn NMR (CDCl_3): -83.1 p.p.m. Mass spectrum (*m/e*): 515 (M-1) $[\text{Ph}_3\text{SnO}_2\text{CCH}(\text{CH}_2\text{O})\text{Ph}]^+$, 439 (M-Ph) $[\text{Ph}_3\text{SnO}_2\text{CCH}(\text{CH}_2\text{OH})]^+$.

Refinement

The hydrogen atoms were placed in calculated positions (C—H = 0.95–1.00 Å, O—H = 0.84 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{carrier})$.

Figures




Fig. 1. The molecular structure of (I) showing displacement ellipsoids at the 70% probability level; hydrogen atoms are shown as spheres of arbitrary radius.

***rac*-(3-Hydroxy-2-phenylpropionato- κ O)triphenyltin(IV)**

Crystal data

$[\text{Sn}(\text{C}_6\text{H}_5)_3(\text{C}_9\text{H}_9\text{O}_3)]$

$M_r = 515.15$

$Z = 2$

$F_{000} = 520$

supplementary materials

Triclinic, $P\bar{1}$	$D_x = 1.503 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation
$a = 9.3880 (2) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 9.4899 (2) \text{ \AA}$	Cell parameters from 9425 reflections
$c = 14.3399 (2) \text{ \AA}$	$\theta = 2.3\text{--}32.6^\circ$
$\alpha = 90.087 (1)^\circ$	$\mu = 1.15 \text{ mm}^{-1}$
$\beta = 103.664 (1)^\circ$	$T = 112 (2) \text{ K}$
$\gamma = 112.758 (1)^\circ$	Block, colourless
$V = 1138.58 (4) \text{ \AA}^3$	$0.39 \times 0.32 \times 0.31 \text{ mm}$

Data collection

Bruker APEXII diffractometer	6612 independent reflections
Radiation source: medium-focus sealed tube	6339 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.020$
$T = 112(2) \text{ K}$	$\theta_{\text{max}} = 30.0^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.5^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -13 \rightarrow 13$
$T_{\text{min}} = 0.663$, $T_{\text{max}} = 0.717$	$k = -13 \rightarrow 13$
33531 measured reflections	$l = -19 \rightarrow 20$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.019$	H-atom parameters constrained
$wR(F^2) = 0.054$	$w = 1/[\sigma^2(F_o^2) + (0.0304P)^2 + 0.5255P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
6612 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
281 parameters	$\Delta\rho_{\text{max}} = 0.64 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.35 \text{ e \AA}^{-3}$
	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn1	0.588742 (10)	0.636637 (10)	0.826066 (6)	0.02043 (3)
O1	0.74512 (13)	0.76508 (12)	0.74695 (7)	0.0268 (2)
O2	0.52531 (13)	0.67054 (13)	0.62949 (8)	0.0303 (2)
O3	0.71606 (16)	0.55344 (13)	0.51412 (10)	0.0394 (3)
H3	0.6449	0.4826	0.4734	0.059*
C1	0.54423 (16)	0.40565 (15)	0.78448 (9)	0.0222 (2)

C2	0.4203 (2)	0.28743 (18)	0.81068 (12)	0.0330 (3)
H2	0.3515	0.3111	0.8411	0.040*
C3	0.3973 (3)	0.1349 (2)	0.79226 (15)	0.0441 (4)
H3A	0.3129	0.0549	0.8102	0.053*
C4	0.4965 (2)	0.09935 (19)	0.74803 (13)	0.0400 (4)
H4	0.4812	-0.0048	0.7365	0.048*
C5	0.6186 (2)	0.21545 (19)	0.72034 (12)	0.0334 (3)
H5	0.6856	0.1908	0.6888	0.040*
C6	0.64278 (18)	0.36822 (17)	0.73878 (11)	0.0274 (3)
H6	0.7269	0.4476	0.7201	0.033*
C7	0.38486 (17)	0.68487 (16)	0.82227 (12)	0.0275 (3)
C8	0.25851 (18)	0.64580 (18)	0.73831 (13)	0.0333 (3)
H8	0.2686	0.6072	0.6802	0.040*
C9	0.1186 (2)	0.6635 (2)	0.74001 (16)	0.0425 (4)
H9	0.0335	0.6372	0.6831	0.051*
C10	0.1040 (2)	0.7191 (2)	0.82417 (19)	0.0480 (5)
H10	0.0082	0.7305	0.8252	0.058*
C11	0.2273 (2)	0.7587 (2)	0.90783 (18)	0.0458 (4)
H11	0.2160	0.7978	0.9655	0.055*
C12	0.3687 (2)	0.74112 (19)	0.90717 (14)	0.0358 (3)
H12	0.4531	0.7675	0.9644	0.043*
C13	0.76077 (16)	0.71155 (16)	0.96187 (10)	0.0231 (2)
C14	0.8361 (2)	0.61586 (19)	1.00105 (11)	0.0323 (3)
H14	0.8035	0.5161	0.9696	0.039*
C15	0.9590 (2)	0.6657 (2)	1.08601 (13)	0.0427 (4)
H15	1.0100	0.6001	1.1120	0.051*
C16	1.0063 (2)	0.8105 (2)	1.13220 (12)	0.0407 (4)
H16	1.0911	0.8451	1.1894	0.049*
C17	0.9304 (2)	0.9052 (2)	1.09543 (12)	0.0366 (3)
H17	0.9617	1.0038	1.1282	0.044*
C18	0.80848 (19)	0.85653 (17)	1.01073 (11)	0.0296 (3)
H18	0.7571	0.9223	0.9858	0.036*
C19	0.67104 (17)	0.73973 (16)	0.65568 (10)	0.0242 (2)
C20	0.77747 (17)	0.80071 (16)	0.58653 (10)	0.0240 (2)
H20	0.8833	0.7969	0.6167	0.029*
C21	0.7052 (2)	0.69608 (18)	0.49250 (11)	0.0310 (3)
H21A	0.5922	0.6808	0.4664	0.037*
H21B	0.7649	0.7415	0.4441	0.037*
C22	0.80597 (18)	0.96785 (16)	0.57297 (10)	0.0248 (3)
C23	0.6983 (2)	1.0097 (2)	0.50719 (13)	0.0394 (4)
H23	0.6010	0.9326	0.4703	0.047*
C24	0.7320 (3)	1.1637 (2)	0.49495 (15)	0.0481 (5)
H24	0.6583	1.1908	0.4490	0.058*
C25	0.8706 (3)	1.2768 (2)	0.54866 (16)	0.0439 (4)
H25	0.8932	1.3818	0.5399	0.053*
C26	0.9769 (2)	1.2367 (2)	0.61539 (17)	0.0444 (4)
H26	1.0719	1.3146	0.6538	0.053*
C27	0.9458 (2)	1.08338 (19)	0.62678 (13)	0.0348 (3)
H27	1.0212	1.0571	0.6719	0.042*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn1	0.01937 (5)	0.01980 (5)	0.02268 (5)	0.00779 (3)	0.00640 (3)	0.00547 (3)
O1	0.0292 (5)	0.0248 (5)	0.0234 (4)	0.0066 (4)	0.0084 (4)	0.0063 (4)
O2	0.0288 (5)	0.0332 (5)	0.0280 (5)	0.0116 (4)	0.0067 (4)	0.0078 (4)
O3	0.0394 (6)	0.0223 (5)	0.0536 (7)	0.0103 (5)	0.0104 (6)	-0.0018 (5)
C1	0.0223 (6)	0.0217 (6)	0.0209 (5)	0.0075 (5)	0.0044 (4)	0.0048 (4)
C2	0.0350 (8)	0.0257 (7)	0.0369 (8)	0.0060 (6)	0.0173 (6)	0.0051 (6)
C3	0.0518 (11)	0.0232 (7)	0.0515 (10)	0.0028 (7)	0.0239 (9)	0.0067 (7)
C4	0.0555 (11)	0.0224 (7)	0.0398 (8)	0.0140 (7)	0.0108 (8)	0.0027 (6)
C5	0.0403 (8)	0.0319 (7)	0.0325 (7)	0.0193 (7)	0.0089 (6)	0.0017 (6)
C6	0.0268 (6)	0.0253 (6)	0.0306 (7)	0.0099 (5)	0.0093 (5)	0.0052 (5)
C7	0.0222 (6)	0.0217 (6)	0.0409 (8)	0.0092 (5)	0.0115 (6)	0.0083 (5)
C8	0.0246 (7)	0.0277 (7)	0.0467 (9)	0.0106 (6)	0.0073 (6)	0.0088 (6)
C9	0.0257 (7)	0.0337 (8)	0.0679 (12)	0.0139 (6)	0.0081 (8)	0.0113 (8)
C10	0.0301 (8)	0.0326 (8)	0.0894 (16)	0.0169 (7)	0.0221 (9)	0.0141 (9)
C11	0.0437 (10)	0.0322 (8)	0.0731 (13)	0.0175 (7)	0.0318 (10)	0.0050 (8)
C12	0.0335 (8)	0.0299 (7)	0.0485 (9)	0.0126 (6)	0.0186 (7)	0.0053 (7)
C13	0.0240 (6)	0.0240 (6)	0.0216 (5)	0.0086 (5)	0.0082 (5)	0.0050 (5)
C14	0.0395 (8)	0.0328 (7)	0.0267 (7)	0.0191 (7)	0.0040 (6)	0.0042 (6)
C15	0.0461 (10)	0.0534 (11)	0.0311 (8)	0.0281 (9)	0.0000 (7)	0.0070 (7)
C16	0.0326 (8)	0.0551 (11)	0.0251 (7)	0.0110 (8)	0.0016 (6)	0.0002 (7)
C17	0.0330 (8)	0.0321 (8)	0.0335 (8)	0.0014 (6)	0.0084 (6)	-0.0048 (6)
C18	0.0292 (7)	0.0246 (6)	0.0326 (7)	0.0075 (5)	0.0089 (6)	0.0032 (5)
C19	0.0301 (6)	0.0199 (6)	0.0240 (6)	0.0107 (5)	0.0083 (5)	0.0067 (5)
C20	0.0238 (6)	0.0228 (6)	0.0255 (6)	0.0083 (5)	0.0080 (5)	0.0061 (5)
C21	0.0367 (8)	0.0260 (7)	0.0300 (7)	0.0096 (6)	0.0133 (6)	0.0034 (5)
C22	0.0308 (7)	0.0214 (6)	0.0237 (6)	0.0090 (5)	0.0122 (5)	0.0063 (5)
C23	0.0438 (9)	0.0311 (8)	0.0345 (8)	0.0124 (7)	-0.0016 (7)	0.0065 (6)
C24	0.0682 (13)	0.0386 (9)	0.0429 (10)	0.0293 (9)	0.0105 (9)	0.0182 (8)
C25	0.0600 (11)	0.0237 (7)	0.0595 (11)	0.0159 (8)	0.0375 (10)	0.0160 (7)
C26	0.0344 (8)	0.0268 (8)	0.0681 (13)	0.0036 (6)	0.0210 (8)	-0.0049 (8)
C27	0.0306 (7)	0.0296 (7)	0.0431 (9)	0.0118 (6)	0.0075 (6)	-0.0005 (6)

Geometric parameters (\AA , $^\circ$)

Sn1—O1	2.082 (1)	C12—H12	0.9500
Sn1—C1	2.123 (1)	C13—C18	1.399 (2)
Sn1—C7	2.122 (1)	C13—C14	1.397 (2)
Sn1—C13	2.128 (1)	C14—C15	1.397 (2)
O1—C19	1.3016 (17)	C14—H14	0.9500
O2—C19	1.2269 (18)	C15—C16	1.383 (3)
O3—C21	1.4254 (19)	C15—H15	0.9500
O3—H3	0.8400	C16—C17	1.383 (3)
C1—C6	1.397 (2)	C16—H16	0.9500
C1—C2	1.3970 (19)	C17—C18	1.390 (2)
C2—C3	1.394 (2)	C17—H17	0.9500

C2—H2	0.9500	C18—H18	0.9500
C3—C4	1.380 (3)	C19—C20	1.5294 (19)
C3—H3A	0.9500	C20—C22	1.5240 (19)
C4—C5	1.387 (3)	C20—C21	1.519 (2)
C4—H4	0.9500	C20—H20	1.0000
C5—C6	1.392 (2)	C21—H21A	0.9900
C5—H5	0.9500	C21—H21B	0.9900
C6—H6	0.9500	C22—C23	1.389 (2)
C7—C12	1.392 (2)	C22—C27	1.388 (2)
C7—C8	1.406 (2)	C23—C24	1.392 (2)
C8—C9	1.393 (2)	C23—H23	0.9500
C8—H8	0.9500	C24—C25	1.372 (3)
C9—C10	1.373 (3)	C24—H24	0.9500
C9—H9	0.9500	C25—C26	1.378 (3)
C10—C11	1.388 (3)	C25—H25	0.9500
C10—H10	0.9500	C26—C27	1.387 (2)
C11—C12	1.402 (2)	C26—H26	0.9500
C11—H11	0.9500	C27—H27	0.9500
O1—Sn1—C1	104.08 (5)	C15—C14—H14	119.7
O1—Sn1—C7	117.51 (5)	C13—C14—H14	119.7
O1—Sn1—C13	94.95 (5)	C16—C15—C14	119.92 (17)
C1—Sn1—C7	116.29 (5)	C16—C15—H15	120.0
C1—Sn1—C13	109.17 (5)	C14—C15—H15	120.0
C7—Sn1—C13	112.54 (6)	C15—C16—C17	120.18 (16)
C19—O1—Sn1	109.50 (9)	C15—C16—H16	119.9
C21—O3—H3	109.5	C17—C16—H16	119.9
C6—C1—C2	118.99 (13)	C18—C17—C16	120.14 (16)
C6—C1—Sn1	122.04 (10)	C18—C17—H17	119.9
C2—C1—Sn1	118.77 (11)	C16—C17—H17	119.9
C3—C2—C1	120.14 (15)	C17—C18—C13	120.60 (15)
C3—C2—H2	119.9	C17—C18—H18	119.7
C1—C2—H2	119.9	C13—C18—H18	119.7
C4—C3—C2	120.38 (16)	O2—C19—O1	120.51 (13)
C4—C3—H3A	119.8	O2—C19—C20	123.92 (13)
C2—C3—H3A	119.8	O1—C19—C20	115.57 (12)
C3—C4—C5	120.09 (15)	C22—C20—C21	113.37 (12)
C3—C4—H4	120.0	C22—C20—C19	110.50 (11)
C5—C4—H4	120.0	C21—C20—C19	109.60 (12)
C6—C5—C4	119.92 (15)	C22—C20—H20	107.7
C6—C5—H5	120.0	C21—C20—H20	107.7
C4—C5—H5	120.0	C19—C20—H20	107.7
C5—C6—C1	120.48 (14)	O3—C21—C20	106.66 (13)
C5—C6—H6	119.8	O3—C21—H21A	110.4
C1—C6—H6	119.8	C20—C21—H21A	110.4
C12—C7—C8	119.39 (15)	O3—C21—H21B	110.4
C12—C7—Sn1	119.13 (12)	C20—C21—H21B	110.4
C8—C7—Sn1	121.13 (12)	H21A—C21—H21B	108.6
C9—C8—C7	120.32 (17)	C23—C22—C27	118.32 (14)
C9—C8—H8	119.8	C23—C22—C20	122.50 (14)

supplementary materials

C7—C8—H8	119.8	C27—C22—C20	119.17 (14)
C10—C9—C8	119.76 (18)	C22—C23—C24	120.45 (17)
C10—C9—H9	120.1	C22—C23—H23	119.8
C8—C9—H9	120.1	C24—C23—H23	119.8
C11—C10—C9	120.90 (17)	C25—C24—C23	120.65 (18)
C11—C10—H10	119.5	C25—C24—H24	119.7
C9—C10—H10	119.5	C23—C24—H24	119.7
C10—C11—C12	119.91 (19)	C24—C25—C26	119.37 (16)
C10—C11—H11	120.0	C24—C25—H25	120.3
C12—C11—H11	120.0	C26—C25—H25	120.3
C7—C12—C11	119.71 (18)	C25—C26—C27	120.40 (18)
C7—C12—H12	120.1	C25—C26—H26	119.8
C11—C12—H12	120.1	C27—C26—H26	119.8
C18—C13—C14	118.58 (14)	C26—C27—C22	120.79 (17)
C18—C13—Sn1	122.64 (11)	C26—C27—H27	119.6
C14—C13—Sn1	118.67 (11)	C22—C27—H27	119.6
C15—C14—C13	120.55 (16)		
C1—Sn1—O1—C19	67.61 (10)	C7—Sn1—C13—C18	-45.80 (13)
C7—Sn1—O1—C19	-62.58 (11)	O1—Sn1—C13—C14	-99.31 (12)
C13—Sn1—O1—C19	178.80 (10)	C1—Sn1—C13—C14	7.46 (13)
O1—Sn1—C1—C6	17.13 (12)	C7—Sn1—C13—C14	138.15 (12)
C7—Sn1—C1—C6	148.04 (11)	C18—C13—C14—C15	-1.5 (2)
C13—Sn1—C1—C6	-83.32 (12)	Sn1—C13—C14—C15	174.73 (14)
O1—Sn1—C1—C2	-168.17 (11)	C13—C14—C15—C16	0.3 (3)
C7—Sn1—C1—C2	-37.26 (13)	C14—C15—C16—C17	1.1 (3)
C13—Sn1—C1—C2	91.39 (12)	C15—C16—C17—C18	-1.4 (3)
C6—C1—C2—C3	0.7 (2)	C16—C17—C18—C13	0.2 (2)
Sn1—C1—C2—C3	-174.13 (14)	C14—C13—C18—C17	1.2 (2)
C1—C2—C3—C4	0.0 (3)	Sn1—C13—C18—C17	-174.81 (12)
C2—C3—C4—C5	-0.9 (3)	Sn1—O1—C19—O2	11.24 (16)
C3—C4—C5—C6	1.1 (3)	Sn1—O1—C19—C20	-168.49 (9)
C4—C5—C6—C1	-0.4 (2)	O2—C19—C20—C22	93.95 (17)
C2—C1—C6—C5	-0.5 (2)	O1—C19—C20—C22	-86.33 (15)
Sn1—C1—C6—C5	174.18 (11)	O2—C19—C20—C21	-31.69 (19)
O1—Sn1—C7—C12	-116.37 (12)	O1—C19—C20—C21	148.03 (13)
C1—Sn1—C7—C12	119.36 (12)	C22—C20—C21—O3	168.43 (12)
C13—Sn1—C7—C12	-7.63 (14)	C19—C20—C21—O3	-67.59 (15)
O1—Sn1—C7—C8	70.48 (13)	C21—C20—C22—C23	39.8 (2)
C1—Sn1—C7—C8	-53.78 (13)	C19—C20—C22—C23	-83.70 (18)
C13—Sn1—C7—C8	179.23 (11)	C21—C20—C22—C27	-138.98 (15)
C12—C7—C8—C9	0.1 (2)	C19—C20—C22—C27	97.53 (16)
Sn1—C7—C8—C9	173.25 (12)	C27—C22—C23—C24	1.0 (3)
C7—C8—C9—C10	-0.1 (3)	C20—C22—C23—C24	-177.80 (17)
C8—C9—C10—C11	0.3 (3)	C22—C23—C24—C25	-1.1 (3)
C9—C10—C11—C12	-0.5 (3)	C23—C24—C25—C26	-0.1 (3)
C8—C7—C12—C11	-0.3 (2)	C24—C25—C26—C27	1.4 (3)
Sn1—C7—C12—C11	-173.55 (13)	C25—C26—C27—C22	-1.5 (3)
C10—C11—C12—C7	0.5 (3)	C23—C22—C27—C26	0.3 (3)
O1—Sn1—C13—C18	76.74 (12)	C20—C22—C27—C26	179.09 (15)

C1—Sn1—C13—C18 -176.49 (11)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O3—H3···O2 ⁱ	0.84	1.98	2.819 (2)	175

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

Fig. 1

