

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(E)-Ethyl N'-(3,4,5-trimethoxybenzylidene)hydrazinecarboxylate

 Lu-Ping Lv,^a Xiao-Ming Ding,^a Wen-Bo Yu,^a Wei-Wei Li^a and Xian-Chao Hu^{b*}
^aDepartment of Chemical Engineering, Hangzhou Vocational and Technical College, Hangzhou 310018, People's Republic of China, and ^bResearch Center of Analysis and Measurement, Zhejiang University of Technology, Hangzhou 310014, People's Republic of China

Correspondence e-mail: zgdxhc@126.com


Received 13 October 2008; accepted 15 October 2008

 Key indicators: single-crystal X-ray study; $T = 273$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.035; wR factor = 0.094; data-to-parameter ratio = 13.5.

The molecule of the title compound, $\text{C}_{13}\text{H}_{18}\text{N}_2\text{O}_5$, adopts a *trans* configuration with respect to the $\text{C}=\text{N}$ bond. The dihedral angle between the benzene ring and the hydrazine-carboxylic acid plane is 49.75 (5)° and an intramolecular $\text{C}-\text{H}\cdots\text{O}$ interaction occurs. In the crystal structure, the molecules are linked into a chain along [010] by $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, and a $\text{C}-\text{H}\cdots\pi$ contact further stabilizes the structure.

Related literature

For general background, see: Parashar *et al.* (1988); Hadjoudis *et al.* (1987). For a related structure, see: Shang *et al.* (2007).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{18}\text{N}_2\text{O}_5$
 $M_r = 282.29$
 Monoclinic, $P2_1/c$
 $a = 22.037$ (2) Å
 $b = 4.8782$ (5) Å
 $c = 14.0212$ (15) Å
 $\beta = 108.239$ (4)°
 $V = 1431.5$ (3) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 273$ (2) K
 $0.26 \times 0.24 \times 0.22$ mm

Data collection

Bruker SMART CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2002)
 $T_{\min} = 0.965$, $T_{\max} = 0.968$
 14449 measured reflections
 2520 independent reflections
 2108 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.094$
 $S = 1.03$
 2520 reflections
 186 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.17$ e Å⁻³
 $\Delta\rho_{\min} = -0.13$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the C4–C9 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2}-\text{H2}\cdots\text{O4}^i$	0.86	1.98	2.8167 (17)	165
$\text{C2}-\text{H2A}\cdots\text{O1}$	0.96	2.53	3.062 (2)	115
$\text{C1}-\text{H1C}\cdots\text{Cg1}^i$	0.96	2.92	3.755 (2)	146

 Symmetry code: (i) $x, y + 1, z$.

Data collection: SMART (Bruker, 2002); cell refinement: SAINT (Bruker, 2002); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The authors thank Hangzhou Vocational and Technical College, China, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2820).

References

- Bruker (2002). SADABS, SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Hadjoudis, E., Vittorakis, M. & Moustakali-Mavridis, J. (1987). *Tetrahedron*, **43**, 1345–1360.
 Parashar, R. K., Sharma, R. C., Kumar, A. & Mohanm, G. (1988). *Inorg. Chim. Acta*, **151**, 201–208.
 Shang, Z.-H., Zhang, H.-L. & Ding, Y. (2007). *Acta Cryst. E* **63**, o3394.
 Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, o2127 [doi:10.1107/S1600536808033461]

(*E*)-Ethyl *N'*-(3,4,5-trimethoxybenzylidene)hydrazinecarboxylate

L.-P. Lv, X.-M. Ding, W.-B. Yu, W.-W. Li and X.-C. Hu

Comment

Benzaldehydehydrazone derivatives are of interest due to their pharmacological activity (Parashar *et al.*, 1988) and photochromic properties (Hadjoudis *et al.*, 1987). As part of our studies of this type of derivatives, we report herein the crystal structure of the title compound (I).

The title molecule (Fig. 1) adopts a *trans* configuration with respect to the C=N bond. The hydrazine carboxylic acid methyl ester group is slightly twisted away from the attached ring. The dihedral angle between the benzene ring and the C11/C12//N1/N2/O4/O5 plane is 49.75 (5)°. The bond lengths and angles agree with those observed for (*E*)-Methyl *N'*-(4-hydroxybenzylidene)hydrazinecarboxylate (Shang *et al.*, 2007).

The molecules are linked into a chain along [010] by intermolecular and intramolecular N—H···O, C—H···O hydrogen bonds (Fig. 2, Table 1) and a C—H··· π contact further stabilizes the structure.

Experimental

3,4,5-trimethoxybenzaldehyde (1.96 g, 0.01 mol) and methyl hydrazinecarboxylate (1.04 g, 0.01 mol) were dissolved in stirred methanol (25 ml) and left for 3 h at room temperature. The resulting solid was filtered off and recrystallized from ethanol to give the title compound in 90% yield. Colourless blocks of (I) were obtained by slow evaporation of a ethanol solution at room temperature (m.p. 452–454 K).

Refinement

The H atoms were geometrically placed (N—H = 0.86 Å, C—H = 0.93–0.96 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C,N})$ or $1.5U_{\text{eq}}(\text{methyl C})$.

Figures




Fig. 1. The molecular structure of (I), showing 40% probability displacement ellipsoids for the non-hydrogen atoms. The C—H···O interaction is shown as a dashed line.




Fig. 2. Crystal packing of the title compound, viewed approximately down the *a* axis. Dashed lines indicate intermolecular hydrogen bonds.

(E)-Ethyl N'-(3,4,5-trimethoxybenzylidene)hydrazinecarboxylate

Crystal data

$C_{13}H_{18}N_2O_5$

$M_r = 282.29$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 22.037$ (2) Å

$b = 4.8782$ (5) Å

$c = 14.0212$ (15) Å

$\beta = 108.239$ (4)°

$V = 1431.5$ (3) Å³

$Z = 4$

$F_{000} = 600$

$D_x = 1.310$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 2520 reflections

$\theta = 1.0$ – 25.0 °

$\mu = 0.10$ mm⁻¹

$T = 273$ (2) K

Block, colourless

$0.26 \times 0.24 \times 0.22$ mm

Data collection

Bruker SMART CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 273$ (2) K

ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2002)

$T_{\min} = 0.965$, $T_{\max} = 0.968$

14449 measured reflections

2520 independent reflections

2108 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

$\theta_{\max} = 25.1$ °

$\theta_{\min} = 1.0$ °

$h = -24 \rightarrow 25$

$k = -5 \rightarrow 5$

$l = -16 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.035$

$wR(F^2) = 0.094$

$S = 1.03$

2520 reflections

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0431P)^2 + 0.3563P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.17$ e Å⁻³

$\Delta\rho_{\min} = -0.13$ e Å⁻³

186 parameters

Extinction correction: SHELXL97 (Sheldrick, 2008),

$$F_c^* = kFc[1+0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$$

Primary atom site location: structure-invariant direct methods

Extinction coefficient: 0.0159 (15)

Secondary atom site location: difference Fourier map

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	U_{iso}^*/U_{eq}
C7	0.19238 (7)	0.8025 (3)	0.61046 (10)	0.0417 (3)
H7	0.1961	0.9206	0.6641	0.050*
C9	0.24447 (7)	0.6457 (3)	0.60759 (10)	0.0399 (3)
C4	0.13069 (7)	0.6200 (3)	0.44959 (10)	0.0407 (3)
C8	0.23981 (7)	0.4730 (3)	0.52645 (10)	0.0446 (4)
H8	0.2743	0.3643	0.5255	0.054*
C6	0.13492 (7)	0.7831 (3)	0.53350 (10)	0.0400 (3)
C11	0.44480 (7)	0.3407 (3)	0.82320 (10)	0.0423 (3)
C5	0.18340 (7)	0.4648 (3)	0.44715 (10)	0.0439 (4)
C10	0.30489 (7)	0.6728 (3)	0.68892 (10)	0.0424 (3)
H10	0.3141	0.8365	0.7245	0.051*
C3	0.08165 (8)	1.0795 (4)	0.61651 (12)	0.0587 (4)
H3A	0.0958	0.9706	0.6764	0.088*
H3B	0.0396	1.1488	0.6086	0.088*
H3C	0.1105	1.2300	0.6215	0.088*
C12	0.55019 (7)	0.2692 (4)	0.93180 (12)	0.0548 (4)
H12A	0.5554	0.1264	0.8869	0.066*
H12B	0.5409	0.1838	0.9881	0.066*
C2	0.03622 (8)	0.3905 (3)	0.35788 (13)	0.0584 (4)
H2A	0.0603	0.2298	0.3535	0.088*
H2B	0.0004	0.4072	0.2979	0.088*
H2C	0.0212	0.3755	0.4150	0.088*
C1	0.22489 (9)	0.1433 (4)	0.35569 (13)	0.0716 (6)
H1A	0.2618	0.2520	0.3588	0.107*
H1B	0.2117	0.0432	0.2936	0.107*
H1C	0.2353	0.0171	0.4110	0.107*
C13	0.60936 (8)	0.4386 (4)	0.96774 (14)	0.0681 (5)
H13A	0.6185	0.5183	0.9112	0.102*

supplementary materials

H13B	0.6445	0.3248	1.0044	0.102*
H13C	0.6031	0.5816	1.0108	0.102*
O5	0.49926 (5)	0.4522 (2)	0.87979 (8)	0.0497 (3)
O2	0.07592 (5)	0.6268 (2)	0.36863 (7)	0.0485 (3)
O3	0.08020 (5)	0.9148 (2)	0.53188 (7)	0.0511 (3)
O1	0.17482 (5)	0.3161 (3)	0.36105 (8)	0.0617 (3)
O4	0.43507 (6)	0.0998 (2)	0.81180 (10)	0.0752 (4)
N2	0.40242 (6)	0.5408 (2)	0.78322 (9)	0.0447 (3)
H2	0.4111	0.7079	0.8021	0.054*
N1	0.34512 (6)	0.4774 (2)	0.71207 (8)	0.0425 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C7	0.0445 (8)	0.0412 (8)	0.0355 (7)	-0.0013 (6)	0.0070 (6)	-0.0019 (6)
C9	0.0396 (8)	0.0395 (8)	0.0365 (7)	-0.0036 (6)	0.0062 (6)	0.0028 (6)
C4	0.0411 (8)	0.0409 (8)	0.0333 (7)	0.0000 (6)	0.0020 (6)	0.0044 (6)
C8	0.0439 (8)	0.0452 (8)	0.0416 (8)	0.0037 (7)	0.0087 (7)	0.0005 (6)
C6	0.0402 (8)	0.0389 (8)	0.0390 (7)	0.0016 (6)	0.0093 (6)	0.0043 (6)
C11	0.0454 (9)	0.0363 (8)	0.0398 (7)	-0.0023 (6)	0.0055 (6)	-0.0018 (6)
C5	0.0495 (9)	0.0442 (8)	0.0338 (7)	0.0021 (7)	0.0071 (6)	-0.0012 (6)
C10	0.0418 (8)	0.0418 (8)	0.0402 (7)	-0.0041 (7)	0.0077 (6)	-0.0022 (6)
C3	0.0558 (10)	0.0653 (11)	0.0527 (9)	0.0110 (8)	0.0137 (8)	-0.0094 (8)
C12	0.0469 (9)	0.0577 (10)	0.0524 (9)	0.0147 (8)	0.0048 (7)	0.0007 (8)
C2	0.0537 (10)	0.0530 (10)	0.0546 (9)	-0.0050 (8)	-0.0031 (8)	0.0000 (8)
C1	0.0766 (13)	0.0769 (13)	0.0560 (10)	0.0208 (10)	0.0130 (9)	-0.0177 (9)
C13	0.0409 (9)	0.0941 (14)	0.0626 (10)	0.0069 (9)	0.0064 (8)	-0.0054 (10)
O5	0.0398 (6)	0.0418 (6)	0.0562 (6)	0.0018 (4)	-0.0014 (5)	-0.0038 (5)
O2	0.0477 (6)	0.0469 (6)	0.0386 (5)	0.0012 (5)	-0.0043 (4)	0.0037 (4)
O3	0.0443 (6)	0.0595 (7)	0.0434 (5)	0.0111 (5)	0.0048 (5)	-0.0053 (5)
O1	0.0615 (7)	0.0723 (8)	0.0408 (6)	0.0176 (6)	0.0010 (5)	-0.0162 (5)
O4	0.0734 (8)	0.0320 (6)	0.0930 (9)	-0.0032 (6)	-0.0131 (7)	-0.0022 (6)
N2	0.0404 (7)	0.0334 (6)	0.0491 (7)	-0.0041 (5)	-0.0020 (6)	-0.0040 (5)
N1	0.0397 (7)	0.0410 (7)	0.0396 (6)	-0.0060 (5)	0.0019 (5)	-0.0008 (5)

Geometric parameters (\AA , $^\circ$)

C7—C6	1.3862 (19)	C3—H3B	0.9600
C7—C9	1.391 (2)	C3—H3C	0.9600
C7—H7	0.9300	C12—O5	1.4409 (17)
C9—C8	1.393 (2)	C12—C13	1.492 (2)
C9—C10	1.4630 (19)	C12—H12A	0.9700
C4—O2	1.3744 (16)	C12—H12B	0.9700
C4—C5	1.396 (2)	C2—O2	1.4264 (19)
C4—C6	1.399 (2)	C2—H2A	0.9600
C8—C5	1.385 (2)	C2—H2B	0.9600
C8—H8	0.9300	C2—H2C	0.9600
C6—O3	1.3602 (17)	C1—O1	1.409 (2)
C11—O4	1.1965 (18)	C1—H1A	0.9600

C11—O5	1.3303 (17)	C1—H1B	0.9600
C11—N2	1.3457 (18)	C1—H1C	0.9600
C5—O1	1.3697 (17)	C13—H13A	0.9600
C10—N1	1.2726 (18)	C13—H13B	0.9600
C10—H10	0.9300	C13—H13C	0.9600
C3—O3	1.4250 (18)	N2—N1	1.3772 (16)
C3—H3A	0.9600	N2—H2	0.8600
C6—C7—C9	120.07 (13)	O5—C12—H12A	110.4
C6—C7—H7	120.0	C13—C12—H12A	110.4
C9—C7—H7	120.0	O5—C12—H12B	110.4
C7—C9—C8	120.37 (13)	C13—C12—H12B	110.4
C7—C9—C10	119.07 (13)	H12A—C12—H12B	108.6
C8—C9—C10	120.51 (13)	O2—C2—H2A	109.5
O2—C4—C5	121.02 (12)	O2—C2—H2B	109.5
O2—C4—C6	119.39 (13)	H2A—C2—H2B	109.5
C5—C4—C6	119.49 (12)	O2—C2—H2C	109.5
C5—C8—C9	119.44 (14)	H2A—C2—H2C	109.5
C5—C8—H8	120.3	H2B—C2—H2C	109.5
C9—C8—H8	120.3	O1—C1—H1A	109.5
O3—C6—C7	124.73 (13)	O1—C1—H1B	109.5
O3—C6—C4	115.43 (12)	H1A—C1—H1B	109.5
C7—C6—C4	119.84 (13)	O1—C1—H1C	109.5
O4—C11—O5	124.87 (14)	H1A—C1—H1C	109.5
O4—C11—N2	125.79 (14)	H1B—C1—H1C	109.5
O5—C11—N2	109.33 (12)	C12—C13—H13A	109.5
O1—C5—C8	124.38 (14)	C12—C13—H13B	109.5
O1—C5—C4	114.98 (12)	H13A—C13—H13B	109.5
C8—C5—C4	120.61 (13)	C12—C13—H13C	109.5
N1—C10—C9	121.50 (13)	H13A—C13—H13C	109.5
N1—C10—H10	119.2	H13B—C13—H13C	109.5
C9—C10—H10	119.2	C11—O5—C12	117.58 (12)
O3—C3—H3A	109.5	C4—O2—C2	114.74 (11)
O3—C3—H3B	109.5	C6—O3—C3	117.63 (11)
H3A—C3—H3B	109.5	C5—O1—C1	117.99 (12)
O3—C3—H3C	109.5	C11—N2—N1	119.88 (12)
H3A—C3—H3C	109.5	C11—N2—H2	120.1
H3B—C3—H3C	109.5	N1—N2—H2	120.1
O5—C12—C13	106.72 (14)	C10—N1—N2	115.09 (12)
C6—C7—C9—C8	1.5 (2)	C7—C9—C10—N1	153.39 (14)
C6—C7—C9—C10	179.09 (13)	C8—C9—C10—N1	-29.0 (2)
C7—C9—C8—C5	1.9 (2)	O4—C11—O5—C12	-0.4 (2)
C10—C9—C8—C5	-175.66 (13)	N2—C11—O5—C12	178.13 (13)
C9—C7—C6—O3	175.98 (13)	C13—C12—O5—C11	166.29 (13)
C9—C7—C6—C4	-4.5 (2)	C5—C4—O2—C2	77.45 (18)
O2—C4—C6—O3	7.15 (19)	C6—C4—O2—C2	-106.16 (16)
C5—C4—C6—O3	-176.39 (13)	C7—C6—O3—C3	-1.6 (2)
O2—C4—C6—C7	-172.45 (13)	C4—C6—O3—C3	178.83 (14)
C5—C4—C6—C7	4.0 (2)	C8—C5—O1—C1	5.2 (2)

supplementary materials

C9—C8—C5—O1	175.76 (14)	C4—C5—O1—C1	-176.66 (15)
C9—C8—C5—C4	-2.3 (2)	O4—C11—N2—N1	-9.3 (2)
O2—C4—C5—O1	-2.5 (2)	O5—C11—N2—N1	172.11 (11)
C6—C4—C5—O1	-178.86 (13)	C9—C10—N1—N2	173.88 (12)
O2—C4—C5—C8	175.78 (13)	C11—N2—N1—C10	170.86 (13)
C6—C4—C5—C8	-0.6 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H2 \cdots O4 ⁱ	0.86	1.98	2.8167 (17)	165
C2—H2A \cdots O1	0.96	2.53	3.062 (2)	115
C1—H1C \cdots Cg1 ⁱ	0.96	2.92	3.755 (2)	146

Symmetry codes: (i) $x, y+1, z$.

Fig. 1




Fig. 2

