

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1,3-Bis(4-methoxyphenyl)imidazolidium chloride monohydrate

Yu Wan, Haiqiang Xin, Xiumei Chen, Huahong Xu and Hui Wu*

School of Chemistry and Chemical Engineering, Xuzhou Normal University, Xuzhou 221116, People's Republic of China

Correspondence e-mail: wuhui72@yahoo.com.cn


Received 27 September 2008; accepted 17 October 2008

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.029; wR factor = 0.076; data-to-parameter ratio = 7.0.

The asymmetric unit of the title compound, $\text{C}_{17}\text{H}_{17}\text{N}_2\text{O}_2^+\cdot\text{Cl}^-\cdot\text{H}_2\text{O}$, contains one-half of the cation, one-half of a water molecule and a chloride anion. The complete cation is generated by crystallographic two-fold symmetry, with one C atom lying on the rotation axis. The O and Cl atoms have site symmetry 2. The imidazolidium ring is oriented at a dihedral angle of $4.15(3)^\circ$ with respect to the 4-methoxyphenyl ring and an intramolecular $\text{C}-\text{H}\cdots\text{O}$ interaction occurs. In the crystal structure, intermolecular $\text{O}-\text{H}\cdots\text{Cl}$ and $\text{C}-\text{H}\cdots\text{Cl}$ hydrogen bonds link the molecules. There is a $\pi-\pi$ contact between the imidazolidium and 4-methoxyphenyl rings [centroid-to-centroid distance = $3.625(3)$ Å]. There is also a $\text{C}-\text{H}\cdots\pi$ contact between the methyl group and the 4-methoxyphenyl ring.

Related literature

For general background, see: Lin & Vasam (2005). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{17}\text{N}_2\text{O}_2^+\cdot\text{Cl}^-\cdot\text{H}_2\text{O}$
 $M_r = 334.79$
 Monoclinic, $C2$
 $a = 15.6706(19)$ Å
 $b = 9.4198(9)$ Å
 $c = 5.4026(4)$ Å
 $\beta = 90.156(1)^\circ$

$V = 797.50(14)$ Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.26$ mm⁻¹
 $T = 298(2)$ K
 $0.20 \times 0.11 \times 0.09$ mm

Data collection

Bruker SMART CCD area-detector diffractometer
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.951$, $T_{\max} = 0.977$
 2026 measured reflections
 749 independent reflections
 688 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.029$
 $wR(F^2) = 0.076$
 $S = 1.01$
 749 reflections
 107 parameters
 1 restraint
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.11$ e Å⁻³
 $\Delta\rho_{\min} = -0.22$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

 $Cg2$ is the centroid of the C3–C8 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O2}-\text{H2}\cdots\text{Cl1}$	0.85	2.29	3.133 (3)	173
$\text{C1}-\text{H1}\cdots\text{O2}$	0.93	2.13	3.060 (3)	180
$\text{C2}-\text{H2A}\cdots\text{Cl1}^{\text{i}}$	0.93	2.69	3.474 (3)	142
$\text{C4}-\text{H4}\cdots\text{O2}$	0.93	2.47	3.391 (3)	170
$\text{C9}-\text{H9C}\cdots\text{Cg2}^{\text{ii}}$	0.96	2.91	3.629 (3)	133

 Symmetry codes: (i) $x + \frac{1}{2}, y - \frac{1}{2}, z$; (ii) $x, y, z + 1$.

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINTE* (Bruker, 1999); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2003); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008) and *PLATON*.

The authors are grateful to the National Natural Science Foundation of China (grant No. 20772103), the Natural Science Foundation of Jiangsu Province (grant No. BK2007028) and the Surpassing Project of Jiangsu Province (grant No. CX07S_016z) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2542).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
 Bruker (1998). *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (1999). *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Lin, I. J. B. & Vasam, C. S. (2005). *J. Organomet. Chem.* **690**, 3498–3512.
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2008). E64, o2159 [doi:10.1107/S1600536808033965]

1,3-Bis(4-methoxyphenyl)imidazolidium chloride monohydrate

Y. Wan, H. Xin, X. Chen, H. Xu and H. Wu

Comment

Imidazole and its derivatives such as imidazolium cation are important compounds playing important roles in medical, organic and material chemistry (Lin & Vasam, 2005). A broad application of imidazolium now is to synthesize ionic liquids. Recently, ionic liquids are attracting much attention as alternative reaction media for synthesis and catalysis. Its applications in many different areas including separation processes, catalyst, electrochemistry, electrolytes in solar cells and lubricants are widely recognized. Therefore, the need of ionic liquids with specific chemical and physical properties become stronger. We report herein the synthesis and crystal structure of the title compound.

The asymmetric unit of the title compound (Fig. 1) contains one half-molecule, one half-water molecule and a chloride atom. The bond lengths (Allen *et al.*, 1987) and angles are within normal ranges. Rings A (N1/N1'/C1/C2/C2') and B (C3–C8) are, of course, planar and the dihedral angle between them is $A/B = 4.15(3)^\circ$ [symmetry code: (') -x, y, -z]. Intramolecular C—H \cdots O and O—H \cdots Cl hydrogen bonds (Table 1) link the molecules.

In the crystal structure, intramolecular C—H \cdots O and O—H \cdots Cl and intermolecular C—H \cdots Cl hydrogen bonds (Table 1) link the molecules (Fig. 2), in which they may be effective in the stabilization of the structure. The π – π contact between the imidazolidium and 4-methoxyphenyl rings, Cg1 \cdots Cg2ⁱ [symmetry code: (i) 1 - x, y, 1 - z, where Cg1 and Cg2 are the centroids of the rings A (N1/N1'/C1/C2/C2') and B (C3–C8), respectively] may further stabilize the structure, with centroid-centroid distance of 3.625(3) Å. There also exist a C—H \cdots π contact (Table 1) between the methyl group and the 4-methoxyphenyl ring.

Experimental

The reaction of 4-methoxybenzenamine (2 mmol) with formaldehyde (aq. 37%, 1 mmol) and glyoxal (aq. 40%, 1 mmol) in ethanol (95%) at 273–278 K for 8 h afforded 1-(2,3-diethoxy-4-(4-methoxyphenyl)cyclopentyl)-4-methoxybenzene (yield; 89%). The title compound was obtained through the oxidization of 1-(2,3-diethoxy-4-(4-methoxyphenyl)cyclopentyl)-4-methoxybenzene by phosgene in DMF at 268–273 K (yield 95%).

Refinement

H atoms were positioned geometrically, with O—H = 0.85 Å (for H₂O) and C—H = 0.93 and 0.96 Å for aromatic and methyl H, respectively, and constrained to ride on their parent atoms with $U_{iso}(H) = xU_{eq}(C,O)$, where $x = 1.5$ for methyl H and $x = 1.2$ for all other H atoms.

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme [symmetry code: (') $-x, y, -z$].




Fig. 2. A partial packing diagram. Hydrogen bonds are shown as dashed lines.

1,3-Bis(4-methoxyphenyl)imidazolidium chloride monohydrate

Crystal data

$C_{17}H_{17}N_2O_2^+ \cdot Cl^- \cdot H_2O$

$M_r = 334.79$

Monoclinic, $C2$

Hall symbol: $-C\ 2y$

$a = 15.6706$ (19) Å

$b = 9.4198$ (9) Å

$c = 5.4026$ (4) Å

$\beta = 90.156$ (1)°

$V = 797.50$ (14) Å³

$Z = 2$

$F_{000} = 352$

$D_x = 1.394$ Mg m⁻³

Melting point = 492–494 K

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 1340 reflections

$\theta = 2.5$ – 28.3 °

$\mu = 0.26$ mm⁻¹

$T = 298$ (2) K

Block, colourless

$0.20 \times 0.11 \times 0.09$ mm

Data collection

Bruker SMART CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 298$ (2) K

φ and ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.951$, $T_{\max} = 0.977$

2026 measured reflections

749 independent reflections

688 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.021$

$\theta_{\max} = 25.0$ °

$\theta_{\min} = 2.5$ °

$h = -18 \rightarrow 13$

$k = -9 \rightarrow 11$

$l = -6 \rightarrow 6$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.029$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$wR(F^2) = 0.076$	$w = 1/[\sigma^2(F_o^2) + (0.0503P)^2 + 0.1521P]$
$S = 1.01$	where $P = (F_o^2 + 2F_c^2)/3$
749 reflections	$(\Delta/\sigma)_{\max} < 0.001$
107 parameters	$\Delta\rho_{\max} = 0.11 \text{ e } \text{\AA}^{-3}$
1 restraint	$\Delta\rho_{\min} = -0.22 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.5000	0.84927 (12)	0.5000	0.0800 (5)
O1	0.24434 (13)	0.4740 (2)	0.9080 (3)	0.0519 (5)
O2	0.5000	0.6808 (3)	0.0000	0.0567 (8)
H2	0.5000	0.7335	0.1277	0.068*
N1	0.45802 (12)	0.27396 (19)	0.1594 (3)	0.0334 (5)
C1	0.5000	0.3560 (4)	0.0000	0.0350 (7)
H1	0.5000	0.4547	0.0000	0.042*
C2	0.47435 (17)	0.1342 (3)	0.0983 (5)	0.0444 (6)
H2A	0.4534	0.0545	0.1795	0.053*
C3	0.40340 (14)	0.3225 (3)	0.3567 (4)	0.0335 (5)
C4	0.39574 (17)	0.4670 (3)	0.4036 (5)	0.0420 (6)
H4	0.4262	0.5322	0.3094	0.050*
C5	0.34270 (17)	0.5135 (3)	0.5903 (5)	0.0449 (6)
H5	0.3378	0.6101	0.6225	0.054*
C6	0.29642 (15)	0.4164 (3)	0.7309 (5)	0.0386 (6)
C7	0.30558 (17)	0.2728 (3)	0.6859 (5)	0.0446 (6)
H7	0.2758	0.2073	0.7812	0.054*
C8	0.35934 (16)	0.2262 (3)	0.4979 (5)	0.0440 (6)
H8	0.3654	0.1296	0.4678	0.053*
C9	0.19340 (19)	0.3780 (4)	1.0497 (5)	0.0580 (8)
H9A	0.1588	0.3219	0.9402	0.087*
H9B	0.1573	0.4309	1.1594	0.087*
H9C	0.2299	0.3168	1.1447	0.087*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.1591 (13)	0.0423 (5)	0.0387 (5)	0.000	0.0069 (6)	0.000
O1	0.0545 (11)	0.0522 (12)	0.0491 (11)	0.0011 (9)	0.0158 (9)	0.0059 (9)
O2	0.092 (2)	0.0344 (15)	0.0434 (14)	0.000	0.0109 (14)	0.000
N1	0.0365 (11)	0.0278 (10)	0.0358 (10)	-0.0015 (8)	-0.0027 (8)	0.0024 (8)
C1	0.0400 (18)	0.0271 (15)	0.0379 (15)	0.000	0.0009 (14)	0.000
C2	0.0572 (16)	0.0300 (13)	0.0461 (13)	-0.0031 (11)	0.0030 (11)	0.0012 (11)
C3	0.0319 (12)	0.0362 (13)	0.0323 (10)	-0.0010 (10)	-0.0018 (9)	0.0024 (10)
C4	0.0473 (15)	0.0333 (14)	0.0455 (14)	-0.0024 (11)	0.0086 (11)	0.0080 (11)
C5	0.0510 (15)	0.0339 (13)	0.0499 (14)	0.0029 (12)	0.0085 (11)	0.0012 (12)
C6	0.0356 (14)	0.0442 (15)	0.0360 (12)	-0.0008 (11)	0.0001 (11)	0.0027 (11)
C7	0.0480 (15)	0.0421 (16)	0.0438 (14)	-0.0081 (12)	0.0045 (11)	0.0072 (12)
C8	0.0518 (16)	0.0319 (13)	0.0484 (15)	-0.0044 (12)	0.0012 (13)	0.0019 (12)
C9	0.0485 (16)	0.069 (2)	0.0566 (16)	-0.0036 (14)	0.0137 (13)	0.0109 (15)

Geometric parameters (\AA , $^\circ$)

O1—C6	1.371 (3)	C4—C5	1.380 (4)
O1—C9	1.430 (3)	C4—H4	0.9300
O2—H2	0.8500	C5—C6	1.394 (4)
N1—C1	1.332 (3)	C5—H5	0.9300
N1—C2	1.382 (3)	C6—C7	1.382 (4)
N1—C3	1.443 (3)	C7—C8	1.392 (4)
C1—N1 ⁱ	1.332 (3)	C7—H7	0.9300
C1—H1	0.9300	C8—H8	0.9300
C2—C2 ⁱ	1.334 (5)	C9—H9A	0.9600
C2—H2A	0.9300	C9—H9B	0.9600
C3—C8	1.372 (3)	C9—H9C	0.9600
C3—C4	1.390 (4)		
C6—O1—C9	117.2 (2)	C4—C5—H5	119.8
C1—N1—C2	107.8 (2)	C6—C5—H5	119.8
C1—N1—C3	126.1 (2)	O1—C6—C7	124.9 (2)
C2—N1—C3	126.1 (2)	O1—C6—C5	115.6 (2)
N1 ⁱ —C1—N1	109.1 (3)	C7—C6—C5	119.5 (2)
N1 ⁱ —C1—H1	125.5	C6—C7—C8	120.0 (2)
N1—C1—H1	125.5	C6—C7—H7	120.0
C2 ⁱ —C2—N1	107.63 (14)	C8—C7—H7	120.0
C2 ⁱ —C2—H2A	126.2	C3—C8—C7	120.2 (2)
N1—C2—H2A	126.2	C3—C8—H8	119.9
C8—C3—C4	120.1 (2)	C7—C8—H8	119.9
C8—C3—N1	120.1 (2)	O1—C9—H9A	109.5
C4—C3—N1	119.8 (2)	O1—C9—H9B	109.5
C5—C4—C3	119.8 (2)	H9A—C9—H9B	109.5
C5—C4—H4	120.1	O1—C9—H9C	109.5

C3—C4—H4	120.1	H9A—C9—H9C	109.5
C4—C5—C6	120.4 (2)	H9B—C9—H9C	109.5
C2—N1—C1—N1 ⁱ	0.12 (13)	C3—C4—C5—C6	-0.4 (4)
C3—N1—C1—N1 ⁱ	-178.4 (2)	C9—O1—C6—C7	-2.5 (4)
C1—N1—C2—C2 ⁱ	-0.3 (3)	C9—O1—C6—C5	177.7 (2)
C3—N1—C2—C2 ⁱ	178.2 (2)	C4—C5—C6—O1	-178.7 (2)
C1—N1—C3—C8	175.65 (18)	C4—C5—C6—C7	1.5 (4)
C2—N1—C3—C8	-2.6 (3)	O1—C6—C7—C8	178.9 (2)
C1—N1—C3—C4	-4.5 (3)	C5—C6—C7—C8	-1.3 (4)
C2—N1—C3—C4	177.3 (2)	C4—C3—C8—C7	1.1 (3)
C8—C3—C4—C5	-0.9 (4)	N1—C3—C8—C7	-179.0 (2)
N1—C3—C4—C5	179.2 (2)	C6—C7—C8—C3	0.0 (4)

Symmetry codes: (i) $-x+1, y, -z$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H2 \cdots C11	0.85	2.29	3.133 (3)	173
C1—H1 \cdots O2	0.93	2.13	3.060 (3)	180
C2—H2A \cdots C11 ⁱⁱ	0.93	2.69	3.474 (3)	142
C4—H4 \cdots O2	0.93	2.47	3.391 (3)	170
C9—H9C \cdots Cg2 ⁱⁱⁱ	0.96	2.91	3.629 (3)	133

Symmetry codes: (ii) $x+1/2, y-1/2, z$; (iii) $x, y, z+1$.

Fig. 1




Fig. 2

