

## (Cyanomethyl)triphenylphosphonium chloride

Muhammad Shafiq,<sup>a</sup> M. Nawaz Tahir,<sup>b\*</sup> Islam Ullah Khan,<sup>a</sup> Muhammad Nadeem Arshad<sup>a</sup> and Zaib-un-Nisa<sup>a</sup>

<sup>a</sup>Government College University, Department of Chemistry, Lahore, Pakistan, and  
<sup>b</sup>University of Sargodha, Department of Physics, Sargodha, Pakistan

Correspondence e-mail: dmntahir\_uos@yahoo.com


Received 21 October 2008; accepted 23 October 2008

Key indicators: single-crystal X-ray study;  $T = 296\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ;  $R$  factor = 0.041;  $wR$  factor = 0.111; data-to-parameter ratio = 21.5.

In the molecule of the title compound,  $\text{C}_{20}\text{H}_{17}\text{NP}^+\cdot\text{Cl}^-$ , the coordination around the P atom is slightly distorted tetrahedral. In the crystal structure, intermolecular  $\text{C}-\text{H}\cdots\text{N}$  and  $\text{C}-\text{H}\cdots\text{Cl}$  hydrogen bonds link the molecules. There is a  $\pi-\pi$  contact between the phenyl rings [centroid–centroid distance = 3.702 (3)  $\text{\AA}$ ].

### Related literature

For related structures, see: Czerwinski (2004); Czerwinski & Ponnuswamy (1988); de Dubourg *et al.* (1986); Fischer & Wiebelhaus (1997); Shafiq *et al.* (2008); Skapski & Stephens (1974); Tahir *et al.* (2008).



### Experimental

#### Crystal data



$M_r = 337.77$

Monoclinic,  $P2_1/n$

$a = 11.8269(5)\text{ \AA}$

$b = 11.8130(4)\text{ \AA}$

$c = 12.8918(5)\text{ \AA}$

$\beta = 92.213(2)^\circ$

$V = 1799.79(12)\text{ \AA}^3$

$Z = 4$

Mo  $K\alpha$  radiation

$\mu = 0.30\text{ mm}^{-1}$

$T = 296(2)\text{ K}$

$0.26 \times 0.20 \times 0.16\text{ mm}$

#### Data collection

Bruker Kappa APEXII CCD diffractometer

Absorption correction: multi-scan (*SADABS*; Bruker, 2005)

$T_{\min} = 0.928$ ,  $T_{\max} = 0.950$

19927 measured reflections

4465 independent reflections

3145 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.034$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$

$wR(F^2) = 0.111$

$S = 1.03$

4465 reflections

208 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.37\text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.31\text{ e \AA}^{-3}$

**Table 1**  
Selected geometric parameters ( $\text{\AA}$ ,  $^\circ$ ).

P1—C1	1.7923 (18)	P1—C13	1.7851 (17)
P1—C7	1.7845 (18)	P1—C19	1.8046 (17)
C1—P1—C7	111.03 (8)	C7—P1—C13	110.71 (8)
C1—P1—C13	109.26 (8)	C7—P1—C19	106.81 (8)
C1—P1—C19	108.56 (8)	C13—P1—C19	110.43 (8)

**Table 2**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C12—H12 $\cdots$ Cl1	0.93	2.66	3.479 (2)	147
C17—H17 $\cdots$ N1 <sup>i</sup>	0.93	2.61	3.530 (3)	171
C19—H19A $\cdots$ Cl1 <sup>ii</sup>	0.97	2.34	3.3076 (17)	173
C19—H19B $\cdots$ Cl1 <sup>iii</sup>	0.97	2.46	3.3830 (19)	160

Symmetry codes: (i)  $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$ ; (ii)  $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$ ; (iii)  $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$ ; (iv)  $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$ .

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2003); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

MS greatfully acknowledges the Higher Education Commission, Islamabad, Pakistan, for providing his Scholarship under the Indigenous PhD Program (PIN 042-120567-PS2-276).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2558).

### References

- Bruker (2005). *SADABS*. Bruker AXS Inc. Madison, Wisconsin, USA.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc. Madison, Wisconsin, USA.
- Czerwinski, E. W. (2004). *Acta Cryst. E60*, o1442–o1443.
- Czerwinski, E. W. & Ponnuswamy, M. N. (1988). *Acta Cryst. C44*, 862–865.
- Dubourg, A., De Castro Dantas, T. N., Klaébé, A. & Declercq, J.-P. (1986). *Acta Cryst. C42*, 112–114.
- Farrugia, L. J. (1997). *J. Appl. Cryst. 30*, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst. 32*, 837–838.
- Fischer, A. & Wiebelhaus, D. (1997). *Z. Kristallogr. New Cryst. Struct. 212*, 335–336.
- Shafiq, M., Khan, I. U., Tahir, M. N. & Siddiqui, W. A. (2008). *Acta Cryst. E64*, o558.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Skapski, A. C. & Stephens, F. A. (1974). *J. Cryst. Mol. Struct. 4*, 77–85.
- Spek, A. L. (2003). *J. Appl. Cryst. 36*, 7–13.
- Tahir, M. N., Shafiq, M., Khan, I. U., Siddiqui, W. A. & Arshad, M. N. (2008). *Acta Cryst. E64*, o557.

# supporting information

*Acta Cryst.* (2008). E64, o2213 [doi:10.1107/S1600536808034673]

## (Cyanomethyl)triphenylphosphonium chloride

**Muhammad Shafiq, M. Nawaz Tahir, Islam Ullah Khan, Muhammad Nadeem Arshad and Zaib-un-Nisa**

### S1. Comment

Triphenyl phosphonium compounds are key reagents in the Wittig reactions, used to convert aldehydes and ketones into alkenes. The Wittig reaction has seen use in applications ranging from the synthesis of simple alkenes to the construction of complex biologically active molecules for the pharmaceutical industry. The title compound is synthesized for the derivatization of our already published structures (Shafiq *et al.*, 2008; Tahir *et al.*, 2008) using this particular reaction. Various structures have been published having the similar geometry around P atom (Skapski & Stephens, 1974; de Dubourg *et al.*, 1986; Czerwinski & Ponnuswamy, 1988; Fischer & Wiebelhaus, 1997; Czerwinski, 2004).

In the molecule of the title compound (Fig 1), the geometry around P atom is slightly distorted tetrahedral (Table 1). Rings A (C1-C6), B (C7-C12) and C (C13-C18) are of course planar. The dihedral angles between them are A/B = 86.10 (11) $^{\circ}$ , A/C = 89.78 (10) $^{\circ}$  and B/C = 76.23 (12) $^{\circ}$ .


In the crystal structure, intramolecular C-H $\cdots$ Cl and intermolecular C-H $\cdots$ N and C-H $\cdots$ Cl hydrogen bonds (Table 2) link the molecules (Fig. 2), in which they may be effective in the stabilization of the structure. The  $\pi$ - $\pi$  contact between the phenyl rings, Cg3 $\cdots$ Cg3<sup>i</sup> [symmetry code: (i) 2 - x, -y, 1 - z, where Cg3 is the centroid of the ring C (C13-C18)] may further stabilize the structure, with centroid-centroid distance of 3.702 (3) Å. There also exist a C—H $\cdots$  $\pi$  contact (Table 2) between the phenyl rings.

### S2. Experimental


Triphenylphosphine (10 g, 0.038 mol) was dissolved in benzene (20 ml) under stirring at room temperature. To this solution, chloroacetonitrile (4 g, 0.0514 mole) was added dropwise. After complete addition, clear solution formed was left in the darkness for 2-3 d. Colorless crystals formed were separated for X-ray diffraction studies.

### S3. Refinement

H-atoms were positioned geometrically, with C-H = 0.93 and 0.97 Å for aromatic and methylene H, and constrained to ride on their parent atoms, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

**Figure 1**

The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids drawn at the 50% probability level.

**Figure 2**

A partial packing diagram. Hydrogen bonds are shown as dashed lines.

### (Cyanomethyl)triphenylphosphonium chloride

#### *Crystal data*

$C_{20}H_{17}NP^+\cdot Cl^-$   
 $M_r = 337.77$   
Monoclinic,  $P2_1/n$   
Hall symbol: -P 2yn  
 $a = 11.8269 (5) \text{ \AA}$   
 $b = 11.8130 (4) \text{ \AA}$   
 $c = 12.8918 (5) \text{ \AA}$   
 $\beta = 92.213 (2)^\circ$   
 $V = 1799.79 (12) \text{ \AA}^3$   
 $Z = 4$

$F(000) = 704$   
 $D_x = 1.247 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$   
Cell parameters from 4467 reflections  
 $\theta = 2.3\text{--}28.3^\circ$   
 $\mu = 0.30 \text{ mm}^{-1}$   
 $T = 296 \text{ K}$   
Prismatic, colorless  
 $0.26 \times 0.20 \times 0.16 \text{ mm}$

#### *Data collection*

Bruker KappaAPEXII CCD  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator

Detector resolution: 7.40 pixels  $\text{mm}^{-1}$   
 $\omega$  scans  
Absorption correction: multi-scan  
(SADABS; Bruker, 2005)

$T_{\min} = 0.928$ ,  $T_{\max} = 0.950$   
 19927 measured reflections  
 4465 independent reflections  
 3145 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.034$

$\theta_{\max} = 28.3^\circ$ ,  $\theta_{\min} = 2.3^\circ$   
 $h = -15 \rightarrow 15$   
 $k = -10 \rightarrow 15$   
 $l = -17 \rightarrow 14$

#### Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.041$   
 $wR(F^2) = 0.111$   
 $S = 1.03$   
 4465 reflections  
 208 parameters  
 0 restraints  
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map  
 Hydrogen site location: inferred from neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0464P)^2 + 0.4839P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.37 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.31 \text{ e } \text{\AA}^{-3}$

#### Special details

**Geometry.** Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.64049 (5)	0.13296 (4)	0.08789 (5)	0.0652 (2)
P1	0.81585 (4)	0.28430 (4)	0.41401 (3)	0.0347 (1)
N1	0.7328 (2)	0.38278 (18)	0.67432 (16)	0.0773 (8)
C1	0.66854 (15)	0.30928 (15)	0.38497 (13)	0.0396 (5)
C2	0.59647 (17)	0.22052 (19)	0.35858 (15)	0.0539 (7)
C3	0.48330 (19)	0.2418 (3)	0.33767 (18)	0.0708 (9)
C4	0.4421 (2)	0.3502 (3)	0.34392 (19)	0.0758 (9)
C5	0.51305 (19)	0.4389 (2)	0.36860 (17)	0.0674 (8)
C6	0.62701 (17)	0.41952 (18)	0.38895 (15)	0.0530 (7)
C7	0.89873 (15)	0.31735 (14)	0.30527 (13)	0.0392 (5)
C8	1.00699 (17)	0.36037 (18)	0.31824 (15)	0.0537 (7)
C9	1.06821 (19)	0.3851 (2)	0.23237 (18)	0.0710 (9)
C10	1.0223 (2)	0.3654 (3)	0.13446 (19)	0.0809 (10)
C11	0.9166 (2)	0.3196 (3)	0.12143 (17)	0.0825 (12)
C12	0.85329 (18)	0.2966 (2)	0.20620 (15)	0.0620 (8)
C13	0.83557 (14)	0.14007 (14)	0.45185 (14)	0.0386 (5)
C14	0.86489 (19)	0.05919 (17)	0.37959 (17)	0.0576 (7)
C15	0.8752 (2)	-0.05313 (17)	0.40849 (19)	0.0640 (8)
C16	0.85644 (18)	-0.08481 (17)	0.50791 (19)	0.0611 (8)
C17	0.8289 (2)	-0.00608 (19)	0.58004 (18)	0.0663 (8)
C18	0.81820 (19)	0.10688 (17)	0.55274 (16)	0.0552 (7)

C19	0.86206 (15)	0.37784 (14)	0.51789 (13)	0.0404 (5)
C20	0.78895 (19)	0.37952 (16)	0.60511 (16)	0.0501 (7)
H2	0.62426	0.14704	0.35500	0.0647*
H3	0.43471	0.18264	0.31927	0.0849*
H4	0.36530	0.36362	0.33129	0.0908*
H5	0.48444	0.51206	0.37164	0.0809*
H6	0.67549	0.47949	0.40513	0.0636*
H8	1.03834	0.37259	0.38456	0.0644*
H9	1.14070	0.41504	0.24066	0.0853*
H10	1.06347	0.38349	0.07668	0.0970*
H11	0.88740	0.30382	0.05498	0.0990*
H12	0.78055	0.26742	0.19721	0.0744*
H14	0.87764	0.08059	0.31161	0.0690*
H15	0.89494	-0.10715	0.35994	0.0767*
H16	0.86246	-0.16063	0.52674	0.0733*
H17	0.81722	-0.02839	0.64798	0.0795*
H18	0.79937	0.16027	0.60220	0.0662*
H19A	0.86739	0.45405	0.49048	0.0485*
H19B	0.93730	0.35522	0.54233	0.0485*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl1	0.0647 (3)	0.0431 (3)	0.0854 (4)	-0.0019 (2)	-0.0289 (3)	-0.0085 (2)
P1	0.0349 (2)	0.0353 (2)	0.0335 (2)	0.0008 (2)	-0.0029 (2)	-0.0028 (2)
N1	0.0872 (15)	0.0903 (16)	0.0549 (12)	0.0070 (12)	0.0102 (11)	-0.0203 (11)
C1	0.0354 (9)	0.0521 (10)	0.0312 (9)	0.0023 (8)	-0.0011 (7)	-0.0010 (7)
C2	0.0478 (11)	0.0621 (12)	0.0512 (12)	-0.0067 (10)	-0.0057 (9)	-0.0032 (10)
C3	0.0468 (13)	0.0974 (19)	0.0669 (15)	-0.0187 (12)	-0.0128 (11)	0.0019 (13)
C4	0.0408 (12)	0.122 (2)	0.0638 (15)	0.0114 (14)	-0.0064 (10)	0.0088 (15)
C5	0.0521 (13)	0.0833 (16)	0.0662 (14)	0.0248 (12)	-0.0052 (11)	-0.0011 (12)
C6	0.0466 (11)	0.0597 (12)	0.0522 (12)	0.0102 (9)	-0.0052 (9)	-0.0023 (10)
C7	0.0379 (9)	0.0427 (9)	0.0367 (9)	0.0010 (7)	-0.0009 (7)	-0.0030 (7)
C8	0.0439 (11)	0.0720 (14)	0.0451 (11)	-0.0083 (10)	0.0007 (9)	-0.0115 (10)
C9	0.0478 (12)	0.103 (2)	0.0630 (15)	-0.0196 (12)	0.0138 (11)	-0.0119 (13)
C10	0.0630 (16)	0.129 (2)	0.0521 (14)	-0.0080 (15)	0.0210 (12)	0.0016 (14)
C11	0.0623 (16)	0.148 (3)	0.0372 (12)	-0.0081 (16)	0.0014 (10)	-0.0048 (14)
C12	0.0448 (11)	0.1007 (18)	0.0402 (11)	-0.0116 (11)	-0.0036 (9)	-0.0066 (11)
C13	0.0354 (9)	0.0345 (8)	0.0458 (10)	0.0013 (7)	-0.0001 (7)	-0.0017 (7)
C14	0.0790 (15)	0.0447 (11)	0.0492 (12)	0.0012 (10)	0.0054 (10)	-0.0090 (9)
C15	0.0814 (16)	0.0399 (10)	0.0704 (15)	0.0028 (10)	0.0010 (12)	-0.0138 (10)
C16	0.0608 (13)	0.0360 (10)	0.0862 (17)	0.0014 (9)	-0.0024 (12)	0.0041 (10)
C17	0.0838 (16)	0.0533 (12)	0.0625 (14)	0.0078 (12)	0.0131 (12)	0.0171 (11)
C18	0.0703 (14)	0.0457 (10)	0.0505 (12)	0.0111 (10)	0.0143 (10)	0.0026 (9)
C19	0.0452 (10)	0.0372 (9)	0.0382 (9)	0.0018 (7)	-0.0078 (8)	-0.0047 (7)
C20	0.0595 (12)	0.0485 (11)	0.0418 (11)	0.0057 (9)	-0.0060 (10)	-0.0116 (8)

Geometric parameters ( $\text{\AA}$ ,  $\text{^{\circ}}$ )

P1—C1	1.7923 (18)	C16—C17	1.364 (3)
P1—C7	1.7845 (18)	C17—C18	1.385 (3)
P1—C13	1.7851 (17)	C19—C20	1.445 (3)
P1—C19	1.8046 (17)	C2—H2	0.9300
N1—C20	1.133 (3)	C3—H3	0.9300
C1—C2	1.385 (3)	C4—H4	0.9300
C1—C6	1.394 (3)	C5—H5	0.9300
C2—C3	1.378 (3)	C6—H6	0.9300
C3—C4	1.374 (5)	C8—H8	0.9300
C4—C5	1.372 (4)	C9—H9	0.9300
C5—C6	1.382 (3)	C10—H10	0.9300
C7—C8	1.382 (3)	C11—H11	0.9300
C7—C12	1.388 (3)	C12—H12	0.9300
C8—C9	1.377 (3)	C14—H14	0.9300
C9—C10	1.375 (3)	C15—H15	0.9300
C10—C11	1.367 (4)	C16—H16	0.9300
C11—C12	1.376 (3)	C17—H17	0.9300
C13—C14	1.388 (3)	C18—H18	0.9300
C13—C18	1.381 (3)	C19—H19A	0.9700
C14—C15	1.382 (3)	C19—H19B	0.9700
C15—C16	1.362 (3)		
C11…C19 <sup>i</sup>	3.3076 (17)	C14…H2	3.0300
C11…C16 <sup>ii</sup>	3.556 (2)	C15…H12 <sup>i</sup>	3.0900
C11…C19 <sup>iii</sup>	3.3830 (19)	C18…H10 <sup>viii</sup>	3.0400
C11…C12	3.479 (2)	C19…H18	2.9000
C11…H16 <sup>ii</sup>	2.8500	C19…H8	2.7500
C11…H19A <sup>i</sup>	2.3400	C19…H6	2.8600
C11…H8 <sup>iii</sup>	2.8400	C20…H6	3.0900
C11…H19B <sup>iii</sup>	2.4600	C20…H18	2.5900
C11…H12	2.6600	H2…C14	3.0300
C11…H6 <sup>i</sup>	2.8300	H2…C13	2.7500
N1…H18	2.9100	H5…N1 <sup>v</sup>	2.8900
N1…H17 <sup>iv</sup>	2.6100	H6…C11 <sup>ii</sup>	2.8300
N1…H5 <sup>v</sup>	2.8900	H6…H19A	2.5000
C6…C20	3.353 (3)	H6…C19	2.8600
C12…C14	3.586 (3)	H6…C20	3.0900
C12…C11	3.479 (2)	H8…H19B	2.4100
C12…C15 <sup>ii</sup>	3.512 (3)	H8…C19	2.7500
C14…C12	3.586 (3)	H8…C11 <sup>vii</sup>	2.8400
C14…C16 <sup>vi</sup>	3.562 (3)	H10…C18 <sup>ix</sup>	3.0400
C15…C17 <sup>vi</sup>	3.566 (3)	H12…C11	2.6600
C15…C12 <sup>i</sup>	3.512 (3)	H12…C15 <sup>ii</sup>	3.0900
C16…C11 <sup>i</sup>	3.556 (2)	H12…C1	2.8500
C16…C14 <sup>vi</sup>	3.562 (3)	H14…C12	2.9000
C17…C15 <sup>vi</sup>	3.566 (3)	H14…C7	2.8100

C18···C20	3.312 (3)	H16···Cl1 <sup>i</sup>	2.8500
C19···Cl1 <sup>vii</sup>	3.3830 (19)	H17···N1 <sup>x</sup>	2.6100
C19···Cl1 <sup>ii</sup>	3.3076 (17)	H18···N1	2.9100
C20···C6	3.353 (3)	H18···C20	2.5900
C20···C18	3.312 (3)	H18···C19	2.9000
C1···H12	2.8500	H19A···Cl1 <sup>ii</sup>	2.3400
C7···H14	2.8100	H19A···H6	2.5000
C8···H19A	3.0300	H19A···C8	3.0300
C8···H19B	3.0400	H19B···C8	3.0400
C12···H14	2.9000	H19B···H8	2.4100
C13···H2	2.7500	H19B···Cl1 <sup>vii</sup>	2.4600
C1—P1—C7	111.03 (8)	C2—C3—H3	120.00
C1—P1—C13	109.26 (8)	C4—C3—H3	120.00
C1—P1—C19	108.56 (8)	C3—C4—H4	120.00
C7—P1—C13	110.71 (8)	C5—C4—H4	120.00
C7—P1—C19	106.81 (8)	C4—C5—H5	120.00
C13—P1—C19	110.43 (8)	C6—C5—H5	120.00
P1—C1—C2	120.70 (14)	C1—C6—H6	120.00
P1—C1—C6	119.17 (14)	C5—C6—H6	120.00
C2—C1—C6	120.13 (17)	C7—C8—H8	120.00
C1—C2—C3	119.6 (2)	C9—C8—H8	120.00
C2—C3—C4	120.2 (3)	C8—C9—H9	120.00
C3—C4—C5	120.7 (2)	C10—C9—H9	120.00
C4—C5—C6	120.0 (2)	C9—C10—H10	120.00
C1—C6—C5	119.36 (19)	C11—C10—H10	120.00
P1—C7—C8	121.33 (14)	C10—C11—H11	120.00
P1—C7—C12	118.64 (14)	C12—C11—H11	120.00
C8—C7—C12	120.02 (17)	C7—C12—H12	120.00
C7—C8—C9	119.62 (18)	C11—C12—H12	120.00
C8—C9—C10	120.1 (2)	C13—C14—H14	120.00
C9—C10—C11	120.5 (2)	C15—C14—H14	120.00
C10—C11—C12	120.3 (2)	C14—C15—H15	120.00
C7—C12—C11	119.5 (2)	C16—C15—H15	120.00
P1—C13—C14	120.47 (14)	C15—C16—H16	120.00
P1—C13—C18	120.37 (14)	C17—C16—H16	120.00
C14—C13—C18	119.13 (17)	C16—C17—H17	120.00
C13—C14—C15	120.1 (2)	C18—C17—H17	120.00
C14—C15—C16	120.1 (2)	C13—C18—H18	120.00
C15—C16—C17	120.4 (2)	C17—C18—H18	120.00
C16—C17—C18	120.4 (2)	P1—C19—H19A	109.00
C13—C18—C17	119.83 (19)	P1—C19—H19B	109.00
P1—C19—C20	114.38 (13)	C20—C19—H19A	109.00
N1—C20—C19	178.6 (2)	C20—C19—H19B	109.00
C1—C2—H2	120.00	H19A—C19—H19B	108.00
C3—C2—H2	120.00		
C7—P1—C1—C2	100.11 (16)	P1—C1—C6—C5	-178.74 (15)

C7—P1—C1—C6	−79.73 (16)	C2—C1—C6—C5	1.4 (3)
C13—P1—C1—C2	−22.28 (17)	C1—C2—C3—C4	−0.6 (3)
C13—P1—C1—C6	157.89 (14)	C2—C3—C4—C5	1.5 (4)
C19—P1—C1—C2	−142.77 (15)	C3—C4—C5—C6	−0.9 (4)
C19—P1—C1—C6	37.40 (17)	C4—C5—C6—C1	−0.5 (3)
C1—P1—C7—C8	146.77 (15)	P1—C7—C8—C9	−179.84 (17)
C1—P1—C7—C12	−34.67 (18)	C12—C7—C8—C9	1.6 (3)
C13—P1—C7—C8	−91.68 (17)	P1—C7—C12—C11	−178.8 (2)
C13—P1—C7—C12	86.88 (17)	C8—C7—C12—C11	−0.3 (3)
C19—P1—C7—C8	28.58 (18)	C7—C8—C9—C10	−1.0 (4)
C19—P1—C7—C12	−152.86 (16)	C8—C9—C10—C11	−1.0 (4)
C1—P1—C13—C14	95.07 (17)	C9—C10—C11—C12	2.4 (5)
C1—P1—C13—C18	−82.80 (17)	C10—C11—C12—C7	−1.8 (4)
C7—P1—C13—C14	−27.51 (18)	P1—C13—C14—C15	−177.19 (17)
C7—P1—C13—C18	154.62 (15)	C18—C13—C14—C15	0.7 (3)
C19—P1—C13—C14	−145.59 (16)	P1—C13—C18—C17	177.16 (17)
C19—P1—C13—C18	36.54 (18)	C14—C13—C18—C17	−0.7 (3)
C1—P1—C19—C20	48.05 (15)	C13—C14—C15—C16	0.1 (3)
C7—P1—C19—C20	167.85 (13)	C14—C15—C16—C17	−0.9 (3)
C13—P1—C19—C20	−71.71 (15)	C15—C16—C17—C18	0.9 (3)
P1—C1—C2—C3	179.31 (16)	C16—C17—C18—C13	0.0 (3)
C6—C1—C2—C3	−0.9 (3)		

Symmetry codes: (i)  $-x+3/2, y-1/2, -z+1/2$ ; (ii)  $-x+3/2, y+1/2, -z+1/2$ ; (iii)  $x-1/2, -y+1/2, z-1/2$ ; (iv)  $-x+3/2, y+1/2, -z+3/2$ ; (v)  $-x+1, -y+1, -z+1$ ; (vi)  $-x+2, -y, -z+1$ ; (vii)  $x+1/2, -y+1/2, z+1/2$ ; (viii)  $x-1/2, -y+1/2, z+1/2$ ; (ix)  $x+1/2, -y+1/2, z-1/2$ ; (x)  $-x+3/2, y-1/2, -z+3/2$ .

#### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C12—H12 $\cdots$ Cl1	0.93	2.66	3.479 (2)	147
C17—H17 $\cdots$ N1 <sup>x</sup>	0.93	2.61	3.530 (3)	171
C19—H19A $\cdots$ Cl1 <sup>ii</sup>	0.97	2.34	3.3076 (17)	173
C19—H19B $\cdots$ Cl1 <sup>vii</sup>	0.97	2.46	3.3830 (19)	160
C15—H15 $\cdots$ Cg1 <sup>i</sup>	0.93	3.06	3.890 (3)	150

Symmetry codes: (i)  $-x+3/2, y-1/2, -z+1/2$ ; (ii)  $-x+3/2, y+1/2, -z+1/2$ ; (vii)  $x+1/2, -y+1/2, z+1/2$ ; (x)  $-x+3/2, y-1/2, -z+3/2$ .