

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bis{4-chloro-6-formyl-2-[(*E*)-2-(1*H*-imidazol-4-yl- κ N³)ethyliminomethyl- κ N]}phenolato- κ O¹}nickel(II)

 Jia-Wei Mao,^a Hong Zhou,^a Zhi-Quan Pan^{a*} and Xiang-Gao Meng^b
^aKey Laboratory for Green Chemical Process of Ministry of Education, Wuhan Institute of Technology, Wuhan 430073, People's Republic of China, and

^bDepartment of Chemistry, Central China Normal University, Wuhan 430079, People's Republic of China

Correspondence e-mail: zhiqpan@163.com


Received 6 September 2008; accepted 30 September 2008

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.008$ Å; R factor = 0.045; wR factor = 0.081; data-to-parameter ratio = 13.0.

In the title compound, $[\text{Ni}(\text{C}_{13}\text{H}_{11}\text{ClN}_3\text{O}_2)_2]$, the Ni^{II} atom is located on a twofold rotation axis and is six-coordinated by four N atoms and two phenolate O atoms from the two equal Schiff base ligands in a distorted octahedral coordination geometry. The complex molecules are connected by C—H \cdots Cl, C—H \cdots O and N—H \cdots O hydrogen bonds.

Related literature

For related literature on transition metal–Schiff base complexes, see: Casella & Gullotti (1986); Hodnett & Dunn (1970); Kim *et al.* (2005); May *et al.* (2004). For literature related to the synthesis, see: Taniguchi (1984).



Experimental

Crystal data

$[\text{Ni}(\text{C}_{13}\text{H}_{11}\text{ClN}_3\text{O}_2)_2]$
 $M_r = 612.11$
 Tetragonal, $P4_32_12$
 $a = 13.5883$ (16) Å
 $c = 14.0392$ (16) Å
 $V = 2592.2$ (5) Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.00$ mm⁻¹
 $T = 293$ (2) K
 $0.10 \times 0.04 \times 0.02$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.901$, $T_{\max} = 0.978$

21136 measured reflections
 2294 independent reflections
 1253 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.154$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.081$
 $S = 0.82$
 2294 reflections
 177 parameters
 H-atom parameters constrained

$\Delta\rho_{\max} = 0.35$ e Å⁻³
 $\Delta\rho_{\min} = -0.24$ e Å⁻³
 Absolute structure: Flack (1983),
 920 Friedel pairs
 Flack parameter: 0.02 (3)

Table 1

Selected geometric parameters (Å, °).

Ni1—O1	2.054 (3)	Ni1—N1	2.102 (4)
Ni1—N2	2.068 (4)		
O1—Ni1—O1 ⁱ	87.37 (17)	N2—Ni1—N1	90.27 (15)
O1—Ni1—N2 ⁱ	91.40 (14)	O1—Ni1—N1 ⁱ	89.72 (13)
O1—Ni1—N2	178.49 (14)	N2—Ni1—N1 ⁱ	91.14 (15)
N2 ⁱ —Ni1—N2	89.8 (2)	N1—Ni1—N1 ⁱ	178.0 (2)
O1—Ni1—N1	88.84 (14)		

 Symmetry code: (i) $y, x, -z$.

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C9—H9B \cdots Cl1 ⁱⁱ	0.97	2.82	3.475 (5)	125
C12—H12 \cdots O2 ⁱⁱⁱ	0.93	2.36	3.287 (7)	174
N3—H3A \cdots O1 ^{iv}	0.86	2.06	2.899 (5)	166

Symmetry codes: (ii) $-x + 1, -y, z - \frac{1}{2}$; (iii) $y + \frac{1}{2}, -x + \frac{1}{2}, z - \frac{3}{4}$; (iv) $-y + \frac{1}{2}, x + \frac{1}{2}, z - \frac{1}{4}$.

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors gratefully acknowledge financial support from the Midlife and Youth Excellent Innovation Group of Hubei Province, China (grant No. T200802), the Key Foundation of the Education Department of Hubei Province, China (grant No. D20081503), and the Graduate Innovation Foundation of Wuhan Institute of Technology (RGCT200804).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2154).

References

- Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Casella, L. & Gullotti, M. (1986). *Inorg. Chem.* **25**, 1293–1303.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Hodnett, E. M. & Dunn, W. J. (1970). *J. Med. Chem.* **13**, 768–770.
- Kim, H.-J., Kim, W., Lough, A. J., Kim, B. M. & Chin, J. (2005). *J. Am. Chem. Soc.* **127**, 16776–16777.
- May, J. P., Ting, R., Lermer, L., Thomas, J. M., Roupioz, Y. & Perrin, D. M. (2004). *J. Am. Chem. Soc.* **126**, 4145–4156.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Taniguchi, S. (1984). *Bull. Chem. Soc. Jpn.*, **57**, 2683–2689.

supplementary materials

Acta Cryst. (2008). E64, m1367-m1368 [doi:10.1107/S1600536808031577]

Bis{4-chloro-6-formyl-2-[(*E*)-2-(1*H*-imidazol-4-yl- κ N³)ethyliminomethyl- κ N]}phenolato- κ O¹}nickel(II)

J.-W. Mao, H. Zhou, Z.-Q. Pan and X.-G. Meng

Comment

Transition metal–Schiff base complexes have been an interesting field for a long time due to their striking biological activities (Casella & Gullotti, 1986; Hodnett & Dunn, 1970; Kim *et al.*, 2005; May *et al.*, 2004). In this paper, we report the crystal structure of a new nickel(II) complex with a Schiff base ligand, 2-[(*E*)-2-(1*H*-imidazol-4-yl)ethylimino)methyl]-4-chloro-6-formylphenolate.

In the title compound, the Ni^{II} atom is located on a twofold rotation axis and six-coordinated by four N atoms and two phenolate O atoms from two Schiff base ligands (Fig. 1). The coordination geometry of the Ni atom can be described as distorted octahedral. The two phenolate O atoms and the two imidazole N atoms are located in the equatorial plane, with Ni—O distance of 2.054 (3) Å and Ni—N distance of 2.068 (4) Å (Table 1), and with the mean plane deviation of 0.0147 (2) Å. The other two N atoms from the imino groups of the Schiff base ligands occupy the axial positions, with somewhat long Ni—N distance of 2.102 (4) Å. The complex molecules are connected by C—H···Cl, C—H···O and N—H···O hydrogen bonds (Table 2).

Experimental

2,6-Diformyl-4-chlorophenol was prepared using the method of Taniguchi (1984). The title compound was synthesized by the following procedure: To an acetonitrile solution (10 ml) of 2,6-diformyl-4-chlorophenol (0.092 g, 0.5 mmol) and Ni(ClO₄)₂·6H₂O (0.018 g, 0.25 mmol), a solution of NaOH (0.041 g, 1 mmol) and histamine dihydrochloride (0.092 g, 0.5 mmol) in 15 ml of absolute methanol was added dropwise. After the mixture was stirred at ambient temperature for about 1 h, a red solution appeared and then the stirring was continued for 3 h. Red needle crystals of the title compound suitable for X-ray diffraction were obtained in about a month.

Refinement

H atoms were positioned geometrically and refined as riding atoms, with C—H = 0.93(CH), 0.97(CH₂) Å and N—H = 0.86 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$.

Figures




Fig. 1. Molecular structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level. [Symmetry code: (i) $x, y, -z$.]

Bis{4-chloro-6-formyl-2-[(E)-2-(1H-imidazol-4-yl-κN³)ethyliminomethyl-κN]phenolato-κO¹}nickel(II)

Crystal data

[Ni(C ₁₃ H ₁₁ ClN ₃ O ₂) ₂]	Z = 4
<i>M_r</i> = 612.11	<i>F</i> ₀₀₀ = 1256
Tetragonal, <i>P</i> 4 ₃ 2 ₁ 2	<i>D_x</i> = 1.568 Mg m ⁻³
Hall symbol: P 4nw 2abw	Mo <i>K</i> α radiation
<i>a</i> = 13.5883 (16) Å	λ = 0.71073 Å
<i>b</i> = 13.5883 (16) Å	Cell parameters from 1360 reflections
<i>c</i> = 14.0392 (16) Å	θ = 2.6–15.1°
α = 90°	μ = 1.00 mm ⁻¹
β = 90°	<i>T</i> = 293 (2) K
γ = 90°	Needle, red
<i>V</i> = 2592.2 (5) Å ³	0.10 × 0.04 × 0.02 mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer	2294 independent reflections
Radiation source: fine-focus sealed tube	1253 reflections with <i>I</i> > 2σ(<i>I</i>)
Monochromator: graphite	<i>R</i> _{int} = 0.154
<i>T</i> = 293(2) K	θ _{max} = 25.0°
φ and ω scans	θ _{min} = 2.1°
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	<i>h</i> = -16→16
<i>T</i> _{min} = 0.901, <i>T</i> _{max} = 0.978	<i>k</i> = -16→16
21136 measured reflections	<i>l</i> = -14→16

Refinement

Refinement on <i>F</i> ²	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.045$	$w = 1/[\sigma^2(F_o^2) + (0.0298P)^2]$
$wR(F^2) = 0.081$	where $P = (F_o^2 + 2F_c^2)/3$
<i>S</i> = 0.82	(Δ/σ) _{max} = 0.003
2294 reflections	Δρ _{max} = 0.35 e Å ⁻³
177 parameters	Δρ _{min} = -0.24 e Å ⁻³
Primary atom site location: structure-invariant direct methods	Extinction correction: none
Secondary atom site location: difference Fourier map	Absolute structure: Flack (1983), 920 Friedel pairs
	Flack parameter: 0.02 (3)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. The reason of the large R_{int} value is the poor quality and small size of the crystal sample. Although many efforts were made to select better crystal for experiment, each time we failed.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.23545 (4)	0.23545 (4)	0.0000	0.0433 (3)
C1	0.2375 (4)	0.0575 (3)	0.1277 (3)	0.0428 (12)
C2	0.2104 (4)	0.0113 (4)	0.2139 (4)	0.0564 (15)
C3	0.2586 (4)	-0.0728 (4)	0.2478 (4)	0.0656 (14)
H3	0.2381	-0.1026	0.3041	0.079*
C4	0.3356 (4)	-0.1106 (4)	0.1977 (5)	0.0642 (17)
C5	0.3666 (4)	-0.0659 (4)	0.1150 (4)	0.0593 (15)
H5	0.4202	-0.0920	0.0825	0.071*
C6	0.3205 (4)	0.0165 (3)	0.0790 (4)	0.0455 (13)
C7	0.1300 (4)	0.0521 (5)	0.2722 (4)	0.0800 (19)
H7	0.1037	0.1125	0.2545	0.096*
C8	0.3581 (3)	0.0522 (4)	-0.0096 (4)	0.0519 (13)
H8	0.4055	0.0129	-0.0384	0.062*
C9	0.3904 (4)	0.1472 (4)	-0.1444 (4)	0.0700 (17)
H9A	0.4022	0.0839	-0.1743	0.084*
H9B	0.4539	0.1760	-0.1294	0.084*
C10	0.3368 (4)	0.2136 (4)	-0.2143 (3)	0.0636 (15)
H10A	0.3701	0.2112	-0.2754	0.076*
H10B	0.2704	0.1892	-0.2233	0.076*
C11	0.3326 (4)	0.3166 (4)	-0.1810 (4)	0.0500 (14)
C12	0.3728 (4)	0.4001 (4)	-0.2169 (4)	0.0620 (16)
H12	0.4097	0.4056	-0.2724	0.074*
C13	0.2960 (3)	0.4340 (4)	-0.0856 (4)	0.0504 (14)
H13	0.2707	0.4695	-0.0345	0.060*
Cl1	0.39581 (12)	-0.21688 (11)	0.23773 (13)	0.1081 (7)
N1	0.3352 (3)	0.1319 (3)	-0.0552 (3)	0.0487 (11)
N2	0.2844 (3)	0.3389 (3)	-0.0971 (3)	0.0480 (11)
N3	0.3486 (3)	0.4739 (3)	-0.1563 (3)	0.0584 (12)
H3A	0.3641	0.5349	-0.1622	0.070*
O1	0.1862 (2)	0.1302 (2)	0.0935 (2)	0.0494 (9)
O2	0.0962 (3)	0.0119 (3)	0.3416 (3)	0.1061 (15)

Atomic displacement parameters (\AA^2)

U^{11} U^{22} U^{33} U^{12} U^{13} U^{23}

supplementary materials

Ni1	0.0453 (3)	0.0453 (3)	0.0393 (5)	0.0010 (4)	0.0040 (3)	-0.0040 (3)
C1	0.038 (3)	0.050 (3)	0.040 (3)	-0.012 (3)	-0.004 (3)	0.003 (3)
C2	0.057 (4)	0.056 (4)	0.056 (4)	-0.014 (3)	-0.008 (3)	0.009 (3)
C3	0.067 (4)	0.064 (4)	0.066 (4)	-0.024 (3)	-0.020 (5)	0.016 (4)
C4	0.059 (4)	0.055 (4)	0.079 (5)	0.001 (3)	-0.026 (4)	0.017 (4)
C5	0.041 (3)	0.060 (4)	0.077 (5)	-0.003 (3)	-0.011 (3)	-0.002 (3)
C6	0.043 (3)	0.043 (3)	0.050 (4)	-0.003 (3)	-0.006 (3)	0.001 (3)
C7	0.080 (5)	0.115 (5)	0.045 (5)	-0.021 (4)	0.001 (4)	0.030 (4)
C8	0.048 (3)	0.044 (3)	0.064 (4)	0.007 (3)	0.004 (3)	-0.013 (3)
C9	0.094 (4)	0.056 (4)	0.060 (4)	0.015 (3)	0.028 (4)	0.004 (3)
C10	0.082 (4)	0.069 (4)	0.040 (4)	0.011 (3)	0.020 (3)	-0.006 (3)
C11	0.060 (4)	0.053 (4)	0.038 (4)	0.003 (3)	0.002 (3)	0.002 (3)
C12	0.070 (4)	0.068 (4)	0.048 (4)	0.015 (3)	0.018 (3)	0.000 (3)
C13	0.052 (4)	0.054 (4)	0.045 (4)	0.004 (3)	0.009 (3)	0.006 (3)
Cl1	0.1068 (12)	0.0770 (11)	0.1405 (16)	0.0095 (10)	-0.0362 (12)	0.0368 (12)
N1	0.052 (3)	0.054 (3)	0.041 (3)	-0.001 (2)	0.009 (2)	-0.003 (2)
N2	0.063 (3)	0.042 (3)	0.039 (3)	-0.003 (2)	0.003 (2)	0.000 (2)
N3	0.063 (3)	0.053 (3)	0.059 (3)	-0.009 (2)	0.006 (3)	0.018 (3)
O1	0.043 (2)	0.059 (2)	0.046 (2)	0.0090 (17)	0.0044 (17)	0.0083 (18)
O2	0.116 (4)	0.135 (4)	0.068 (4)	-0.013 (3)	0.017 (3)	0.026 (3)

Geometric parameters (\AA , $^\circ$)

Ni1—O1	2.054 (3)	C7—H7	0.9300
Ni1—O1 ⁱ	2.054 (3)	C8—N1	1.296 (5)
Ni1—N2 ⁱ	2.068 (4)	C8—H8	0.9300
Ni1—N2	2.068 (4)	C9—N1	1.474 (5)
Ni1—N1	2.102 (4)	C9—C10	1.519 (6)
Ni1—N1 ⁱ	2.102 (4)	C9—H9A	0.9700
C1—O1	1.301 (5)	C9—H9B	0.9700
C1—C2	1.412 (6)	C10—C11	1.477 (6)
C1—C6	1.431 (6)	C10—H10A	0.9700
C2—C3	1.401 (6)	C10—H10B	0.9700
C2—C7	1.474 (7)	C11—C12	1.355 (6)
C3—C4	1.362 (7)	C11—N2	1.382 (5)
C3—H3	0.9300	C12—N3	1.356 (5)
C4—C5	1.376 (7)	C12—H12	0.9300
C4—Cl1	1.752 (5)	C13—N2	1.312 (5)
C5—C6	1.380 (6)	C13—N3	1.337 (5)
C5—H5	0.9300	C13—H13	0.9300
C6—C8	1.429 (6)	N3—H3A	0.8600
C7—O2	1.208 (5)		
O1—Ni1—O1 ⁱ	87.37 (17)	C2—C7—H7	117.9
O1—Ni1—N2 ⁱ	91.40 (14)	N1—C8—C6	128.9 (5)
O1 ⁱ —Ni1—N2 ⁱ	178.49 (14)	N1—C8—H8	115.5
O1—Ni1—N2	178.49 (14)	C6—C8—H8	115.5
O1 ⁱ —Ni1—N2	91.40 (14)	N1—C9—C10	112.8 (4)

N2 ⁱ —Ni1—N2	89.8 (2)	N1—C9—H9A	109.0
O1—Ni1—N1	88.84 (14)	C10—C9—H9A	109.0
O1 ⁱ —Ni1—N1	89.72 (13)	N1—C9—H9B	109.0
N2 ⁱ —Ni1—N1	91.14 (15)	C10—C9—H9B	109.0
N2—Ni1—N1	90.27 (15)	H9A—C9—H9B	107.8
O1—Ni1—N1 ⁱ	89.72 (13)	C11—C10—C9	112.2 (4)
O1 ⁱ —Ni1—N1 ⁱ	88.84 (14)	C11—C10—H10A	109.2
N2 ⁱ —Ni1—N1 ⁱ	90.27 (15)	C9—C10—H10A	109.2
N2—Ni1—N1 ⁱ	91.14 (15)	C11—C10—H10B	109.2
N1—Ni1—N1 ⁱ	178.0 (2)	C9—C10—H10B	109.2
O1—C1—C2	120.9 (5)	H10A—C10—H10B	107.9
O1—C1—C6	122.8 (4)	C12—C11—N2	108.9 (5)
C2—C1—C6	116.2 (5)	C12—C11—C10	131.3 (5)
C3—C2—C1	122.2 (5)	N2—C11—C10	119.7 (5)
C3—C2—C7	117.6 (5)	C11—C12—N3	106.8 (5)
C1—C2—C7	120.2 (5)	C11—C12—H12	126.6
C4—C3—C2	119.4 (5)	N3—C12—H12	126.6
C4—C3—H3	120.3	N2—C13—N3	111.9 (5)
C2—C3—H3	120.3	N2—C13—H13	124.1
C3—C4—C5	120.3 (5)	N3—C13—H13	124.1
C3—C4—C11	120.3 (5)	C8—N1—C9	114.6 (4)
C5—C4—C11	119.4 (5)	C8—N1—Ni1	122.1 (3)
C4—C5—C6	121.9 (5)	C9—N1—Ni1	123.2 (3)
C4—C5—H5	119.0	C13—N2—C11	105.3 (4)
C6—C5—H5	119.0	C13—N2—Ni1	128.9 (4)
C5—C6—C8	115.6 (5)	C11—N2—Ni1	124.4 (3)
C5—C6—C1	119.9 (5)	C13—N3—C12	107.2 (4)
C8—C6—C1	124.4 (4)	C13—N3—H3A	126.4
O2—C7—C2	124.1 (6)	C12—N3—H3A	126.4
O2—C7—H7	117.9	C1—O1—Ni1	126.1 (3)
O1—C1—C2—C3	173.9 (4)	C10—C9—N1—Ni1	-28.0 (6)
C6—C1—C2—C3	-3.1 (6)	O1—Ni1—N1—C8	-17.2 (4)
O1—C1—C2—C7	-7.0 (7)	O1 ⁱ —Ni1—N1—C8	-104.6 (4)
C6—C1—C2—C7	176.0 (4)	N2 ⁱ —Ni1—N1—C8	74.1 (4)
C1—C2—C3—C4	1.5 (7)	N2—Ni1—N1—C8	164.0 (4)
C7—C2—C3—C4	-177.6 (5)	O1—Ni1—N1—C9	167.8 (4)
C2—C3—C4—C5	0.8 (8)	O1 ⁱ —Ni1—N1—C9	80.4 (4)
C2—C3—C4—C11	-179.0 (3)	N2 ⁱ —Ni1—N1—C9	-100.8 (4)
C3—C4—C5—C6	-1.3 (8)	N2—Ni1—N1—C9	-11.0 (4)
C11—C4—C5—C6	178.5 (4)	N3—C13—N2—C11	-0.2 (5)
C4—C5—C6—C8	-177.3 (5)	N3—C13—N2—Ni1	166.5 (3)
C4—C5—C6—C1	-0.4 (7)	C12—C11—N2—C13	0.6 (6)
O1—C1—C6—C5	-174.4 (4)	C10—C11—N2—C13	178.1 (5)
C2—C1—C6—C5	2.5 (6)	C12—C11—N2—Ni1	-166.8 (3)
O1—C1—C6—C8	2.2 (7)	C10—C11—N2—Ni1	10.7 (6)
C2—C1—C6—C8	179.2 (4)	N2 ⁱ —Ni1—N2—C13	-52.1 (4)

supplementary materials

C3—C2—C7—O2	-8.3 (8)	N1—Ni1—N2—C13	-143.3 (4)
C1—C2—C7—O2	172.5 (5)	N2 ⁱ —Ni1—N2—C11	112.2 (4)
C5—C6—C8—N1	-173.5 (5)	N1—Ni1—N2—C11	21.0 (4)
C1—C6—C8—N1	9.8 (8)	N1 ⁱ —Ni1—N2—C11	-157.6 (4)
N1—C9—C10—C11	69.3 (6)	N2—C13—N3—C12	-0.3 (6)
C9—C10—C11—C12	115.1 (6)	C11—C12—N3—C13	0.6 (6)
C9—C10—C11—N2	-61.8 (6)	C2—C1—O1—Ni1	157.3 (3)
N2—C11—C12—N3	-0.7 (6)	C6—C1—O1—Ni1	-25.9 (6)
C10—C11—C12—N3	-177.9 (5)	O1 ⁱ —Ni1—O1—C1	118.8 (4)
C6—C8—N1—C9	178.7 (5)	N1—Ni1—O1—C1	29.0 (4)
C6—C8—N1—Ni1	3.3 (7)	N1 ⁱ —Ni1—O1—C1	-152.4 (3)
C10—C9—N1—C8	156.6 (4)		

Symmetry codes: (i) $y, x, -z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C9—H9B \cdots C11 ⁱⁱ	0.97	2.82	3.475 (5)	125
C12—H12 \cdots O2 ⁱⁱⁱ	0.93	2.36	3.287 (7)	174
N3—H3A \cdots O1 ^{iv}	0.86	2.06	2.899 (5)	166

Symmetry codes: (ii) $-x+1, -y, z-1/2$; (iii) $y+1/2, -x+1/2, z-3/4$; (iv) $-y+1/2, x+1/2, z-1/4$.

