

catena-Poly[[[aqua(pyridine-2,6-dicarboxylato *N*-oxide- κ^2O^1,O^2)cobalt(II)]- μ -1,3-di-4-pyridylpropane- $\kappa^2N:N'$] dihydrate]

Li-Jin Wang

Department of Chemistry, Lishui University, Lishui 323000, Zhejiang, People's Republic of China

Correspondence e-mail: lswlj2008@yahoo.cn


Received 4 October 2008; accepted 10 October 2008

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.008$ Å; R factor = 0.059; wR factor = 0.175; data-to-parameter ratio = 13.4.

In the title compound, $\{[Co(C_7H_3NO_5)(C_{13}H_{14}N_2)(H_2O)] \cdot 2H_2O\}_n$, the Co^{II} atom is coordinated by two O atoms from a pyridine-2,6-dicarboxylate *N*-oxide ligand, two N atoms from two 1,3-di-4-pyridylpropane ligands and one water molecule, and displays a distorted square-pyramidal coordination geometry. The 1,3-di-4-pyridylpropane ligands link the Co^{II} atoms into an infinite zigzag chain parallel to [010]. The chains are further linked through $O-H \cdots O$ and $C-H \cdots O$ hydrogen bonds, forming a three-dimensional supramolecular network.

Related literature

For related literature on metal complexes with pyridine-2,6-dicarboxylate *N*-oxide, see: Nathan *et al.* (1985); Wen *et al.* (2005, 2006); Wu *et al.* (2007). For related literature on metal complexes with 1,3-di-4-pyridylpropane, see: Konar *et al.* (2003); Lai & Tiekink (2004); Li *et al.* (2004).



Experimental

Crystal data

$[Co(C_7H_3NO_5)(C_{13}H_{14}N_2)(H_2O)] \cdot 2H_2O$
 $M_r = 492.34$
 Monoclinic, $P2_1/c$
 $a = 10.2712$ (12) Å
 $b = 11.5251$ (13) Å
 $c = 18.309$ (2) Å

$\beta = 90.521$ (2)°
 $V = 2167.3$ (4) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.84$ mm⁻¹
 $T = 296$ (2) K
 $0.35 \times 0.29 \times 0.25$ mm

Data collection

Bruker SMART APEXII CCD
 area-detector diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{min} = 0.751$, $T_{max} = 0.817$

10818 measured reflections
 3897 independent reflections
 2170 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.064$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.059$
 $wR(F^2) = 0.175$
 $S = 1.00$
 3897 reflections
 290 parameters

9 restraints
 H-atom parameters constrained
 $\Delta\rho_{max} = 0.66$ e Å⁻³
 $\Delta\rho_{min} = -0.40$ e Å⁻³

Table 1

Selected bond lengths (Å).

Co1—O1	1.927 (4)	Co1—N3 ⁱ	2.000 (4)
Co1—O5	1.932 (3)	Co1—O1W	2.184 (4)
Co1—N2	1.994 (4)		

 Symmetry code: (i) $x, y - 1, z$.

Table 2

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O1W—H1W ⁱⁱ ···O3 ⁱⁱ	0.82	1.83	2.635 (5)	165
O1W—H2W ⁱⁱ ···O2W	0.82	1.99	2.732 (6)	150
O2W—H3W ⁱⁱ ···O3W ⁱⁱⁱ	0.84	2.55	3.018 (9)	117
O2W—H4W ⁱⁱ ···O3	0.84	1.98	2.778 (6)	157
O3W—H5W ⁱⁱ ···O2	0.84	2.01	2.828 (7)	166
O3W—H6W ⁱⁱ ···O4 ^{iv}	0.82	2.19	2.935 (7)	150
C3—H3 ⁱⁱ ···O4 ^v	0.93	2.46	3.364 (8)	165
C9—H9 ⁱⁱ ···O5 ^{vi}	0.93	2.46	3.366 (7)	164
C15—H15A ⁱⁱ ···O2 ^{vii}	0.97	2.57	3.492 (7)	159
C18—H18 ⁱⁱ ···O3W ^{vi}	0.93	2.55	3.369 (7)	149

Symmetry codes: (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (iv) $x, y + 1, z$; (v) $-x + 1, -y, -z + 1$; (vi) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$; (vii) $x, -y + \frac{3}{2}, z - \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The author is grateful to Lishui University for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2158).

References

- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Konar, S., Zangrando, E., Drew, M. G. B., Mallah, T., Ribas, J. & Chaudhuri, N. R. (2003). *Inorg. Chem.* **42**, 5966–5973.
- Lai, C. S. & Tiekink, E. R. T. (2004). *CrystEngComm*, **6**, 593–605.
- Li, X., Cao, R., Sun, D., Bi, W., Wang, Y., Li, X. & Hong, M. (2004). *Cryst. Growth Des.* **4**, 775–780.
- Nathan, L. C., Doyle, C. A., Mooring, A. M., Zapfen, D. C., Larsen, S. K. & Pierpont, C. G. (1985). *Inorg. Chem.* **24**, 2763–2766.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wen, L.-L., Dang, D.-B., Duan, C.-Y., Li, Y.-Z., Tian, Z.-F. & Meng, Q.-J. (2005). *Inorg. Chem.* **44**, 7161–7170.
- Wen, L.-L., Tian, Z.-F., Lin, J.-G., Zhu, H.-Z. & Meng, Q.-J. (2006). *Z. Anorg. Allg. Chem.* **632**, 689–694.
- Wu, W.-P., Wang, Y.-Y., Wu, Y.-P., Liu, J.-Q., Zeng, X.-R., Shi, Q.-Z. & Peng, S.-M. (2007). *CrystEngComm*, **9**, 753–757.

supplementary materials

Acta Cryst. (2008). E64, m1415-m1416 [doi:10.1107/S1600536808032741]

catena-Poly[[[aqua(pyridine-2,6-dicarboxylato *N*-oxide- $\kappa^2 O^1, O^2$)cobalt(II)]- μ -1,3-di-4-pyridylpropane- $\kappa^2 N:N'$] dihydrate]

L.-J. Wang

Comment

In the structural investigation of metal complexes with pyridine-2,6-dicarboxylate-*N*-oxide (pdco), it has been found that pdco functions as a multidentate ligand with versatile coordination modes (Nathan *et al.*, 1985; Wen *et al.*, 2005, 2006; Wu *et al.*, 2007). As is well known, 1,3-di-4-pyridylpropane may act in bidentate bridging or monodentate terminal modes, leading to the formation of one-, two- or three-dimensional network (Konar *et al.*, 2003; Lai & Tiekink, 2004; Li *et al.*, 2004). On the basis of these observations, we utilize pdco, 1,3-di-4-pyridylpropane and Co^{II} ion as building blocks. A new one-dimensional coordination framework has been obtained from the hydrothermal treatment in an alkaline aqueous solution.

As illustrated in Fig. 1, the Co^{II} atom exists in a distorted square-pyramidal environment, defined by two O atoms from one pdco ligand, two N atoms from two 1,3-di-4-pyridylpropane ligands and one water molecule (Table 1). The O1, O5, N2, N3^{*i*} atoms (*i* = *x*, -1 + *y*, *z*) in the basal plane are almost coplanar, and a water molecule lies at the apical position. The 1,3-di-4-pyridylpropane ligand in a bidentate bridging mode links the Co^{II} atoms into an infinite zigzag chain, with the shortest Co...Co separation of 11.525 (3) Å and a Co—C13—Coⁱⁱ angle (*ii* = *x*, 1 + *y*, *z*) of 100.06 (4)°. The chains are further self-assembled into a three-dimensional supramolecular network through O—H...O and C—H...O hydrogen bonds (Table 2; Fig. 2).

Experimental

A mixture of cobalt chloride (0.238 g, 1 mmol), pyridine-2,6-dicarboxylic acid *N*-oxide (0.181 g, 1 mmol), 1,3-di-4-pyridylpropane (0.198 g, 1 mmol), NaOH (0.06 g, 1.5 mmol) and H₂O (12 ml) was placed in a 23 ml Teflon-lined reactor, which was heated to 433 K for 3 d and then cooled to room temperature at a rate of 10 K h⁻¹. The crystals obtained were washed with water and dried in air.

Refinement

H atoms bound to C atoms were positioned geometrically and refined as riding atoms, with C—H = 0.93(CH) and 0.97(CH₂) Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. H atoms of water molecules were located on a difference Fourier map and fixed in the refinements, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. The structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level. H atoms have been omitted for clarity. [Symmetry code: (i) $x, -1 + y, z$.]




Fig. 2. A packing view of the title compound. Hydrogen bonds are shown as dashed lines.

catena-Poly[[[aqua(pyridine-2,6-dicarboxylato N-oxide- κ^2O^1, O^2)cobalt(II)]- μ -1,3-di-4-pyridylpropane- $\kappa^2N:N'$] dihydrate]

Crystal data

$[\text{Co}(\text{C}_7\text{H}_3\text{NO}_5)(\text{C}_{13}\text{H}_{14}\text{N}_2)(\text{H}_2\text{O})] \cdot 2\text{H}_2\text{O}$

$M_r = 492.34$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 10.2712\ (12)\ \text{\AA}$

$b = 11.5251\ (13)\ \text{\AA}$

$c = 18.309\ (2)\ \text{\AA}$

$\beta = 90.521\ (2)^\circ$

$V = 2167.3\ (4)\ \text{\AA}^3$

$Z = 4$

$F_{000} = 1020$

$D_x = 1.509\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 5837 reflections

$\theta = 2.8\text{--}27.9^\circ$

$\mu = 0.84\ \text{mm}^{-1}$

$T = 296\ (2)\ \text{K}$

Block, colorless

$0.35 \times 0.29 \times 0.25\ \text{mm}$

Data collection

Bruker SMART APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 296\ (2)\ \text{K}$

3897 independent reflections

2170 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.064$

$\theta_{\text{max}} = 25.2^\circ$

φ and ω scans $\theta_{\min} = 2.0^\circ$
 Absorption correction: multi-scan $h = -12 \rightarrow 12$
 (SADABS; Sheldrick, 1996) $k = -13 \rightarrow 13$
 $T_{\min} = 0.751$, $T_{\max} = 0.817$ $l = -20 \rightarrow 21$
 10818 measured reflections

Refinement

Refinement on F^2 Secondary atom site location: difference Fourier map
 Least-squares matrix: full Hydrogen site location: inferred from neighbouring sites
 $R[F^2 > 2\sigma(F^2)] = 0.059$ H-atom parameters constrained
 $wR(F^2) = 0.175$ $w = 1/[\sigma^2(F_o^2) + (0.0828P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $S = 1.00$ $(\Delta/\sigma)_{\max} < 0.001$
 3897 reflections $\Delta\rho_{\max} = 0.66 \text{ e } \text{\AA}^{-3}$
 290 parameters $\Delta\rho_{\min} = -0.40 \text{ e } \text{\AA}^{-3}$
 9 restraints Extinction correction: SHELXL97 (Sheldrick, 2008),
 $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
 Primary atom site location: structure-invariant direct methods Extinction coefficient: 0.0030 (8)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O4	0.3287 (4)	-0.0221 (4)	0.4108 (2)	0.0776 (13)
N3	0.2102 (4)	1.1756 (4)	0.1902 (2)	0.0519 (11)
C19	0.2965 (6)	1.0894 (5)	0.1927 (3)	0.0622 (16)
H19	0.3729	1.1009	0.2194	0.075*
C20	0.2794 (6)	0.9841 (5)	0.1583 (3)	0.0609 (15)
H20	0.3432	0.9272	0.1621	0.073*
C18	0.1003 (5)	1.1549 (5)	0.1514 (3)	0.0568 (15)
H18	0.0376	1.2129	0.1486	0.068*
C17	0.0769 (6)	1.0515 (5)	0.1159 (3)	0.0597 (15)
H17	-0.0007	1.0411	0.0902	0.072*
C16	0.1677 (5)	0.9631 (5)	0.1180 (3)	0.0565 (14)
C15	0.1470 (6)	0.8479 (5)	0.0794 (3)	0.0647 (16)
H15A	0.1842	0.8532	0.0310	0.078*
H15B	0.1948	0.7885	0.1059	0.078*
Co1	0.24328 (6)	0.32149 (5)	0.24652 (3)	0.0423 (3)
O5	0.2788 (4)	0.2215 (3)	0.32898 (19)	0.0565 (10)
N2	0.1568 (4)	0.4237 (4)	0.1728 (2)	0.0522 (11)
O2	0.2658 (5)	0.5550 (4)	0.4120 (2)	0.0892 (15)
O1	0.2384 (4)	0.4531 (3)	0.3113 (2)	0.0777 (13)
O3	0.4937 (4)	0.0354 (3)	0.3417 (2)	0.0761 (12)
C6	0.3381 (5)	0.3644 (5)	0.4174 (3)	0.0513 (14)
C12	0.1618 (6)	0.5381 (5)	0.0652 (3)	0.0593 (15)
H12	0.2043	0.5575	0.0223	0.071*

supplementary materials

N1	0.3424 (4)	0.2545 (4)	0.3900 (2)	0.0494 (11)
C3	0.4691 (6)	0.1910 (6)	0.4901 (3)	0.0702 (18)
H3	0.5156	0.1326	0.5137	0.084*
C7	0.2755 (6)	0.4653 (5)	0.3766 (3)	0.0607 (15)
C2	0.4082 (5)	0.1683 (5)	0.4248 (3)	0.0496 (13)
C13	-0.0066 (6)	0.6897 (4)	0.0347 (3)	0.0660 (17)
H13A	-0.0981	0.6750	0.0252	0.079*
H13B	0.0381	0.6903	-0.0117	0.079*
C10	0.0487 (5)	0.5928 (5)	0.0824 (3)	0.0540 (14)
C1	0.4076 (6)	0.0499 (5)	0.3885 (3)	0.0559 (14)
C5	0.3971 (6)	0.3843 (5)	0.4831 (3)	0.0664 (16)
H5	0.3931	0.4585	0.5029	0.080*
C11	0.2145 (5)	0.4541 (5)	0.1107 (3)	0.0547 (14)
H11	0.2918	0.4180	0.0976	0.066*
C8	0.0451 (6)	0.4774 (5)	0.1890 (3)	0.0675 (17)
H8	0.0037	0.4577	0.2322	0.081*
C4	0.4619 (6)	0.2997 (6)	0.5211 (3)	0.0730 (18)
H4	0.4999	0.3150	0.5663	0.088*
C9	-0.0109 (6)	0.5589 (5)	0.1456 (3)	0.0653 (16)
H9	-0.0899	0.5918	0.1587	0.078*
C14	0.0088 (6)	0.8095 (4)	0.0717 (3)	0.0641 (17)
H14A	-0.0387	0.8667	0.0432	0.077*
H14B	-0.0299	0.8064	0.1198	0.077*
O1W	0.4465 (4)	0.3311 (4)	0.2130 (3)	0.0969 (16)
H1W	0.4767	0.3944	0.2011	0.145*
H2W	0.4891	0.3034	0.2471	0.145*
O2W	0.6417 (5)	0.2145 (5)	0.2840 (3)	0.1170 (19)
H3W	0.6669	0.1724	0.2494	0.175*
H4W	0.5887	0.1753	0.3084	0.175*
O3W	0.2003 (5)	0.7719 (5)	0.3494 (4)	0.141 (2)
H5W	0.2281	0.7061	0.3618	0.212*
H6W	0.2190	0.8220	0.3795	0.212*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O4	0.089 (3)	0.053 (3)	0.091 (3)	-0.012 (2)	0.005 (2)	0.013 (2)
N3	0.058 (3)	0.044 (3)	0.053 (3)	0.005 (2)	-0.012 (2)	0.000 (2)
C19	0.074 (4)	0.043 (4)	0.069 (4)	0.008 (3)	-0.015 (3)	-0.002 (3)
C20	0.066 (4)	0.045 (4)	0.071 (4)	0.008 (3)	-0.010 (3)	-0.005 (3)
C18	0.061 (4)	0.038 (3)	0.071 (4)	0.006 (3)	-0.012 (3)	0.002 (3)
C17	0.070 (4)	0.038 (3)	0.071 (4)	-0.008 (3)	-0.014 (3)	-0.004 (3)
C16	0.067 (4)	0.043 (3)	0.059 (3)	0.003 (3)	-0.002 (3)	-0.003 (3)
C15	0.080 (4)	0.045 (4)	0.069 (4)	-0.005 (3)	-0.003 (3)	-0.004 (3)
Co1	0.0543 (5)	0.0245 (4)	0.0478 (4)	0.0021 (3)	-0.0085 (3)	0.0006 (3)
O5	0.075 (3)	0.040 (2)	0.054 (2)	0.0000 (19)	-0.0150 (19)	-0.0013 (18)
N2	0.059 (3)	0.042 (3)	0.056 (3)	0.001 (2)	-0.002 (2)	-0.003 (2)
O2	0.133 (4)	0.046 (3)	0.088 (3)	0.019 (3)	-0.033 (3)	-0.024 (2)

O1	0.121 (4)	0.039 (2)	0.072 (3)	0.010 (2)	-0.026 (3)	-0.005 (2)
O3	0.074 (3)	0.048 (3)	0.107 (3)	0.005 (2)	0.022 (3)	-0.008 (2)
C6	0.055 (3)	0.040 (3)	0.059 (3)	0.007 (3)	-0.009 (3)	-0.008 (3)
C12	0.080 (4)	0.048 (4)	0.050 (3)	-0.005 (3)	-0.001 (3)	0.002 (3)
N1	0.055 (3)	0.040 (3)	0.053 (3)	0.001 (2)	-0.007 (2)	0.002 (2)
C3	0.076 (4)	0.066 (5)	0.068 (4)	0.010 (3)	-0.018 (3)	0.010 (3)
C7	0.071 (4)	0.045 (4)	0.066 (4)	0.009 (3)	-0.011 (3)	-0.013 (3)
C2	0.050 (3)	0.044 (3)	0.055 (3)	0.004 (3)	-0.001 (3)	0.004 (3)
C13	0.087 (4)	0.042 (3)	0.069 (4)	0.003 (3)	-0.027 (3)	0.003 (3)
C10	0.059 (3)	0.037 (3)	0.066 (4)	0.001 (3)	-0.013 (3)	-0.003 (3)
C1	0.064 (4)	0.041 (4)	0.062 (4)	0.007 (3)	-0.013 (3)	0.008 (3)
C5	0.086 (4)	0.047 (4)	0.066 (4)	0.006 (3)	-0.014 (3)	-0.008 (3)
C11	0.058 (3)	0.048 (4)	0.058 (3)	0.008 (3)	0.000 (3)	-0.003 (3)
C8	0.068 (4)	0.065 (4)	0.070 (4)	0.015 (3)	0.008 (3)	0.009 (3)
C4	0.088 (5)	0.065 (4)	0.066 (4)	-0.001 (4)	-0.029 (3)	-0.004 (3)
C9	0.068 (4)	0.056 (4)	0.072 (4)	0.015 (3)	0.006 (3)	0.012 (3)
C14	0.078 (4)	0.033 (3)	0.081 (4)	-0.004 (3)	-0.027 (3)	0.005 (3)
O1W	0.074 (3)	0.070 (3)	0.147 (4)	-0.008 (2)	-0.002 (3)	0.047 (3)
O2W	0.116 (4)	0.098 (4)	0.136 (5)	-0.009 (3)	-0.017 (3)	0.034 (4)
O3W	0.115 (4)	0.075 (4)	0.233 (7)	-0.015 (3)	-0.046 (4)	0.024 (4)

Geometric parameters (Å, °)

O4—C1	1.232 (7)	C12—C10	1.361 (7)
N3—C19	1.332 (6)	C12—C11	1.384 (7)
N3—C18	1.348 (6)	C12—H12	0.9300
N3—Co1 ⁱ	2.000 (4)	N1—C2	1.358 (6)
C19—C20	1.378 (7)	C3—C2	1.370 (7)
C19—H19	0.9300	C3—C4	1.377 (8)
C20—C16	1.379 (7)	C3—H3	0.9300
C20—H20	0.9300	C2—C1	1.518 (7)
C18—C17	1.378 (7)	C13—C10	1.524 (7)
C18—H18	0.9300	C13—C14	1.545 (7)
C17—C16	1.381 (7)	C13—H13A	0.9700
C17—H17	0.9300	C13—H13B	0.9700
C16—C15	1.519 (7)	C10—C9	1.371 (7)
C15—C14	1.492 (7)	C5—C4	1.368 (8)
C15—H15A	0.9700	C5—H5	0.9300
C15—H15B	0.9700	C11—H11	0.9300
Co1—O1	1.927 (4)	C8—C9	1.354 (7)
Co1—O5	1.932 (3)	C8—H8	0.9300
Co1—N2	1.994 (4)	C4—H4	0.9300
Co1—N3 ⁱⁱ	2.000 (4)	C9—H9	0.9300
Co1—O1W	2.184 (4)	C14—H14A	0.9700
O5—N1	1.344 (5)	C14—H14B	0.9700
N2—C11	1.334 (6)	O1W—H1W	0.8200
N2—C8	1.339 (6)	O1W—H2W	0.8200
O2—C7	1.224 (6)	O2W—H3W	0.8400
O1—C7	1.259 (6)	O2W—H4W	0.8400

supplementary materials

O3—C1	1.248 (7)	O3W—H6W	0.8200
C6—C5	1.361 (7)	O3W—H5W	0.8400
C6—N1	1.363 (6)	O3W—H6W	0.8200
C6—C7	1.522 (7)		
C19—N3—C18	116.1 (5)	C2—C3—C4	120.5 (6)
C19—N3—Co1 ⁱ	119.9 (4)	C2—C3—H3	119.7
C18—N3—Co1 ⁱ	123.9 (4)	C4—C3—H3	119.7
N3—C19—C20	123.9 (5)	O2—C7—O1	124.8 (6)
N3—C19—H19	118.0	O2—C7—C6	115.0 (5)
C20—C19—H19	118.0	O1—C7—C6	120.2 (5)
C16—C20—C19	120.1 (5)	N1—C2—C3	119.4 (5)
C16—C20—H20	119.9	N1—C2—C1	116.9 (4)
C19—C20—H20	119.9	C3—C2—C1	123.7 (5)
N3—C18—C17	122.9 (5)	C10—C13—C14	111.6 (4)
N3—C18—H18	118.5	C10—C13—H13A	109.3
C17—C18—H18	118.5	C14—C13—H13A	109.3
C18—C17—C16	120.6 (5)	C10—C13—H13B	109.3
C18—C17—H17	119.7	C14—C13—H13B	109.3
C16—C17—H17	119.7	H13A—C13—H13B	108.0
C20—C16—C17	116.3 (5)	C12—C10—C9	116.8 (5)
C20—C16—C15	121.0 (5)	C12—C10—C13	121.4 (5)
C17—C16—C15	122.7 (5)	C9—C10—C13	121.8 (5)
C14—C15—C16	115.6 (5)	O4—C1—O3	127.6 (6)
C14—C15—H15A	108	O4—C1—C2	117.4 (6)
C16—C15—H15A	108	O3—C1—C2	114.9 (5)
C14—C15—H15B	108	C6—C5—C4	122.7 (6)
C16—C15—H15B	108	C6—C5—H5	118.7
H15A—C15—H15B	107	C4—C5—H5	118.7
O1—Co1—O5	89.67 (16)	N2—C11—C12	121.5 (5)
O1—Co1—N2	86.43 (16)	N2—C11—H11	119.3
O5—Co1—N2	164.02 (17)	C12—C11—H11	119.3
O1—Co1—N3 ⁱⁱ	166.97 (19)	N2—C8—C9	123.4 (6)
O5—Co1—N3 ⁱⁱ	86.08 (16)	N2—C8—H8	118.3
N2—Co1—N3 ⁱⁱ	94.30 (16)	C9—C8—H8	118.3
O1—Co1—O1W	99.39 (19)	C5—C4—C3	117.8 (5)
O5—Co1—O1W	94.32 (16)	C5—C4—H4	121.1
N2—Co1—O1W	101.61 (17)	C3—C4—H4	121.1
N3 ⁱⁱ —Co1—O1W	93.21 (18)	C8—C9—C10	120.2 (6)
N1—O5—Co1	124.5 (3)	C8—C9—H9	119.9
C11—N2—C8	117.0 (5)	C10—C9—H9	119.9
C11—N2—Co1	122.3 (4)	C15—C14—C13	113.6 (5)
C8—N2—Co1	120.1 (4)	C15—C14—H14A	108.9
C7—O1—Co1	131.5 (4)	C13—C14—H14A	108.9
C5—C6—N1	117.8 (5)	C15—C14—H14B	108.9
C5—C6—C7	119.1 (5)	C13—C14—H14B	108.9
N1—C6—C7	123.0 (4)	H14A—C14—H14B	107.7
C10—C12—C11	121.0 (5)	Co1—O1W—H1W	118.8

C10—C12—H12	119.5	Co1—O1W—H2W	105.7
C11—C12—H12	119.5	H1W—O1W—H2W	110
O5—N1—C2	114.7 (4)	H3W—O2W—H4W	107
O5—N1—C6	123.5 (4)	H5W—O3W—H6W	112
C2—N1—C6	121.7 (4)		

Symmetry codes: (i) $x, y+1, z$; (ii) $x, y-1, z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1W—H1W \cdots O3 ⁱⁱⁱ	0.82	1.83	2.635 (5)	165
O1W—H2W \cdots O2W	0.82	1.99	2.732 (6)	150
O2W—H3W \cdots O3W ^{iv}	0.84	2.55	3.018 (9)	117
O2W—H4W \cdots O3	0.84	1.98	2.778 (6)	157
O3W—H5W \cdots O2	0.84	2.01	2.828 (7)	166
O3W—H6W \cdots O4 ⁱ	0.82	2.19	2.935 (7)	150
C3—H3 \cdots O4 ^v	0.93	2.46	3.364 (8)	165
C9—H9 \cdots O5 ^{vi}	0.93	2.46	3.366 (7)	164
C15—H15A \cdots O2 ^{vii}	0.97	2.57	3.492 (7)	159
C18—H18 \cdots O3W ^{vi}	0.93	2.55	3.369 (7)	149

Symmetry codes: (iii) $-x+1, y+1/2, -z+1/2$; (iv) $-x+1, y-1/2, -z+1/2$; (i) $x, y+1, z$; (v) $-x+1, -y, -z+1$; (vi) $-x, y+1/2, -z+1/2$; (vii) $x, -y+3/2, z-1/2$.

Fig. 1




Fig. 2

