

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bis{2-[(*E*)-benzyliminomethyl]-4,6-dibromophenolato- κ^2 N,O}cobalt(II)

Wei Jiang,* Gui-Di Mo and Lie Jin

School of Chemistry and Life Sciences, Maoming University, Guandu Second Road 139, Maoming 525000, Guangdong, People's Republic of China
Correspondence e-mail: gdmjw@yahoo.cn


Received 2 October 2008; accepted 7 October 2008

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.009$ Å; R factor = 0.052; wR factor = 0.172; data-to-parameter ratio = 18.4.

In the title compound, $[\text{Co}(\text{C}_{14}\text{H}_{10}\text{Br}_2\text{NO})_2]$, the Co^{II} ion is coordinated by an O and an N atom from two equivalent 2-[(*E*)-benzyliminomethyl]-4,6-dibromophenolate ligands, displaying a distorted tetrahedral geometry. The Co^{II} ion occupies a special position on a twofold rotation axis and thus the molecular symmetry of the complex is C_2 . The two phenolate rings are perpendicular [$89.8(3)^\circ$].

Related literature

For general background on the applications of Schiff bases, see: Vigato *et al.* (2007).



Experimental

Crystal data

$[\text{Co}(\text{C}_{14}\text{H}_{10}\text{Br}_2\text{NO})_2]$
 $M_r = 795.03$
Monoclinic, $C2/c$
 $a = 23.875(3)$ Å
 $b = 4.8190(6)$ Å
 $c = 24.209(3)$ Å
 $\beta = 105.8730(1)^\circ$

$V = 2679.1(6)$ Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 6.64$ mm⁻¹
 $T = 296(2)$ K
 $0.30 \times 0.26 \times 0.22$ mm

Data collection

Bruker SMART APEXII CCD diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 2000)
 $T_{\text{min}} = 0.165$, $T_{\text{max}} = 0.232$

11046 measured reflections
3094 independent reflections
2627 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$
 $wR(F^2) = 0.172$
 $S = 1.07$
3094 reflections

168 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.98$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.46$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Co1—O1	1.935 (4)	Co1—N1	2.005 (4)
O1 ⁱ —Co1—O1	126.8 (2)	O1—Co1—N1	113.29 (17)
O1 ⁱ —Co1—N1	94.16 (17)	N1—Co1—N1 ⁱ	117.1 (3)

Symmetry code: (i) $-x, y, -z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: XP in SHELXTL (Sheldrick, 2008); software used to prepare material for publication: XP in SHELXTL.

We are grateful to the Starting Fund for the Doctoral Program of Maoming University for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: KP2194).

References

- Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Sheldrick, G. M. (2000). SADABS. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Vigato, P.A., Tamburini, S. & Bertolo, L. (2007). *Coord. Chem. Rev.* **251**, 1311–1316.

supplementary materials

Acta Cryst. (2008). E64, m1394 [doi:10.1107/S1600536808032303]

Bis{2-[(*E*)-benzyliminomethyl]-4,6-dibromophenolato- κ^2N,O }cobalt(II)

W. Jiang, G.-D. Mo and L. Jin

Comment

The Schiff bases are widely employed as ligands in coordination chemistry. The advantages of Schiff bases enable their use in the synthesis of metal complexes of interest in bioinorganic chemistry, catalysis, encapsulation, transport and separation processes, and magnetochemistry (Choi & Jeon, 2003). Salicylaldehyde and its derivatives are useful carbonyl precursors for the synthesis of a large variety of Schiff bases. In this paper we report on a new cobalt(II) complex (I).

In the title complex Co^{II} atom is tetrahedrally coordinated by two O atoms and two N atoms from two 2-((*E*)-(benzylimino)methyl)-4,6-dibromophenol bidentate chelating ligand. The Co1—O1 distance of 1.935 (4) Å is shorter than the distance of Co1—N1 (2.005 (4) Å) (Table 1). The dihedral angle between two phenol rings is 89.8 (3)°.

Experimental

To a solution containing 2 mmol (0.738 g) 2-((*E*)-(benzylimino)methyl)-4,6-dibromophenol dissolved in 20 mL ethanol, 1 mmol of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ (0.238 g) were added, and the resulting mixture was stirred for about 1 h. The slow evaporation of the solvent after about 3 d yielded dark brown single crystals. Yield: 51.4%. Calcd. for $\text{C}_{28}\text{H}_{20}\text{Br}_4\text{CoN}_2\text{O}_2$: C, 42.30; H, 2.54; N, 3.52; Found: C, 42.24; H, 3.41; N, 3.46%.

Refinement

All H atoms were located from difference Fourier syntheses, H atoms from the C—H groups were placed in geometrically idealized positions and constrained to ride on their parent atoms (C—H = 0.93 Å, 0.96 Å, 0.97 Å) and $U_{\text{iso}}(\text{H})$ values equal to 1.2 $U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The structure of (I), showing displacement ellipsoids drawn at the 30% probability level. [Symmetry code: (i) $-x, y, -z+0.5$]

Bis{2-[(E)-benzyliminomethyl]-4,6-dibromophenolato- κ^2 N,O}cobalt(II)

Crystal data

[Co(C ₁₄ H ₁₀ Br ₂ NO) ₂]	$F_{000} = 1540$
$M_r = 795.03$	$D_x = 1.971 \text{ Mg m}^{-3}$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
Hall symbol: -C 2yc	$\lambda = 0.71073 \text{ \AA}$
$a = 23.875 (3) \text{ \AA}$	Cell parameters from 2269 reflections
$b = 4.8190 (6) \text{ \AA}$	$\theta = 1.0\text{--}27.7^\circ$
$c = 24.209 (3) \text{ \AA}$	$\mu = 6.64 \text{ mm}^{-1}$
$\beta = 105.8730 (10)^\circ$	$T = 296 (2) \text{ K}$
$V = 2679.1 (6) \text{ \AA}^3$	Block, brown
$Z = 4$	$0.30 \times 0.26 \times 0.22 \text{ mm}$

Data collection

Bruker SMART APEXII CCD diffractometer	3094 independent reflections
Radiation source: fine-focus sealed tube	2627 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.027$
$T = 296(2) \text{ K}$	$\theta_{\text{max}} = 27.7^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.8^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 2000)	$h = -30 \rightarrow 30$
$T_{\text{min}} = 0.165$, $T_{\text{max}} = 0.232$	$k = -6 \rightarrow 6$
11046 measured reflections	$l = -30 \rightarrow 31$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.052$	H-atom parameters constrained
$wR(F^2) = 0.172$	$w = 1/[\sigma^2(F_o^2) + (0.1045P)^2 + 20.5054P]$
$S = 1.07$	where $P = (F_o^2 + 2F_c^2)/3$
3094 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
168 parameters	$\Delta\rho_{\text{max}} = 0.98 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -1.45 \text{ e \AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.0000	0.9824 (2)	0.2500	0.0481 (3)
Br1	0.25733 (3)	0.73207 (15)	0.14570 (3)	0.0446 (2)
Br2	0.06423 (3)	1.45190 (13)	0.09822 (2)	0.0389 (2)
O1	0.04168 (16)	1.1622 (8)	0.20173 (16)	0.0316 (8)
N1	-0.06936 (19)	0.7651 (9)	0.20554 (18)	0.0269 (9)
C1	0.0898 (2)	1.0738 (10)	0.1927 (2)	0.0260 (10)
C2	0.1095 (2)	1.1793 (11)	0.1464 (2)	0.0285 (10)
C3	0.1595 (2)	1.0865 (12)	0.1339 (2)	0.0313 (11)
H3A	0.1707	1.1621	0.1032	0.038*
C4	0.1925 (2)	0.8813 (13)	0.1672 (2)	0.0321 (11)
C5	0.1772 (2)	0.7738 (12)	0.2136 (2)	0.0336 (11)
H5A	0.2005	0.6395	0.2364	0.040*
C6	0.1268 (2)	0.8649 (12)	0.2269 (2)	0.0281 (10)
C7	-0.1144 (2)	0.7334 (12)	0.2242 (2)	0.0309 (11)
H7A	-0.1425	0.6106	0.2037	0.037*
C8	-0.0702 (3)	0.6166 (12)	0.1512 (2)	0.0327 (11)
H8A	-0.0934	0.4492	0.1482	0.039*
H8B	-0.0309	0.5631	0.1517	0.039*
C9	-0.0954 (3)	0.7997 (11)	0.1000 (2)	0.0320 (11)
C10	-0.1555 (3)	0.8139 (16)	0.0759 (3)	0.0483 (16)
H10A	-0.1807	0.7055	0.0901	0.058*
C11	-0.1773 (4)	0.9955 (19)	0.0296 (3)	0.059 (2)
H11A	-0.2174	1.0061	0.0137	0.071*
C12	-0.1433 (4)	1.1523 (16)	0.0075 (3)	0.0555 (19)
H12A	-0.1593	1.2715	-0.0229	0.067*
C13	-0.0827 (4)	1.1356 (17)	0.0309 (3)	0.0533 (17)
H13A	-0.0581	1.2426	0.0157	0.064*
C14	-0.0597 (3)	0.9608 (14)	0.0764 (3)	0.0420 (14)
H14A	-0.0195	0.9504	0.0915	0.050*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.0443 (5)	0.0596 (6)	0.0428 (5)	0.000	0.0163 (4)	0.000
Br1	0.0289 (3)	0.0637 (4)	0.0467 (4)	0.0012 (2)	0.0197 (3)	-0.0043 (3)
Br2	0.0460 (4)	0.0389 (3)	0.0348 (3)	0.0015 (2)	0.0160 (3)	0.0099 (2)
O1	0.0320 (19)	0.039 (2)	0.0299 (19)	0.0049 (16)	0.0186 (16)	0.0067 (16)
N1	0.027 (2)	0.033 (2)	0.021 (2)	0.0010 (16)	0.0071 (17)	0.0003 (16)
C1	0.028 (2)	0.030 (2)	0.022 (2)	-0.0039 (19)	0.0092 (19)	-0.0012 (18)
C2	0.033 (3)	0.031 (3)	0.024 (2)	-0.006 (2)	0.011 (2)	-0.0022 (19)
C3	0.032 (3)	0.038 (3)	0.028 (3)	-0.010 (2)	0.014 (2)	-0.005 (2)
C4	0.026 (2)	0.042 (3)	0.032 (3)	-0.004 (2)	0.012 (2)	-0.006 (2)
C5	0.027 (3)	0.044 (3)	0.029 (3)	0.001 (2)	0.006 (2)	0.000 (2)
C6	0.026 (2)	0.038 (3)	0.022 (2)	-0.002 (2)	0.0078 (19)	-0.001 (2)
C7	0.027 (3)	0.042 (3)	0.023 (2)	-0.004 (2)	0.006 (2)	-0.003 (2)
C8	0.041 (3)	0.034 (3)	0.025 (2)	0.006 (2)	0.013 (2)	-0.002 (2)
C9	0.045 (3)	0.031 (3)	0.023 (2)	0.001 (2)	0.013 (2)	-0.0074 (19)
C10	0.043 (4)	0.061 (4)	0.042 (4)	0.007 (3)	0.014 (3)	0.004 (3)
C11	0.048 (4)	0.076 (5)	0.046 (4)	0.016 (4)	0.000 (3)	0.003 (4)
C12	0.084 (6)	0.051 (4)	0.025 (3)	0.005 (4)	0.005 (3)	0.001 (3)
C13	0.065 (5)	0.056 (4)	0.036 (3)	-0.011 (4)	0.010 (3)	0.006 (3)
C14	0.044 (3)	0.052 (4)	0.029 (3)	-0.007 (3)	0.008 (3)	0.000 (2)

Geometric parameters (\AA , $^\circ$)

Co1—O1 ⁱ	1.935 (4)	C6—C7 ⁱ	1.441 (7)
Co1—O1	1.935 (4)	C7—C6 ⁱ	1.441 (7)
Co1—N1	2.005 (4)	C7—H7A	0.9300
Co1—N1 ⁱ	2.005 (4)	C8—C9	1.505 (8)
Br1—C4	1.902 (5)	C8—H8A	0.9700
Br2—C2	1.888 (6)	C8—H8B	0.9700
O1—C1	1.298 (6)	C9—C14	1.388 (8)
N1—C7	1.285 (7)	C9—C10	1.396 (9)
N1—C8	1.493 (7)	C10—C11	1.404 (11)
C1—C2	1.424 (7)	C10—H10A	0.9300
C1—C6	1.442 (7)	C11—C12	1.323 (12)
C2—C3	1.383 (7)	C11—H11A	0.9300
C3—C4	1.379 (8)	C12—C13	1.405 (11)
C3—H3A	0.9300	C12—H12A	0.9300
C4—C5	1.376 (8)	C13—C14	1.375 (10)
C5—C6	1.399 (8)	C13—H13A	0.9300
C5—H5A	0.9300	C14—H14A	0.9300
O1 ⁱ —Co1—O1	126.8 (2)	N1—C7—C6 ⁱ	127.9 (5)
O1 ⁱ —Co1—N1	94.16 (17)	N1—C7—H7A	116.0
O1—Co1—N1	113.29 (17)	C6 ⁱ —C7—H7A	116.0
O1 ⁱ —Co1—N1 ⁱ	113.29 (17)	N1—C8—C9	110.5 (4)

O1—Co1—N1 ⁱ	94.16 (17)	N1—C8—H8A	109.6
N1—Co1—N1 ⁱ	117.1 (3)	C9—C8—H8A	109.6
C1—O1—Co1	125.4 (3)	N1—C8—H8B	109.6
C7—N1—C8	116.2 (5)	C9—C8—H8B	109.6
C7—N1—Co1	121.4 (4)	H8A—C8—H8B	108.1
C8—N1—Co1	122.3 (4)	C14—C9—C10	118.5 (6)
O1—C1—C2	120.9 (5)	C14—C9—C8	121.1 (6)
O1—C1—C6	124.4 (4)	C10—C9—C8	120.5 (6)
C2—C1—C6	114.7 (4)	C9—C10—C11	118.6 (7)
C3—C2—C1	123.3 (5)	C9—C10—H10A	120.7
C3—C2—Br2	118.2 (4)	C11—C10—H10A	120.7
C1—C2—Br2	118.6 (4)	C12—C11—C10	123.0 (7)
C4—C3—C2	119.6 (5)	C12—C11—H11A	118.5
C4—C3—H3A	120.2	C10—C11—H11A	118.5
C2—C3—H3A	120.2	C11—C12—C13	118.8 (7)
C5—C4—C3	120.6 (5)	C11—C12—H12A	120.6
C5—C4—Br1	120.0 (5)	C13—C12—H12A	120.6
C3—C4—Br1	119.3 (4)	C14—C13—C12	120.0 (7)
C4—C5—C6	120.6 (5)	C14—C13—H13A	120.0
C4—C5—H5A	119.7	C12—C13—H13A	120.0
C6—C5—H5A	119.7	C13—C14—C9	121.2 (7)
C5—C6—C7 ⁱ	115.6 (5)	C13—C14—H14A	119.4
C5—C6—C1	121.2 (5)	C9—C14—H14A	119.4
C7 ⁱ —C6—C1	123.2 (5)		

Symmetry codes: (i) $-x, y, -z+1/2$.

Fig. 1

