

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

**(*O,O'*-Diethyl dithiophosphato- $\kappa^2S,S'$ )-  
(hydridotripyrazol-1-ylborato- $\kappa^3N^2,N^2',N^2''$ )(triphenylphosphine- $\kappa P$ )-  
ruthenium(II)**

Hung-Chun Tong,<sup>a</sup> Chih-Yung Chen Hsu,<sup>a</sup> Yao-Ren Liang,<sup>a</sup>  
Yih Hsing Lo<sup>a\*</sup> and Chia-Her Lin<sup>b</sup>

<sup>a</sup>Department of Chemical Engineering, Tatung University, Taipei 104, Taiwan, and

<sup>b</sup>Department of Chemistry, Chung-Yuan Christian University, Chung-Li 320, Taiwan

Correspondence e-mail: yhlo@ttu.edu.tw


Received 16 September 2008; accepted 1 October 2008

Key indicators: single-crystal X-ray study;  $T = 200$  K; mean  $\sigma(C-C) = 0.005$  Å; disorder in main residue;  $R$  factor = 0.036;  $wR$  factor = 0.093; data-to-parameter ratio = 14.0.

Reaction of  $[Ru(Tp)Cl(PPh_3)_2]$  {where Tp is hydridotripyrazolylborate,  $BH[C_3H_3N_2)_3]$ } with  $NH_4[S_2P(OEt)_2]$  in methanol afforded the title compound,  $[Ru(C_9H_{10}BN_6)(C_4H_{10}O_2PS_2)(C_{18}H_{15}P)]$ , in which the  $Ru^{II}$  ion is in a slightly distorted octahedral coordination environment. The  $[S_2P(OEt)_2]^-$  ligand coordinates in a chelating mode with two similar  $Ru-S$  bond lengths and a slightly acute  $S-Ru-S$  angle. The atoms of both  $-OCH_2CH_3$  groups of the diethyl dithiophosphate ligand are disordered over two sites with approximate occupancies of 0.76 and 0.24.

## Related literature

For related structures, see: Alock *et al.* (1992); Burrows (2001); Hidai *et al.* (2000); Gemel *et al.* (1996); Jain & Jakkal (1996); Meno *et al.* (1995); Pavlik *et al.* (2005); Sellmann *et al.* (1999); Slugovc *et al.* (1998); Vit & Zdrzil (1989).



## Experimental

### Crystal data

$[Ru(C_9H_{10}BN_6)(C_4H_{10}O_2PS_2)-$   
 $(C_{18}H_{15}P)]$   
 $M_r = 761.59$

Monoclinic,  $P2_1/c$   
 $a = 12.4408$  (2) Å  
 $b = 13.7386$  (2) Å

$c = 20.3775$  (3) Å  
 $\beta = 99.676$  (1)°  
 $V = 3433.36$  (9) Å<sup>3</sup>  
 $Z = 4$

Mo  $K\alpha$  radiation  
 $\mu = 0.71$  mm<sup>-1</sup>  
 $T = 200$  (2) K  
 $0.42 \times 0.3 \times 0.15$  mm

### Data collection

Nonius KappaCCD diffractometer  
Absorption correction: multi-scan  
(*SORTAV*; Blessing, 1995)  
 $T_{min} = 0.755$ ,  $T_{max} = 0.901$

24009 measured reflections  
6265 independent reflections  
5601 reflections with  $I > 2\sigma(I)$   
 $R_{int} = 0.051$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$   
 $wR(F^2) = 0.093$   
 $S = 1.05$   
6265 reflections  
449 parameters

42 restraints  
H-atom parameters constrained  
 $\Delta\rho_{max} = 1.43$  e Å<sup>-3</sup>  
 $\Delta\rho_{min} = -0.62$  e Å<sup>-3</sup>

**Table 1**

Selected geometric parameters (Å, °).

Ru1—N3	2.086 (2)	Ru1—P1	2.3171 (8)
Ru1—N1	2.088 (2)	Ru1—S1	2.4540 (8)
Ru1—N5	2.144 (3)	Ru1—S2	2.4635 (8)
N3—Ru1—N1	90.09 (9)	N5—Ru1—S1	87.41 (7)
N3—Ru1—N5	83.57 (10)	P1—Ru1—S1	99.23 (3)
N1—Ru1—N5	84.88 (10)	N3—Ru1—S2	94.61 (7)
N3—Ru1—P1	90.07 (7)	N1—Ru1—S2	169.78 (7)
N1—Ru1—P1	90.63 (7)	N5—Ru1—S2	86.63 (7)
N5—Ru1—P1	172.20 (7)	P1—Ru1—S2	98.42 (3)
N3—Ru1—S1	170.14 (7)	S1—Ru1—S2	80.86 (3)
N1—Ru1—S1	93.07 (7)		

Data collection: *COLLECT* (Nonius, 1999); cell refinement: *DENZO* and *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* and *SCALEPACK*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

This research was supported by the National Science Council, Taiwan (NSC 97-2113-M-036-001-MY2), and in part by the project of the specific research fields in Tatung University, Taiwan (B96-C07-081), and the project of the specific research fields in Chung Yuan Christian University, Taiwan, under grant No. CYCU-97-CR-CH. We also thank Mr Ting Shen Kuo (Department of Chemistry, National Taiwan Normal University) for his assistance in X-ray single-crystal structure analysis.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2697).

## References

- Alock, N. W., Burns, I. D., Claire, K. S. & Hill, A. F. (1992). *Inorg. Chem.* **31**, 2906–2908.  
Blessing, R. H. (1995). *Acta Cryst.* **A51**, 33–38.  
Burrows, A. D. (2001). *CrystEngComm*, **46**, 1–5.  
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.  
Gemel, C., Trimmel, G., Slugovc, C., Kremel, S., Mereiter, K., Schmid, R. & Kirchner, K. (1996). *Organometallics*, **15**, 3998–4004.

- Hidai, M., Kuwata, S. & Mizobe, Y. (2000). *Acc. Chem. Res.* **33**, 46–52.
- Jain, V. K. & Jakkal, V. S. (1996). *J. Organomet. Chem.* **515**, 81–87.
- Meno, M., Pramanik, A., Bag, N. & Chakravorty, A. (1995). *J. Chem. Soc. Dalton Trans.* pp. 1543–1547.
- Nonius (1999). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography, Part A*, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Pavlik, S., Mereiter, K., Puchberger, M. & Kirchner, K. (2005). *Organometallics*, **24**, 3561–3575.
- Sellmann, D., Utz, J. & Heineman, F. W. (1999). *Eur. J. Inorg. Chem.* **2**, 341–346.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Slugovc, C., Mereiter, K., Schmidt, R. & Kirchner, K. (1998). *Organometallics*, **17**, 827–831.
- Vit, Z. & Zdrzil, M. (1989). *J. Catal.* **119**, 1–9.

**supplementary materials**

*Acta Cryst.* (2008). E64, m1372-m1373 [ doi:10.1107/S160053680803170X ]

**(*O,O'*-Diethyl dithiophosphato- $\kappa^2$ S,S')(hydridotripyrazol-1-ylborato- $\kappa^3$ N<sup>2</sup>,N<sup>2'</sup>,N<sup>2''</sup>)(triphenylphosphine- $\kappa$ P)ruthenium(II)**

**H.-C. Tong, C.-Y. Chen Hsu, Y.-R. Liang, Y. H. Lo and C.-H. Lin**

**Comment**

The chemistry of transition metal sulfur compounds has attracted interest for their importance in the field of catalysts, metalloenzymes, and material precursor (Hidai *et al.*, 2000). In recent years there has been an increased interest in ruthenium sulfur complexes, in part because of the high catalytic activity of RuS<sub>2</sub> in various hydrotreating processes (Vit & Zdrzil, 1989). As a part of this development, many examples of ruthenium thiolate complexes have been reported, however, the ruthenium complexes with dithio ligands are relatively rare (Sellmann *et al.*, 1999). On the other hand, ruthenium(II)hydridotripyrazolylborate complexes, Ru(Tp), are of interest for stoichiometric and catalytic transformations of organic molecules (Pavlik *et al.*, 2005). The complex [Ru(Tp)Cl(PPh<sub>3</sub>)<sub>2</sub>] (Alock *et al.*, 1992) has been used as the starting material for the synthesis of several complexes because of its substitutionally labile chloride and phosphines (Burrows, 2001). In order to acquire a better understanding of the coordination chemistry of RuS<sub>2</sub>, we have studied the ruthenium phosphine complex containing the ligands hydrotris(pyrazolyl)borate (Tp) and [NH<sub>4</sub>][S<sub>2</sub>P(OEt)<sub>2</sub>]. Interaction of [Ru(Tp)Cl(PPh<sub>3</sub>)<sub>2</sub>] with [NH<sub>4</sub>][S<sub>2</sub>P(OEt)<sub>2</sub>] in MeOH afforded the title compound {Ru(Tp)(PPh<sub>3</sub>)[S<sub>2</sub>P(OEt)<sub>2</sub>]} (I). The <sup>31</sup>P NMR spectrum of (I) in CDCl<sub>3</sub> shows two intense singlets at 50.8 and 105.7 p.p.m., assignable to PPh<sub>3</sub> and [S<sub>2</sub>P(OEt)<sub>2</sub>], respectively. The FAB mass spectrum of (I) shows the molecular ions {Ru(Tp)(PPh<sub>3</sub>)[S<sub>2</sub>P(OEt)<sub>2</sub>]} with the characteristic isotopic distribution patterns. The crystal structure of (I) was established by X-ray crystallography. In the title compound, the environment about the Ru<sup>II</sup> ion is slightly distorted octahedral and the bite angle of the Tp ligand produces an average N—Ru—N angle of ca. 86° only slightly distorted from 90°. The three Ru—N(Tp) bond lengths are slightly longer than the average distance of 2.038 Å in other ruthenium Tp complexes (Gemel *et al.* 1996; Slugovc *et al.* 1998). The [S<sub>2</sub>P(OEt)<sub>2</sub>]<sup>−</sup> ligand chelates the ruthenium centre with two nearly equal Ru—S bonds and the S—Ru—S angle is slightly acute. The Ru—S bond lengths in (I) are comparable to those in [(η<sup>6</sup>-p-cymene)Ru{S<sub>2</sub>P(OMe)<sub>2</sub>}(PPh<sub>3</sub>)] [BPh<sub>4</sub>][av. 2.4311 (12) Å] with a chelated dithiophosphate ligand (Jain & Jakkal, 1996), but slightly longer than for *cis*-[Ru(S<sub>2</sub>CNEt<sub>2</sub>)<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>][av. 2.3952 (5) Å] with chelated dithiocarbamate (Meno *et al.*, 1995). The Ru—P bond length in (I) agrees well with those in related ruthenium(II) complexes with PPh<sub>3</sub> ligands (Jain & Jakkal, 1996, Meno *et al.*, 1995).

**Experimental**

The synthesis of the title compound (I) was carried out as follows. To a solution of [Ru(Tp)Cl(PPh<sub>3</sub>)<sub>2</sub>] (3.95 g, 4.50 mmol) in MeOH (20 ml), an excess of [NH<sub>4</sub>][S<sub>2</sub>P(OEt)<sub>2</sub>] (1.82 g, 9.00 mmol) were added. The reaction mixture was stirred for a further 8 h at room temperature. The solvent was dried under vacuum and 20 ml of CH<sub>2</sub>Cl<sub>2</sub> was added to the residue. The product was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and other salts such as [NH<sub>4</sub>][S<sub>2</sub>P(OEt)<sub>2</sub>] and NH<sub>4</sub>Cl precipitated. After filtration, the solvent was dried under vacuum to give the title compound (I) (3.27 g, 95% yield). Spectroscopic analysis: IR (KBr,

## supplementary materials

cm<sup>-1</sup>):  $\nu(\text{BH})$  2468 cm<sup>-1</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 303 K, d, p.p.m.): d 7.92 (d,  $J_{\text{H-H}} = 2.3$  Hz, 1H, Tp), 7.83 (d,  $J_{\text{H-H}} = 2.3$  Hz, 1H, Tp), 7.71 (d,  $J_{\text{H-H}} = 2.3$  Hz, 1H, Tp), 7.4–6.9 (m, Tp, Ph), 6.83 (d,  $J_{\text{H-H}} = 2.3$  Hz, 1H, Tp), 5.81 (d,  $J_{\text{H-H}} = 2.2$  Hz, 1H, Tp), 5.66 (d,  $J_{\text{H-H}} = 2.2$  Hz, 1H, Tp), 5.63 (t,  $J_{\text{H-H}} = 2.2$  Hz, 1H, Tp), 5.54 (t,  $J_{\text{H-H}} = 2.2$  Hz, 1H, Tp), 4.16 (q,  $J_{\text{H-H}} = 7.2$  Hz, 2H, OCH<sub>2</sub>), 3.11 (q,  $J_{\text{H-H}} = 7.2$  Hz, 2H, OCH<sub>2</sub>), 1.32 (t,  $J_{\text{H-H}} = 7.2$  Hz, 3H, CH<sub>3</sub>), 0.79 (t,  $J_{\text{H-H}} = 7.2$  Hz, 3H, CH<sub>3</sub>). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 303 K, d, p.p.m.): 146.7–104.6 (m, PPh<sub>3</sub>, Tp), 60.6, 61.4 (d, OCH<sub>2</sub>,  $^2J_{\text{P-C}} = 10$  Hz), 15.5, 15.9 (d, OCH<sub>2</sub>CH<sub>3</sub>,  $^3J_{\text{P-C}} = 8.4$  Hz). <sup>31</sup>P NMR (CDCl<sub>3</sub>, 303 K, d, p.p.m.): d 105.7 (PS<sub>2</sub>), 50.9 (s, PPh<sub>3</sub>). MS ( $m/z$ , Ru102): 762.2 ( $M^+$ ), 500.1 ( $M^+ - \text{PPh}_3$ ). Anal. Calcd for C<sub>31</sub>H<sub>35</sub>BN<sub>6</sub>O<sub>2</sub>P<sub>2</sub>RuS<sub>2</sub>: C, 48.89; H, 4.63; N, 11.03. Found: C, 48.73; H, 4.61; N, 11.02. The bright-yellow crystals of (I) for X-ray structure analysis were obtained by recrystallization of the crude product from dichloromethane-hexane.

### Refinement

H atoms were placed in idealized positions and constrained to ride on their parent atoms, with C—H = 0.93 - 0.97 Å and  $U_{\text{iso}}(\text{H}) = 1.2$  or  $1.5U_{\text{eq}}(\text{C})$ , B—H = 0.98 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . The atoms of both -OCH<sub>2</sub>CH<sub>3</sub> groups of the diethyldithiophosphato ligand are disordered over two sites with refined occupancies of 0.764 (3) and 0.236 (3).

### Figures




Fig. 1. Molecular structure of (I) showing displacement ellipsoids at the 35% level and H atoms having arbitrary radius. The disorder is not shown.

### (*O,O'*-Diethyl dithiophosphato- $\kappa^2S,S'$ )(hydridotripyrazol-1-ylborato- $\kappa^3N^2,N^2',N^2''$ )(triphenylphosphine- $\kappa P$ )ruthenium(II)

#### Crystal data

[Ru(C<sub>9</sub>H<sub>10</sub>BN<sub>6</sub>)(C<sub>4</sub>H<sub>10</sub>O<sub>2</sub>PS<sub>2</sub>)(C<sub>18</sub>H<sub>15</sub>P)]

$M_r = 761.59$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 12.4408$  (2) Å

$b = 13.7386$  (2) Å

$c = 20.3775$  (3) Å

$\beta = 99.676$  (1)°

$V = 3433.36$  (9) Å<sup>3</sup>

$Z = 4$

$F_{000} = 1560$

$D_x = 1.473$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation

$\lambda = 0.71073$  Å

Cell parameters from 32408 reflections

$\theta = 2.0$ – $25.4$ °

$\mu = 0.71$  mm<sup>-1</sup>

$T = 200$  (2) K

Prism, yellow

$0.42 \times 0.3 \times 0.15$  mm

*Data collection*

Nonius KappaCCD diffractometer	6265 independent reflections
Radiation source: fine-focus sealed tube	5601 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.051$
Detector resolution: 9 pixels $\text{mm}^{-1}$	$\theta_{\text{max}} = 25.4^\circ$
$T = 200(2)$ K	$\theta_{\text{min}} = 2.4^\circ$
CCD rotation images, thick slices scans	$h = -11 \rightarrow 14$
Absorption correction: multi-scan (SORTAV; Blessing, 1995)	$k = -16 \rightarrow 16$
$T_{\text{min}} = 0.755$ , $T_{\text{max}} = 0.901$	$l = -24 \rightarrow 24$
24009 measured reflections	

*Refinement*

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.036$	H-atom parameters constrained
$wR(F^2) = 0.093$	$w = 1/[\sigma^2(F_o^2) + (0.0373P)^2 + 5.0939P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
6265 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
449 parameters	$\Delta\rho_{\text{max}} = 1.43 \text{ e } \text{\AA}^{-3}$
42 restraints	$\Delta\rho_{\text{min}} = -0.62 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 \geq \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Ru1	0.685497 (18)	0.068530 (16)	0.255762 (11)	0.02384 (9)	
S1	0.79384 (7)	0.17427 (6)	0.19498 (4)	0.0388 (2)	
S2	0.74769 (7)	-0.05385 (6)	0.18210 (4)	0.03638 (19)	

## supplementary materials

---

P1	0.81369 (6)	0.03972 (6)	0.35016 (4)	0.02612 (17)
P2	0.84695 (7)	0.05178 (7)	0.16235 (4)	0.0369 (2)
N1	0.6227 (2)	0.18566 (18)	0.30252 (12)	0.0289 (5)
N2	0.5136 (2)	0.18862 (18)	0.30454 (13)	0.0324 (6)
N3	0.57667 (19)	-0.02517 (18)	0.29151 (12)	0.0272 (5)
N4	0.47495 (19)	0.00904 (18)	0.29637 (13)	0.0301 (6)
N5	0.5506 (2)	0.09368 (19)	0.17715 (13)	0.0314 (6)
N6	0.4510 (2)	0.11145 (19)	0.19433 (14)	0.0339 (6)
C1	0.6649 (3)	0.2667 (2)	0.33239 (16)	0.0351 (7)
H1A	0.7383	0.2835	0.3384	0.042*
C2	0.5841 (3)	0.3225 (3)	0.35311 (19)	0.0454 (9)
H2A	0.5922	0.3822	0.3749	0.054*
C3	0.4901 (3)	0.2709 (2)	0.33460 (17)	0.0406 (8)
H3A	0.4212	0.2897	0.3417	0.049*
C4	0.5802 (3)	-0.1169 (2)	0.31240 (15)	0.0307 (7)
H4A	0.6401	-0.1579	0.3140	0.037*
C5	0.4826 (3)	-0.1432 (2)	0.33140 (17)	0.0396 (8)
H5A	0.4647	-0.2030	0.3480	0.048*
C6	0.4181 (3)	-0.0622 (2)	0.32047 (17)	0.0371 (7)
H6A	0.3469	-0.0571	0.3284	0.045*
C7	0.5363 (3)	0.0975 (3)	0.11105 (17)	0.0410 (8)
H7A	0.5909	0.0872	0.0858	0.049*
C8	0.4285 (3)	0.1189 (3)	0.08504 (19)	0.0493 (9)
H8A	0.3978	0.1263	0.0405	0.059*
C9	0.3772 (3)	0.1267 (3)	0.13905 (18)	0.0445 (9)
H9A	0.3037	0.1403	0.1378	0.053*
C10	0.7727 (2)	0.0899 (2)	0.42642 (15)	0.0295 (6)
C11	0.8439 (3)	0.1437 (2)	0.47297 (15)	0.0351 (7)
H11A	0.9154	0.1535	0.4667	0.042*
C12	0.8085 (3)	0.1827 (3)	0.52858 (17)	0.0419 (8)
H12A	0.8566	0.2186	0.5593	0.050*
C13	0.7026 (3)	0.1686 (3)	0.53855 (17)	0.0437 (8)
H13A	0.6789	0.1958	0.5754	0.052*
C14	0.6322 (3)	0.1137 (3)	0.49359 (17)	0.0427 (8)
H14A	0.5614	0.1026	0.5009	0.051*
C15	0.6662 (3)	0.0751 (2)	0.43786 (16)	0.0354 (7)
H15A	0.6177	0.0389	0.4077	0.042*
C16	0.8413 (2)	-0.0886 (2)	0.37420 (16)	0.0300 (6)
C17	0.8813 (3)	-0.1501 (2)	0.32978 (17)	0.0389 (8)
H17A	0.8871	-0.1272	0.2876	0.047*
C18	0.9128 (3)	-0.2447 (3)	0.34695 (19)	0.0441 (8)
H18A	0.9395	-0.2845	0.3165	0.053*
C19	0.9043 (3)	-0.2796 (2)	0.40899 (19)	0.0446 (9)
H19A	0.9265	-0.3427	0.4210	0.054*
C20	0.8628 (3)	-0.2207 (3)	0.45330 (19)	0.0464 (9)
H20A	0.8553	-0.2448	0.4949	0.056*
C21	0.8320 (3)	-0.1256 (2)	0.43639 (17)	0.0387 (8)
H21A	0.8049	-0.0863	0.4670	0.046*
C22	0.9557 (2)	0.0851 (2)	0.35834 (15)	0.0314 (7)

C23	1.0452 (3)	0.0291 (3)	0.38653 (17)	0.0386 (8)	
H23A	1.0344	-0.0335	0.4015	0.046*	
C24	1.1505 (3)	0.0660 (3)	0.39243 (19)	0.0470 (9)	
H24A	1.2096	0.0273	0.4102	0.056*	
C25	1.1676 (3)	0.1587 (3)	0.37228 (19)	0.0515 (10)	
H25A	1.2381	0.1833	0.3767	0.062*	
C26	1.0798 (3)	0.2161 (3)	0.34532 (19)	0.0494 (9)	
H26A	1.0912	0.2794	0.3318	0.059*	
C27	0.9746 (3)	0.1791 (3)	0.33844 (17)	0.0406 (8)	
H27A	0.9160	0.2180	0.3202	0.049*	
O1	0.9685 (2)	0.0388 (3)	0.20330 (16)	0.0485 (8)	0.764 (3)
C28	1.0361 (4)	-0.0397 (4)	0.1920 (3)	0.0657 (16)	0.764 (3)
H28A	1.0254	-0.0524	0.1446	0.079*	0.764 (3)
H28B	1.0124	-0.0971	0.2134	0.079*	0.764 (3)
C29	1.1497 (4)	-0.0264 (6)	0.2152 (5)	0.079 (2)	0.764 (3)
H29A	1.1884	-0.0839	0.2059	0.118*	0.764 (3)
H29B	1.1619	-0.0149	0.2623	0.118*	0.764 (3)
H29C	1.1753	0.0283	0.1929	0.118*	0.764 (3)
O1A	0.9713 (4)	0.0320 (9)	0.1572 (5)	0.0479 (13)	0.236 (3)
C29A	1.1270 (19)	-0.061 (2)	0.2317 (18)	0.079 (2)	0.236 (3)
H29D	1.1592	-0.0565	0.2779	0.118*	0.236 (3)
H29E	1.1827	-0.0527	0.2047	0.118*	0.236 (3)
H29F	1.0928	-0.1229	0.2229	0.118*	0.236 (3)
C28A	1.0470 (15)	0.0148 (15)	0.2162 (8)	0.0648 (19)	0.236 (3)
H28C	1.0029	0.0108	0.2511	0.078*	0.236 (3)
H28D	1.0878	0.0750	0.2241	0.078*	0.236 (3)
O2	0.8764 (3)	0.0499 (3)	0.08958 (13)	0.0458 (9)	0.764 (3)
C30	0.7920 (4)	0.0556 (4)	0.0330 (2)	0.0476 (13)	0.764 (3)
H30A	0.7472	0.1124	0.0366	0.057*	0.764 (3)
H30B	0.7459	-0.0017	0.0311	0.057*	0.764 (3)
C31	0.8393 (18)	0.0618 (10)	-0.0271 (6)	0.068 (3)	0.764 (3)
H31A	0.7821	0.0649	-0.0650	0.102*	0.764 (3)
H31B	0.8835	0.0054	-0.0305	0.102*	0.764 (3)
H31C	0.8835	0.1193	-0.0255	0.102*	0.764 (3)
O2A	0.8154 (9)	0.0724 (8)	0.0839 (2)	0.0467 (13)	0.236 (3)
C30A	0.8201 (17)	0.0081 (12)	0.0299 (7)	0.0487 (17)	0.236 (3)
H30C	0.7493	-0.0230	0.0193	0.058*	0.236 (3)
H30D	0.8721	-0.0426	0.0462	0.058*	0.236 (3)
C31A	0.848 (7)	0.042 (4)	-0.032 (2)	0.068 (3)	0.236 (3)
H31D	0.8431	-0.0104	-0.0630	0.102*	0.236 (3)
H31E	0.9207	0.0677	-0.0241	0.102*	0.236 (3)
H31F	0.7979	0.0929	-0.0497	0.102*	0.236 (3)
B1	0.4383 (3)	0.1100 (3)	0.26828 (19)	0.0346 (8)	
H1	0.3624	0.1224	0.2729	0.042*	

Atomic displacement parameters ( $\text{\AA}^2$ )

$U^{11}$   $U^{22}$   $U^{33}$   $U^{12}$   $U^{13}$   $U^{23}$

## supplementary materials

---

Ru1	0.01928 (13)	0.02232 (13)	0.03081 (14)	0.00061 (9)	0.00673 (9)	0.00169 (9)
S1	0.0393 (5)	0.0347 (4)	0.0454 (5)	-0.0073 (4)	0.0159 (4)	0.0054 (4)
S2	0.0374 (5)	0.0309 (4)	0.0439 (4)	0.0032 (3)	0.0155 (4)	-0.0018 (3)
P1	0.0202 (4)	0.0268 (4)	0.0319 (4)	0.0005 (3)	0.0060 (3)	0.0010 (3)
P2	0.0284 (4)	0.0498 (5)	0.0346 (4)	-0.0018 (4)	0.0116 (3)	-0.0028 (4)
N1	0.0237 (13)	0.0267 (13)	0.0369 (13)	0.0020 (10)	0.0065 (10)	0.0017 (11)
N2	0.0264 (13)	0.0298 (13)	0.0422 (14)	0.0062 (11)	0.0093 (11)	0.0007 (11)
N3	0.0212 (12)	0.0268 (13)	0.0352 (13)	0.0004 (10)	0.0087 (10)	-0.0018 (10)
N4	0.0188 (12)	0.0309 (13)	0.0418 (14)	-0.0007 (10)	0.0091 (10)	0.0009 (11)
N5	0.0249 (13)	0.0314 (14)	0.0374 (14)	0.0034 (11)	0.0036 (11)	0.0022 (11)
N6	0.0230 (13)	0.0311 (14)	0.0460 (15)	0.0043 (11)	0.0007 (11)	0.0036 (12)
C1	0.0347 (18)	0.0277 (16)	0.0428 (18)	-0.0022 (14)	0.0059 (14)	-0.0026 (14)
C2	0.050 (2)	0.0291 (17)	0.057 (2)	0.0076 (16)	0.0108 (17)	-0.0077 (16)
C3	0.0376 (19)	0.0357 (18)	0.050 (2)	0.0108 (15)	0.0133 (15)	-0.0032 (15)
C4	0.0298 (16)	0.0252 (15)	0.0382 (16)	-0.0014 (13)	0.0084 (13)	0.0010 (13)
C5	0.0370 (19)	0.0324 (17)	0.052 (2)	-0.0065 (15)	0.0159 (15)	0.0062 (15)
C6	0.0289 (17)	0.0407 (19)	0.0445 (18)	-0.0050 (14)	0.0137 (14)	0.0022 (15)
C7	0.042 (2)	0.0434 (19)	0.0359 (17)	0.0025 (16)	0.0016 (15)	0.0014 (15)
C8	0.046 (2)	0.054 (2)	0.043 (2)	-0.0011 (18)	-0.0077 (17)	0.0042 (17)
C9	0.0290 (18)	0.044 (2)	0.055 (2)	0.0008 (15)	-0.0078 (16)	0.0037 (17)
C10	0.0292 (16)	0.0264 (15)	0.0337 (16)	0.0032 (13)	0.0071 (12)	0.0046 (12)
C11	0.0312 (17)	0.0372 (18)	0.0366 (16)	-0.0011 (14)	0.0044 (13)	-0.0006 (14)
C12	0.046 (2)	0.0407 (19)	0.0390 (18)	-0.0037 (16)	0.0087 (15)	-0.0093 (15)
C13	0.051 (2)	0.045 (2)	0.0383 (18)	0.0001 (17)	0.0157 (16)	-0.0076 (15)
C14	0.0345 (19)	0.052 (2)	0.0451 (19)	0.0007 (16)	0.0173 (15)	-0.0001 (16)
C15	0.0321 (17)	0.0408 (18)	0.0333 (16)	-0.0035 (14)	0.0059 (13)	-0.0023 (14)
C16	0.0194 (14)	0.0302 (16)	0.0390 (16)	-0.0006 (12)	0.0008 (12)	0.0025 (13)
C17	0.0341 (18)	0.0380 (18)	0.0455 (19)	0.0100 (15)	0.0091 (15)	0.0048 (15)
C18	0.0373 (19)	0.0357 (18)	0.059 (2)	0.0068 (15)	0.0070 (16)	-0.0009 (16)
C19	0.0356 (19)	0.0291 (17)	0.066 (2)	0.0029 (15)	-0.0017 (17)	0.0054 (16)
C20	0.054 (2)	0.0369 (19)	0.047 (2)	-0.0029 (17)	0.0046 (17)	0.0127 (16)
C21	0.0406 (19)	0.0347 (18)	0.0403 (18)	-0.0024 (15)	0.0053 (15)	0.0012 (14)
C22	0.0236 (15)	0.0388 (17)	0.0331 (16)	-0.0042 (13)	0.0082 (12)	-0.0025 (13)
C23	0.0268 (17)	0.0458 (19)	0.0428 (18)	0.0000 (15)	0.0045 (14)	0.0017 (15)
C24	0.0233 (17)	0.067 (3)	0.051 (2)	-0.0018 (16)	0.0071 (15)	-0.0041 (18)
C25	0.0287 (19)	0.077 (3)	0.051 (2)	-0.0171 (19)	0.0119 (16)	-0.010 (2)
C26	0.046 (2)	0.052 (2)	0.052 (2)	-0.0224 (18)	0.0136 (17)	-0.0027 (17)
C27	0.0346 (18)	0.0390 (19)	0.0474 (19)	-0.0065 (15)	0.0049 (15)	-0.0012 (15)
O1	0.0291 (16)	0.072 (2)	0.0457 (19)	0.0045 (16)	0.0094 (15)	-0.0087 (18)
C28	0.053 (3)	0.060 (4)	0.080 (4)	0.007 (3)	0.001 (3)	-0.006 (3)
C29	0.034 (3)	0.103 (7)	0.103 (6)	0.015 (4)	0.018 (3)	-0.009 (5)
O1A	0.029 (2)	0.070 (3)	0.046 (3)	0.005 (2)	0.012 (2)	-0.007 (3)
C29A	0.034 (3)	0.103 (7)	0.103 (6)	0.015 (4)	0.018 (3)	-0.009 (5)
C28A	0.053 (4)	0.059 (4)	0.079 (5)	0.008 (4)	-0.001 (4)	-0.008 (4)
O2	0.0285 (19)	0.077 (2)	0.0354 (15)	-0.0102 (19)	0.0145 (15)	-0.0088 (15)
C30	0.048 (3)	0.053 (4)	0.040 (2)	0.001 (3)	0.003 (2)	0.002 (3)
C31	0.090 (5)	0.079 (7)	0.038 (3)	0.026 (6)	0.021 (3)	-0.010 (4)
O2A	0.032 (3)	0.076 (3)	0.036 (2)	-0.012 (3)	0.017 (2)	-0.009 (2)
C30A	0.050 (4)	0.054 (4)	0.041 (3)	0.001 (3)	0.002 (3)	0.001 (3)

C31A	0.090 (5)	0.079 (7)	0.038 (3)	0.026 (6)	0.021 (3)	-0.010 (4)
B1	0.0216 (17)	0.0350 (19)	0.047 (2)	0.0031 (15)	0.0065 (15)	0.0024 (16)

*Geometric parameters (Å, °)*

Ru1—N3	2.086 (2)	C15—H15A	0.9300
Ru1—N1	2.088 (2)	C16—C21	1.388 (4)
Ru1—N5	2.144 (3)	C16—C17	1.390 (5)
Ru1—P1	2.3171 (8)	C17—C18	1.385 (5)
Ru1—S1	2.4540 (8)	C17—H17A	0.9300
Ru1—S2	2.4635 (8)	C18—C19	1.372 (5)
S1—P2	1.9643 (13)	C18—H18A	0.9300
S2—P2	1.9899 (12)	C19—C20	1.376 (5)
P1—C16	1.846 (3)	C19—H19A	0.9300
P1—C10	1.848 (3)	C20—C21	1.389 (5)
P1—C22	1.854 (3)	C20—H20A	0.9300
P2—O2	1.587 (3)	C21—H21A	0.9300
P2—O1A	1.591 (4)	C22—C27	1.386 (5)
P2—O2A	1.607 (4)	C22—C23	1.396 (5)
P2—O1	1.609 (3)	C23—C24	1.391 (5)
N1—C1	1.334 (4)	C23—H23A	0.9300
N1—N2	1.366 (3)	C24—C25	1.366 (6)
N2—C3	1.341 (4)	C24—H24A	0.9300
N2—B1	1.535 (5)	C25—C26	1.383 (6)
N3—C4	1.329 (4)	C25—H25A	0.9300
N3—N4	1.369 (3)	C26—C27	1.389 (5)
N4—C6	1.347 (4)	C26—H26A	0.9300
N4—B1	1.540 (4)	C27—H27A	0.9300
N5—C7	1.330 (4)	O1—C28	1.409 (5)
N5—N6	1.365 (4)	C28—C29	1.424 (6)
N6—C9	1.344 (4)	C28—H28A	0.9700
N6—B1	1.541 (5)	C28—H28B	0.9700
C1—C2	1.384 (5)	C29—H29A	0.9600
C1—H1A	0.9300	C29—H29B	0.9600
C2—C3	1.366 (5)	C29—H29C	0.9600
C2—H2A	0.9300	O1A—C28A	1.416 (6)
C3—H3A	0.9300	C29A—C28A	1.435 (6)
C4—C5	1.383 (4)	C29A—H29D	0.9600
C4—H4A	0.9300	C29A—H29E	0.9600
C5—C6	1.368 (5)	C29A—H29F	0.9600
C5—H5A	0.9300	C28A—H28C	0.9700
C6—H6A	0.9300	C28A—H28D	0.9700
C7—C8	1.388 (5)	O2—C30	1.426 (4)
C7—H7A	0.9300	C30—C31	1.446 (5)
C8—C9	1.366 (5)	C30—H30A	0.9700
C8—H8A	0.9300	C30—H30B	0.9700
C9—H9A	0.9300	C31—H31A	0.9600
C10—C11	1.396 (4)	C31—H31B	0.9600
C10—C15	1.399 (4)	C31—H31C	0.9600

## supplementary materials

---

C11—C12	1.390 (5)	O2A—C30A	1.420 (6)
C11—H11A	0.9300	C30A—C31A	1.438 (6)
C12—C13	1.380 (5)	C30A—H30C	0.9700
C12—H12A	0.9300	C30A—H30D	0.9700
C13—C14	1.381 (5)	C31A—H31D	0.9600
C13—H13A	0.9300	C31A—H31E	0.9600
C14—C15	1.383 (5)	C31A—H31F	0.9600
C14—H14A	0.9300	B1—H1	0.9800
N3—Ru1—N1	90.09 (9)	C14—C15—H15A	119.7
N3—Ru1—N5	83.57 (10)	C10—C15—H15A	119.7
N1—Ru1—N5	84.88 (10)	C21—C16—C17	117.8 (3)
N3—Ru1—P1	90.07 (7)	C21—C16—P1	123.5 (2)
N1—Ru1—P1	90.63 (7)	C17—C16—P1	118.6 (2)
N5—Ru1—P1	172.20 (7)	C18—C17—C16	121.5 (3)
N3—Ru1—S1	170.14 (7)	C18—C17—H17A	119.2
N1—Ru1—S1	93.07 (7)	C16—C17—H17A	119.2
N5—Ru1—S1	87.41 (7)	C19—C18—C17	119.9 (3)
P1—Ru1—S1	99.23 (3)	C19—C18—H18A	120.0
N3—Ru1—S2	94.61 (7)	C17—C18—H18A	120.0
N1—Ru1—S2	169.78 (7)	C18—C19—C20	119.6 (3)
N5—Ru1—S2	86.63 (7)	C18—C19—H19A	120.2
P1—Ru1—S2	98.42 (3)	C20—C19—H19A	120.2
S1—Ru1—S2	80.86 (3)	C19—C20—C21	120.6 (3)
P2—S1—Ru1	84.74 (4)	C19—C20—H20A	119.7
P2—S2—Ru1	83.95 (4)	C21—C20—H20A	119.7
C16—P1—C10	101.43 (14)	C16—C21—C20	120.6 (3)
C16—P1—C22	99.51 (14)	C16—C21—H21A	119.7
C10—P1—C22	101.07 (14)	C20—C21—H21A	119.7
C16—P1—Ru1	117.05 (10)	C27—C22—C23	118.1 (3)
C10—P1—Ru1	112.69 (10)	C27—C22—P1	119.6 (2)
C22—P1—Ru1	121.98 (10)	C23—C22—P1	122.3 (2)
O1A—P2—O2A	92.4 (6)	C24—C23—C22	120.6 (3)
O2—P2—O1	97.94 (17)	C24—C23—H23A	119.7
O2A—P2—O1	125.9 (4)	C22—C23—H23A	119.7
O2—P2—S1	118.31 (15)	C25—C24—C23	120.4 (4)
O1A—P2—S1	123.7 (4)	C25—C24—H24A	119.8
O2A—P2—S1	98.4 (4)	C23—C24—H24A	119.8
O1—P2—S1	105.05 (13)	C24—C25—C26	119.9 (3)
O2—P2—S2	115.21 (13)	C24—C25—H25A	120.0
O1A—P2—S2	122.1 (4)	C26—C25—H25A	120.0
O2A—P2—S2	105.9 (4)	C25—C26—C27	119.9 (4)
O1—P2—S2	112.03 (14)	C25—C26—H26A	120.0
S1—P2—S2	107.51 (5)	C27—C26—H26A	120.0
C1—N1—N2	106.2 (2)	C22—C27—C26	121.0 (3)
C1—N1—Ru1	134.7 (2)	C22—C27—H27A	119.5
N2—N1—Ru1	119.14 (19)	C26—C27—H27A	119.5
C3—N2—N1	109.4 (3)	C28—O1—P2	122.3 (3)
C3—N2—B1	130.7 (3)	O1—C28—C29	115.8 (6)
N1—N2—B1	119.5 (2)	O1—C28—H28A	108.3

C4—N3—N4	106.5 (2)	C29—C28—H28A	108.3
C4—N3—Ru1	135.1 (2)	O1—C28—H28B	108.3
N4—N3—Ru1	118.39 (18)	C29—C28—H28B	108.3
C6—N4—N3	109.0 (2)	H28A—C28—H28B	107.4
C6—N4—B1	130.5 (3)	C28—C29—H29A	109.5
N3—N4—B1	120.1 (2)	C28—C29—H29B	109.5
C7—N5—N6	106.2 (3)	H29A—C29—H29B	109.5
C7—N5—Ru1	135.9 (2)	C28—C29—H29C	109.5
N6—N5—Ru1	117.86 (19)	H29A—C29—H29C	109.5
C9—N6—N5	109.5 (3)	H29B—C29—H29C	109.5
C9—N6—B1	130.8 (3)	C28A—O1A—P2	119.2 (11)
N5—N6—B1	119.6 (2)	C28A—C29A—H29D	109.5
N1—C1—C2	110.6 (3)	C28A—C29A—H29E	109.5
N1—C1—H1A	124.7	H29D—C29A—H29E	109.5
C2—C1—H1A	124.7	C28A—C29A—H29F	109.5
C3—C2—C1	105.1 (3)	H29D—C29A—H29F	109.5
C3—C2—H2A	127.5	H29E—C29A—H29F	109.5
C1—C2—H2A	127.5	O1A—C28A—C29A	130 (2)
N2—C3—C2	108.8 (3)	O1A—C28A—H28C	104.7
N2—C3—H3A	125.6	C29A—C28A—H28C	104.7
C2—C3—H3A	125.6	O1A—C28A—H28D	104.7
N3—C4—C5	110.7 (3)	C29A—C28A—H28D	104.7
N3—C4—H4A	124.6	H28C—C28A—H28D	105.7
C5—C4—H4A	124.6	C30—O2—P2	120.1 (3)
C6—C5—C4	105.1 (3)	O2—C30—C31	109.8 (10)
C6—C5—H5A	127.5	O2—C30—H30A	109.7
C4—C5—H5A	127.5	C31—C30—H30A	109.7
N4—C6—C5	108.7 (3)	O2—C30—H30B	109.7
N4—C6—H6A	125.6	C31—C30—H30B	109.7
C5—C6—H6A	125.6	H30A—C30—H30B	108.2
N5—C7—C8	110.6 (3)	C30—C31—H31A	109.5
N5—C7—H7A	124.7	C30—C31—H31B	109.5
C8—C7—H7A	124.7	H31A—C31—H31B	109.5
C9—C8—C7	105.1 (3)	C30—C31—H31C	109.5
C9—C8—H8A	127.4	H31A—C31—H31C	109.5
C7—C8—H8A	127.4	H31B—C31—H31C	109.5
N6—C9—C8	108.5 (3)	C30A—O2A—P2	128.8 (11)
N6—C9—H9A	125.7	O2A—C30A—C31A	121 (3)
C8—C9—H9A	125.7	O2A—C30A—H30C	107.0
C11—C10—C15	118.3 (3)	C31A—C30A—H30C	107.0
C11—C10—P1	122.4 (2)	O2A—C30A—H30D	107.0
C15—C10—P1	119.3 (2)	C31A—C30A—H30D	107.0
C12—C11—C10	120.5 (3)	H30C—C30A—H30D	106.7
C12—C11—H11A	119.8	C30A—C31A—H31D	109.5
C10—C11—H11A	119.8	C30A—C31A—H31E	109.5
C13—C12—C11	120.4 (3)	H31D—C31A—H31E	109.5
C13—C12—H12A	119.8	C30A—C31A—H31F	109.5
C11—C12—H12A	119.8	H31D—C31A—H31F	109.5
C12—C13—C14	119.6 (3)	H31E—C31A—H31F	109.5

## supplementary materials

---

C12—C13—H13A	120.2	N2—B1—N4	109.7 (3)
C14—C13—H13A	120.2	N2—B1—N6	107.9 (3)
C13—C14—C15	120.5 (3)	N4—B1—N6	107.6 (3)
C13—C14—H14A	119.8	N2—B1—H1	110.5
C15—C14—H14A	119.8	N4—B1—H1	110.5
C14—C15—C10	120.7 (3)	N6—B1—H1	110.5

Fig. 1

