

Ethyl 3,7-dichloroquinoline-8-carboxylate**Fengxia Zhu, Li-Tao An,* Min Xia and Jian-Feng Zhou**

Jiangsu Key Laboratory for the Chemistry of Low-dimensional Materials, Department of Chemistry, Huaiyin Teachers College, Huaian 223300, Jiangsu Province, People's Republic of China

Correspondence e-mail: annleet@126.com


Received 8 October 2008; accepted 28 October 2008

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.043; wR factor = 0.117; data-to-parameter ratio = 17.7.

The title compound, $\text{C}_{12}\text{H}_9\text{Cl}_2\text{NO}_2$, was prepared by the esterification of 3,7-dichloroquinoline-8-carboxylic acid with triethyl phosphite. The crystal structure is stabilized by aromatic π - π stacking between the benzene and the pyridine rings of neighbouring molecules [centroid-centroid distances = 3.716 (2) and 3.642 (2) \AA]. In addition, weak intermolecular $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds are present in the structure.

Related literature

For the use of 3,7-dichloroquinoline-8-carboxylic acid as a herbicide, see: Nuria *et al.* (1997); Pornprom *et al.* (2006); Sunohara & Matsumoto (2004); Tresch & Grossmann (2002). For the usual preparative route, see: Yang *et al.* (2002). For related complexes, see: An *et al.* (2008); Che *et al.* (2005); Guo (2008); Li *et al.* (2008); Turel *et al.* (2004); Zhang *et al.* (2007). For 3,7-dichloroquinoline-8-carboxylic acid derivatives, see: Liang *et al.* (2006);

**Experimental***Crystal data*

$\text{C}_{12}\text{H}_9\text{Cl}_2\text{NO}_2$
 $M_r = 270.10$
Tetragonal, $I\bar{4}_1/a$
 $a = 25.4806(3)\text{ \AA}$
 $c = 7.3497(2)\text{ \AA}$
 $V = 4771.87(15)\text{ \AA}^3$

$Z = 16$
Mo $K\alpha$ radiation
 $\mu = 0.53\text{ mm}^{-1}$
 $T = 296(2)\text{ K}$
 $0.10 \times 0.08 \times 0.06\text{ mm}$

Data collection

Bruker SMART APEX2 diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 1999)
 $T_{\min} = 0.950$, $T_{\max} = 0.969$

19332 measured reflections
2750 independent reflections
1625 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.045$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.117$
 $S = 1.05$
2750 reflections

155 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.20\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.25\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C11—H11A \cdots N1 ⁱ	0.97	2.46	3.299 (3)	145

Symmetry code: (i) $-y + \frac{5}{4}, x + \frac{1}{4}, -z + \frac{5}{4}$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXS97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2003).

This work was supported financially by Jiangsu Key Laboratory for the Chemistry of Low-dimensional Materials.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LX2075).

References

- An, L.-T., Zhou, J., Zhou, J.-F. & Xia, M. (2008). *Acta Cryst. E64*, m1170.
- Bruker (1999). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Che, G.-B., Liu, C.-B., Cui, Y.-C. & Li, C.-B. (2005). *Acta Cryst. E61*, m2207–m2208.
- Farrugia, L. J. (1997). *J. Appl. Cryst. 30*, 565.
- Guo, X.-H. (2008). *Acta Cryst. E64*, o1786.
- Li, Z., Wu, F., Gong, Y., Zhang, Y. & Bai, C. (2008). *Acta Cryst. E64*, m227.
- Liang, Y., He, H.-W., Wang, Y.-Z. & Chen, T. (2006). *Acta Cryst. E62*, o3652–o3654.
- Nuria, L. M., George, M. & Rafael, D. P. (1997). *Pestic. Sci. 51*, 171–175.
- Pornprom, T., Mahatamuchoke, P. & Usui, K. (2006). *Pest Manag. Sci. 62*, 1109–1115.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst. 36*, 7–13.
- Sunohara, Y. & Matsumoto, H. (2004). *Plant Sci. 167*, 597–606.
- Tresch, S. & Grossmann, K. (2002). *Pestic. Biochem. Physiol. 75*, 73–78.
- Turel, I., Milena, P., Amalija, G., Enzo, A., Barbara, S., Alberta, B. & Gianni, S. (2004). *Inorg. Chim. Acta, 98*, 239–401.
- Yang, L., Wang, P. & Zhou, D. R. (2002). *J. Harbin Inst. Technol. 9*, 401–404.
- Zhang, Y.-H., Wu, F.-J., Li, X.-M., Zhu, M.-C. & Gong, Y. (2007). *Acta Cryst. E63*, m1557.

supporting information

Acta Cryst. (2008). E64, o2241 [doi:10.1107/S1600536808034995]

Ethyl 3,7-dichloroquinoline-8-carboxylate

Fengxia Zhu, Li-Tao An, Min Xia and Jian-Feng Zhou

S1. Comment

Quinclorac (3,7-dichloroquinoline-8-carboxylic acid) is one of the most effective herbicides (Nuria *et al.*, 1997; Pornprom *et al.*, 2006; Sunohara & Matsumoto, 2004; Tresch & Grossmann, 2002). Usually, it was prepared *via* Skraup cyclization from 2-methyl-3-chloroaniline, followed by chlorination and oxidation (Yang *et al.*, 2002). Furthermore, quinolinecarboxylates can chelate to metal atoms, forming the complexes, such as *trans*-Dimethanolbis(quinoline-8-carboxylato- κ^2N,O)- cobalt(II) (Che *et al.*, 2005), *catena*-Poly[nickel(II)-bis(μ -3,7-dichloroquinoline-8- χ arboxylato- $\kappa^3N,O:O'$)] (Zhang *et al.*, 2007), *catena*-Poly[cobalt(II)-bis (l-3,7-dichloroquinoline-8-carboxylato- $\kappa^3N,O:O'$)] (Li *et al.*, 2008). More recently, we also have reported a Zinc-quinclorac complex (An *et al.*, 2008) and quinclorac (Guo, 2008). But the derivatives of 3,7-dichloroquinoline-8-carboxylic acid have been less reported (Liang *et al.*, 2006). Here we report the crystal structure of the title compound, ethyl 3,7-dichloroquinoline-8-carboxylate (I) (Fig. 1).


In the title compound (I), as shown in Fig. 1, the plane (O1—C10—O2—C11) is nearly vertical to the quinoline ring, in which the dihedral angel is 86.6 (1). The quinoline unit is essentially planar, with a mean deviation of 0.007 (2) Å from the least-squares plane defined by the ten constituent atoms. The molecular packing (Fig. 2) is stabilized by aromatic π — π stackings between the benzene and the pyridine rings of the adjacent molecules. The $Cg1 \cdots Cg2^{ii}$ and $Cg1 \cdots Cg2^{iii}$ distances are 3.716 (2) and 3.642 (2) Å (Fig. 2; $Cg1$ and $Cg2$ are the centroids of the C1/C2/C3/C4/C9/C8 benzene ring and the N1/C7/C6/C5/C9/C8 pyridine ring, respectively, symmetry code as in Fig. 2). The crystal structure is further stabilized by intermolecular C11—H11A···Nⁱ hydrogen bonds (Fig. 2 and Table 1; symmetry code as in Fig. 2).

S2. Experimental


Ethyl 3,7-dichloroquinoline-8-carboxylate was obtained from the reaction of 3,7-dichloroquinoline-8-carboxylic acid with triethyl phosphite in refluxing condition. After recrystallization from ethanol, then it was dissolved the mixture of acetone/petroleum ether (1:4, V/V). The suitable single-crystal for X-ray analysis was obtained by slow evaporation.

S3. Refinement

All H atoms were geometrically positioned and refined using a riding model, with C—H = 0.93 (aromatic), 0.97 (methylene) and 0.96 Å (methyl) H atoms, and with $U_{iso}(H) = 1.2U_{eq}(C)$ (aromatic, methylene) and $1.5U_{eq}(C)$ (methyl) H atoms.

**Figure 1**

The molecular structure of the title compound, showing displacement ellipsoids drawn at the 30% probability level.

**Figure 2**

$\pi-\pi$ stackings and C—H \cdots N interactions (dotted lines) in the title compound. Cg denotes ring centroid. [Symmetry code: (i) $-y+5/4$, $x+1/4$, $-z+5/4$; (ii) $-x+1$, $-y+1$, $-z+2$; (iii) $-x+1$, $-y+1$, $-z+1$.]

Ethyl 3,7-dichloroquinoline-8-carboxylate

Crystal data

$C_{12}H_9Cl_2NO_2$
 $M_r = 270.10$

Tetragonal, $I4_1/a$
Hall symbol: -I_4ad

$a = 25.4806 (3)$ Å
 $c = 7.3497 (2)$ Å
 $V = 4771.87 (15)$ Å³
 $Z = 16$
 $F(000) = 2208$
 $D_x = 1.504$ Mg m⁻³
 Melting point: not measured K

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 2208 reflections
 $\theta = 1.6\text{--}26.0^\circ$
 $\mu = 0.53$ mm⁻¹
 $T = 296$ K
 Needle, colorless
 $0.10 \times 0.08 \times 0.06$ mm

Data collection

Bruker SMART APEX2
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 Detector resolution: 10.0 pixels mm⁻¹
 φ and ω scans
 Absorption correction: multi-scan
 (*SADABS*; Bruker, 1999)
 $T_{\min} = 0.950$, $T_{\max} = 0.969$

19332 measured reflections
 2750 independent reflections
 1625 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.045$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 1.6^\circ$
 $h = -32 \rightarrow 33$
 $k = -33 \rightarrow 32$
 $l = -9 \rightarrow 9$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.117$
 $S = 1.06$
 2750 reflections
 155 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0469P)^2 + 1.2301P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.20$ e Å⁻³
 $\Delta\rho_{\min} = -0.25$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.65771 (2)	0.53635 (3)	0.60570 (9)	0.0725 (2)
Cl2	0.34654 (3)	0.47136 (3)	0.89519 (10)	0.0809 (3)
O1	0.57642 (7)	0.63042 (6)	0.8389 (2)	0.0714 (5)
O2	0.55721 (6)	0.62556 (5)	0.5418 (2)	0.0560 (4)
N1	0.46789 (7)	0.56227 (7)	0.7898 (3)	0.0535 (5)
C1	0.55677 (8)	0.54727 (8)	0.7053 (3)	0.0479 (5)
C2	0.59627 (8)	0.51296 (8)	0.6642 (3)	0.0511 (5)
C3	0.58827 (9)	0.45826 (9)	0.6676 (3)	0.0582 (6)
H3	0.6157	0.4356	0.6390	0.070*

C4	0.54076 (9)	0.43888 (9)	0.7125 (3)	0.0594 (6)
H4	0.5358	0.4027	0.7144	0.071*
C5	0.44775 (9)	0.45445 (8)	0.8039 (3)	0.0574 (6)
H5	0.4405	0.4187	0.8107	0.069*
C6	0.40972 (9)	0.49023 (9)	0.8394 (3)	0.0555 (6)
C7	0.42156 (9)	0.54368 (9)	0.8316 (3)	0.0577 (6)
H7	0.3949	0.5675	0.8577	0.069*
C8	0.50652 (8)	0.52712 (8)	0.7514 (3)	0.0458 (5)
C9	0.49847 (8)	0.47219 (8)	0.7565 (3)	0.0492 (5)
C10	0.56464 (8)	0.60545 (8)	0.7069 (3)	0.0505 (5)
C11	0.56478 (9)	0.68194 (8)	0.5254 (3)	0.0609 (6)
H11A	0.5999	0.6915	0.5639	0.073*
H11B	0.5398	0.7004	0.6015	0.073*
C12	0.55684 (13)	0.69615 (10)	0.3318 (4)	0.0967 (10)
H12A	0.5809	0.6766	0.2574	0.145*
H12B	0.5630	0.7330	0.3160	0.145*
H12C	0.5215	0.6880	0.2967	0.145*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0602 (4)	0.0814 (5)	0.0760 (5)	0.0005 (3)	0.0113 (3)	-0.0024 (3)
Cl2	0.0661 (4)	0.1000 (5)	0.0765 (5)	-0.0247 (4)	-0.0006 (3)	0.0114 (4)
O1	0.0916 (13)	0.0584 (10)	0.0642 (12)	-0.0132 (9)	-0.0061 (9)	-0.0109 (8)
O2	0.0650 (10)	0.0406 (8)	0.0624 (11)	-0.0065 (7)	-0.0037 (8)	0.0032 (7)
N1	0.0524 (11)	0.0480 (10)	0.0601 (12)	0.0009 (9)	-0.0024 (9)	0.0023 (8)
C1	0.0560 (13)	0.0447 (12)	0.0430 (13)	-0.0014 (10)	-0.0049 (10)	0.0001 (9)
C2	0.0580 (13)	0.0529 (13)	0.0424 (12)	0.0009 (10)	-0.0021 (10)	-0.0015 (10)
C3	0.0704 (16)	0.0530 (14)	0.0511 (14)	0.0128 (11)	-0.0018 (11)	-0.0042 (10)
C4	0.0804 (17)	0.0423 (12)	0.0555 (15)	0.0022 (12)	-0.0030 (12)	-0.0035 (10)
C5	0.0778 (17)	0.0462 (12)	0.0483 (14)	-0.0158 (12)	-0.0057 (11)	0.0036 (10)
C6	0.0584 (14)	0.0637 (15)	0.0445 (13)	-0.0129 (11)	-0.0065 (10)	0.0043 (10)
C7	0.0557 (14)	0.0595 (14)	0.0578 (15)	0.0034 (11)	-0.0035 (11)	0.0037 (11)
C8	0.0567 (13)	0.0418 (12)	0.0389 (12)	0.0007 (10)	-0.0072 (9)	0.0002 (9)
C9	0.0667 (15)	0.0412 (12)	0.0397 (13)	-0.0033 (10)	-0.0077 (10)	0.0014 (9)
C10	0.0456 (12)	0.0497 (13)	0.0562 (15)	-0.0038 (10)	0.0016 (10)	-0.0020 (11)
C11	0.0583 (14)	0.0392 (12)	0.0851 (18)	-0.0080 (10)	0.0071 (12)	0.0011 (11)
C12	0.139 (3)	0.0547 (16)	0.097 (2)	-0.0178 (17)	-0.0243 (19)	0.0215 (15)

Geometric parameters (\AA , ^\circ)

C11—C2	1.729 (2)	C4—H4	0.9300
Cl2—C6	1.730 (2)	C5—C6	1.356 (3)
O1—C10	1.198 (2)	C5—C9	1.413 (3)
O2—C10	1.331 (2)	C5—H5	0.9300
O2—C11	1.454 (2)	C6—C7	1.396 (3)
N1—C7	1.309 (3)	C7—H7	0.9300
N1—C8	1.360 (2)	C8—C9	1.415 (3)

C1—C2	1.367 (3)	C11—C12	1.482 (3)
C1—C8	1.420 (3)	C11—H11A	0.9700
C1—C10	1.496 (3)	C11—H11B	0.9700
C2—C3	1.409 (3)	C12—H12A	0.9600
C3—C4	1.348 (3)	C12—H12B	0.9600
C3—H3	0.9300	C12—H12C	0.9600
C4—C9	1.409 (3)		
C10—O2—C11	115.91 (17)	C6—C7—H7	118.1
C7—N1—C8	117.60 (18)	N1—C8—C9	122.72 (19)
C2—C1—C8	119.03 (18)	N1—C8—C1	117.63 (17)
C2—C1—C10	122.47 (18)	C9—C8—C1	119.65 (19)
C8—C1—C10	118.49 (18)	C4—C9—C5	124.3 (2)
C1—C2—C3	121.5 (2)	C4—C9—C8	118.6 (2)
C1—C2—Cl1	120.06 (16)	C5—C9—C8	117.1 (2)
C3—C2—Cl1	118.45 (17)	O1—C10—O2	124.7 (2)
C4—C3—C2	119.8 (2)	O1—C10—C1	124.5 (2)
C4—C3—H3	120.1	O2—C10—C1	110.81 (18)
C2—C3—H3	120.1	O2—C11—C12	107.63 (19)
C3—C4—C9	121.5 (2)	O2—C11—H11A	110.2
C3—C4—H4	119.3	C12—C11—H11A	110.2
C9—C4—H4	119.3	O2—C11—H11B	110.2
C6—C5—C9	119.07 (19)	C12—C11—H11B	110.2
C6—C5—H5	120.5	H11A—C11—H11B	108.5
C9—C5—H5	120.5	C11—C12—H12A	109.5
C5—C6—C7	119.6 (2)	C11—C12—H12B	109.5
C5—C6—Cl2	121.59 (18)	H12A—C12—H12B	109.5
C7—C6—Cl2	118.82 (19)	C11—C12—H12C	109.5
N1—C7—C6	123.9 (2)	H12A—C12—H12C	109.5
N1—C7—H7	118.1	H12B—C12—H12C	109.5

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C11—H11A···N1 ⁱ	0.97	2.46	3.299 (3)	145

Symmetry code: (i) $-y+5/4, x+1/4, -z+5/4$.