

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# Poly[[ $\mu_3$ - $N,N'$ -bis(3-pyridylmethyl)-thiourea- $\kappa^3 N:N':S$ ]iodidocopper(I)]

Shi-Shen Zhang

Department of Applied Chemistry, Zhejiang Sci-Tech University, Hang Zhou, 310018, People's Republic of China


Correspondence e-mail: zhangshishen@126.com

Received 22 September 2008; accepted 26 September 2008

 Key indicators: single-crystal X-ray study;  $T = 294$  K; mean  $\sigma(C-C) = 0.005$  Å;  $R$  factor = 0.026;  $wR$  factor = 0.062; data-to-parameter ratio = 15.2.

In the title coordination polymer,  $[CuI(C_{13}H_{14}N_4S)]_n$ , the Cu<sup>I</sup> atom is coordinated by two N atoms from two  $N,N'$ -bis(3-pyridylmethyl)thiourea ligands, as well as by the S atom of a third ligand and an I atom to confer a distorted tetrahedral coordination at the metal centre. The coordination bonds give rise to a layer structure parallel to (010).

## Related literature

 For related literature, see: Li *et al.* (2002); Zhang *et al.* (2006).


## Experimental

## Crystal data

$[CuI(C_{13}H_{14}N_4S)]$   
 $M_r = 448.78$   
 Monoclinic,  $P2_1/c$   
 $a = 13.3610$  (10) Å  
 $b = 8.3673$  (7) Å  
 $c = 14.2686$  (11) Å  
 $\beta = 102.001$  (2)°

$V = 1560.3$  (2) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 3.51$  mm<sup>-1</sup>  
 $T = 294$  (2) K  
 $0.20 \times 0.15 \times 0.12$  mm

## Data collection

Bruker SMART CCD diffractometer  
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.541$ ,  $T_{\max} = 0.678$

8191 measured reflections  
 2750 independent reflections  
 2418 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.020$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.026$   
 $wR(F^2) = 0.062$   
 $S = 1.06$   
 2750 reflections

181 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.73$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.28$  e Å<sup>-3</sup>

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1998); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by the Foundation for Young Researchers of the Key Laboratory of Advanced Textile Materials and Manufacturing Technology, Ministry of Education (grant No. 2007QN05).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2496).

## References

- Bruker (1998). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Li, G., Hou, H.-W., Niu, Y.-Y., Fan, Y.-T., Liu, Z.-S., Ge, T.-Z. & Xin, X.-Q. (2002). *Inorg. Chim. Acta*, **332**, 216–222.  
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Zhang, X.-J., Zhou, X.-P. & Li, D. (2006). *Cryst. Growth Des.* **6**, 1440–1444.

**supplementary materials**

*Acta Cryst.* (2008). E64, m1395 [ doi:10.1107/S1600536808031188 ]

## Poly[[ $\mu_3$ -*N,N'*-bis(3-pyridylmethyl)thiourea- $\kappa^3$ *N:N':S*]iodidocopper(I)]

S.-S. Zhang

### Comment

Flexible ligand has been considered as one of the most important type of organic ligand for their flexibility and conformational freedom allow for greater structural diversity. *N,N'*-bis(3-pyridylmethyl)thiourea, as one kind of those ligand, has usually been used to construct a great variety of structurally interesting entities. such as helix, macrocycle (Zhang *et al.*, 2006; Li *et al.*, 2002).

The asymmetric unit of the title compound (I) is illustrated in Fig. 1. Single-crystal X-ray diffraction shows that the asymmetric unit contains one Cu crystallographically nonequivalent atom. The Cu(I) atom coordinated by two N atoms from two *N,N'*-bis(3-pyridylmethyl)thiourea ligands as well as by the S atom of a third ligand to confer a tetrahedral geometry at the metal center. The Cu atom coordination by two N atoms to form a one-dimensional helix, and is then linked by the bond of Cu atom and S atom to extend to a two-dimensional structure. The crystal packing is stabilized by intermolecular  $\pi$ - $\pi$  stacking interaction (Fig. 2).

### Experimental

a mixture of CuI (0.038 g, 0.2 nmol) and *N,N'*-bis(3-pyridylmethyl)thiourea (0.026 g, 0.1 nmol) in mole ratio of 2:1 in acetonitrile (6 cm<sup>3</sup>) was sealed in 15 cm<sup>3</sup> Teflon-lined reactor and heated to 110°C for 10 h and then cooled to room temperature at a rate of 5°C/h. the yellow block crystal was obtained in the yield of 35%.

The web of checkcif show one Alert level B (Hirshfeld Test Diff (M—X) II – Cu1.. 43.03 su), we think this is the result of the slightly distorted I atom for his unidentate coordination model.

### Refinement

H atoms were positioned geometrically and refined using a riding model, with C—H = 0.93 Å (aromatic) or 0.97 Å (aliphatic) and N—H = 0.86 Å, and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$

### Figures




Fig. 1. The asymmetric unit of the title compound showing 30° probability ellipsoids.




Fig. 2. The crystal packing of the title compound.

## Poly[[ $\mu_3$ -*N,N'*-bis(3-pyridylmethyl)thiourea- $\kappa^3$ *N:N':S*]iodidocopper(I)]

### Crystal data

[CuI(C <sub>13</sub> H <sub>14</sub> N <sub>4</sub> S)]	$F_{000} = 872$
$M_r = 448.78$	$D_x = 1.910 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 13.3610 (10) \text{ \AA}$	Cell parameters from 3234 reflections
$b = 8.3673 (7) \text{ \AA}$	$\theta = 2.8\text{--}27.7^\circ$
$c = 14.2686 (11) \text{ \AA}$	$\mu = 3.51 \text{ mm}^{-1}$
$\beta = 102.001 (2)^\circ$	$T = 294 (2) \text{ K}$
$V = 1560.3 (2) \text{ \AA}^3$	Block, yellow
$Z = 4$	$0.20 \times 0.15 \times 0.12 \text{ mm}$

### Data collection

Bruker SMART CCD diffractometer	2750 independent reflections
Radiation source: fine-focus sealed tube	2418 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.020$
$T = 294(2) \text{ K}$	$\theta_{\text{max}} = 25.0^\circ$
$\varphi$ and $\omega$ scans	$\theta_{\text{min}} = 2.8^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -12 \rightarrow 15$
$T_{\text{min}} = 0.541$ , $T_{\text{max}} = 0.678$	$k = -9 \rightarrow 9$
8191 measured reflections	$l = -15 \rightarrow 16$

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.026$	H-atom parameters constrained
$wR(F^2) = 0.062$	$w = 1/[\sigma^2(F_o^2) + (0.0275P)^2 + 0.8144P]$
$S = 1.06$	where $P = (F_o^2 + 2F_c^2)/3$
2750 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
181 parameters	$\Delta\rho_{\text{max}} = 0.73 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.28 \text{ e \AA}^{-3}$
	Extinction correction: none

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.4137 (2)	0.8305 (4)	0.3317 (2)	0.0340 (7)
H1A	0.4469	0.7269	0.3368	0.041*
H1B	0.4581	0.9036	0.3739	0.041*
C2	0.4006 (2)	0.8893 (3)	0.2301 (2)	0.0290 (6)
C3	0.3491 (2)	1.0308 (3)	0.2012 (2)	0.0315 (7)
H3	0.3220	1.0879	0.2460	0.038*
C4	0.3768 (2)	1.0054 (4)	0.0489 (2)	0.0361 (7)
H4	0.3685	1.0441	-0.0134	0.043*
C5	0.4292 (2)	0.8663 (4)	0.0718 (2)	0.0402 (7)
H5	0.4565	0.8122	0.0260	0.048*
C6	0.4415 (2)	0.8065 (4)	0.1630 (2)	0.0376 (7)
H6	0.4770	0.7115	0.1795	0.045*
C7	-0.0360 (2)	0.7596 (4)	0.4500 (2)	0.0379 (7)
H7	-0.0490	0.7921	0.3863	0.045*
C8	0.0621 (2)	0.7151 (4)	0.4916 (2)	0.0372 (7)
C9	0.0813 (3)	0.6670 (5)	0.5858 (3)	0.0521 (9)
H9	0.1469	0.6365	0.6166	0.063*
C10	0.0020 (3)	0.6648 (5)	0.6337 (3)	0.0571 (10)
H10	0.0130	0.6307	0.6970	0.068*
C11	-0.0934 (3)	0.7134 (4)	0.5874 (2)	0.0445 (8)
H11	-0.1460	0.7147	0.6210	0.053*
C12	0.1456 (2)	0.7225 (4)	0.4347 (3)	0.0413 (8)
H12A	0.1909	0.6314	0.4502	0.050*
H12B	0.1157	0.7194	0.3667	0.050*
C13	0.2875 (2)	0.9112 (3)	0.4274 (2)	0.0309 (6)
Cu1	0.26574 (3)	1.19634 (5)	0.57370 (3)	0.04022 (12)
N1	0.20322 (18)	0.8710 (3)	0.45837 (19)	0.0397 (6)
H1	0.1810	0.9377	0.4952	0.048*
N2	0.31731 (18)	0.8164 (3)	0.36364 (17)	0.0317 (6)
H2	0.2765	0.7407	0.3393	0.038*
N3	0.33636 (18)	1.0893 (3)	0.11266 (18)	0.0328 (6)
N4	-0.11373 (19)	0.7591 (3)	0.4956 (2)	0.0396 (6)
S3	0.35264 (6)	1.07993 (10)	0.46956 (6)	0.0397 (2)

## supplementary materials

II                    0.24731 (2)                    1.03780 (3)                    0.728706 (18)                    0.05208 (10)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0309 (16)	0.0383 (16)	0.0344 (17)	0.0077 (12)	0.0101 (13)	-0.0001 (13)
C2	0.0242 (14)	0.0350 (16)	0.0285 (16)	-0.0015 (12)	0.0072 (12)	-0.0040 (13)
C3	0.0323 (16)	0.0333 (16)	0.0322 (17)	-0.0006 (12)	0.0140 (13)	-0.0037 (13)
C4	0.0379 (17)	0.0451 (17)	0.0268 (16)	-0.0068 (14)	0.0102 (14)	-0.0002 (14)
C5	0.0462 (19)	0.0439 (18)	0.0348 (18)	0.0034 (15)	0.0184 (15)	-0.0077 (15)
C6	0.0418 (18)	0.0334 (16)	0.0396 (19)	0.0072 (13)	0.0134 (14)	-0.0025 (14)
C7	0.0335 (17)	0.0461 (18)	0.0370 (18)	-0.0018 (14)	0.0139 (14)	-0.0007 (15)
C8	0.0338 (17)	0.0328 (16)	0.047 (2)	-0.0057 (13)	0.0128 (15)	-0.0044 (15)
C9	0.0371 (19)	0.065 (2)	0.052 (2)	0.0022 (16)	0.0035 (16)	0.0058 (19)
C10	0.055 (2)	0.076 (3)	0.040 (2)	-0.0008 (19)	0.0097 (17)	0.0101 (19)
C11	0.0411 (19)	0.056 (2)	0.040 (2)	-0.0058 (16)	0.0171 (15)	0.0028 (17)
C12	0.0329 (17)	0.0378 (17)	0.056 (2)	-0.0043 (14)	0.0151 (15)	-0.0110 (16)
C13	0.0311 (16)	0.0309 (15)	0.0312 (16)	0.0012 (12)	0.0079 (13)	-0.0013 (13)
Cu1	0.0403 (2)	0.0416 (2)	0.0416 (2)	-0.00434 (17)	0.01496 (18)	-0.00960 (18)
N1	0.0378 (14)	0.0374 (14)	0.0495 (17)	-0.0068 (12)	0.0223 (13)	-0.0140 (13)
N2	0.0346 (14)	0.0321 (13)	0.0297 (13)	-0.0022 (10)	0.0097 (11)	-0.0049 (11)
N3	0.0324 (13)	0.0361 (13)	0.0311 (14)	-0.0004 (11)	0.0099 (11)	0.0023 (11)
N4	0.0324 (14)	0.0475 (16)	0.0412 (16)	-0.0010 (12)	0.0129 (12)	0.0011 (13)
S3	0.0369 (4)	0.0383 (4)	0.0488 (5)	-0.0085 (3)	0.0203 (4)	-0.0128 (4)
II	0.07354 (19)	0.04215 (15)	0.04564 (16)	0.00823 (11)	0.02411 (12)	0.01044 (10)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

C1—N2	1.457 (4)	C9—H9	0.9300
C1—C2	1.506 (4)	C10—C11	1.370 (5)
C1—H1A	0.9700	C10—H10	0.9300
C1—H1B	0.9700	C11—N4	1.337 (4)
C2—C6	1.384 (4)	C11—H11	0.9300
C2—C3	1.388 (4)	C12—N1	1.464 (4)
C3—N3	1.332 (4)	C12—H12A	0.9700
C3—H3	0.9300	C12—H12B	0.9700
C4—N3	1.348 (4)	C13—N2	1.330 (4)
C4—C5	1.363 (4)	C13—N1	1.335 (4)
C4—H4	0.9300	C13—S3	1.701 (3)
C5—C6	1.371 (5)	Cu1—N3 <sup>i</sup>	2.049 (2)
C5—H5	0.9300	Cu1—N4 <sup>ii</sup>	2.099 (3)
C6—H6	0.9300	Cu1—S3	2.2852 (8)
C7—N4	1.335 (4)	Cu1—I1	2.6341 (5)
C7—C8	1.372 (4)	N1—H1	0.8600
C7—H7	0.9300	N2—H2	0.8600
C8—C9	1.374 (5)	N3—Cu1 <sup>iii</sup>	2.049 (2)
C8—C12	1.512 (4)	N4—Cu1 <sup>ii</sup>	2.099 (3)
C9—C10	1.376 (5)		

N2—C1—C2	113.3 (2)	C9—C10—H10	120.3
N2—C1—H1A	108.9	N4—C11—C10	122.4 (3)
C2—C1—H1A	108.9	N4—C11—H11	118.8
N2—C1—H1B	108.9	C10—C11—H11	118.8
C2—C1—H1B	108.9	N1—C12—C8	108.8 (2)
H1A—C1—H1B	107.7	N1—C12—H12A	109.9
C6—C2—C3	117.6 (3)	C8—C12—H12A	109.9
C6—C2—C1	121.3 (3)	N1—C12—H12B	109.9
C3—C2—C1	121.1 (2)	C8—C12—H12B	109.9
N3—C3—C2	123.6 (3)	H12A—C12—H12B	108.3
N3—C3—H3	118.2	N2—C13—N1	118.0 (3)
C2—C3—H3	118.2	N2—C13—S3	122.2 (2)
N3—C4—C5	122.7 (3)	N1—C13—S3	119.8 (2)
N3—C4—H4	118.6	N3 <sup>i</sup> —Cu1—N4 <sup>ii</sup>	108.54 (10)
C5—C4—H4	118.6	N3 <sup>i</sup> —Cu1—S3	106.38 (7)
C4—C5—C6	119.6 (3)	N4 <sup>ii</sup> —Cu1—S3	110.01 (8)
C4—C5—H5	120.2	N3 <sup>i</sup> —Cu1—I1	109.34 (7)
C6—C5—H5	120.2	N4 <sup>ii</sup> —Cu1—I1	103.57 (7)
C5—C6—C2	119.1 (3)	S3—Cu1—I1	118.69 (3)
C5—C6—H6	120.4	C13—N1—C12	125.3 (2)
C2—C6—H6	120.4	C13—N1—H1	117.4
N4—C7—C8	123.9 (3)	C12—N1—H1	117.4
N4—C7—H7	118.0	C13—N2—C1	125.2 (3)
C8—C7—H7	118.0	C13—N2—H2	117.4
C7—C8—C9	118.0 (3)	C1—N2—H2	117.4
C7—C8—C12	120.1 (3)	C3—N3—C4	117.3 (3)
C9—C8—C12	121.9 (3)	C3—N3—Cu1 <sup>iii</sup>	122.7 (2)
C8—C9—C10	118.8 (3)	C4—N3—Cu1 <sup>iii</sup>	119.9 (2)
C8—C9—H9	120.6	C7—N4—C11	117.2 (3)
C10—C9—H9	120.6	C7—N4—Cu1 <sup>ii</sup>	123.0 (2)
C11—C10—C9	119.5 (3)	C11—N4—Cu1 <sup>ii</sup>	119.4 (2)
C11—C10—H10	120.3	C13—S3—Cu1	106.95 (10)

Symmetry codes: (i)  $x, -y+5/2, z+1/2$ ; (ii)  $-x, -y+2, -z+1$ ; (iii)  $x, -y+5/2, z-1/2$ .

Fig. 1




Fig. 2

