

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

catena-Poly[[*(1,12,15,26-tetraaza-5,8,19,22-tetraoxa-3,4:9,10:17,18:-23,24-tetrabenzocyclooctacosane-κ⁴N¹,N¹²,N¹⁵,N²⁶)nickel(II)-μ-terephthalato-κ²O¹:O⁴*]]

Shuang-Ming Meng, Yue-Qin Fan and Yong Guo*

College of Chemistry and Chemical Engineering, Shanxi Datong University, Datong 037009, People's Republic of China
Correspondence e-mail: ybsysm@126.com


Received 17 September 2008; accepted 13 October 2008

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.063; wR factor = 0.155; data-to-parameter ratio = 17.8.

In the title compound, $[\text{Ni}(\text{C}_8\text{H}_4\text{O}_4)(\text{C}_{36}\text{H}_{44}\text{N}_4\text{O}_4)]_n$, the Ni^{II} atom is coordinated in a distorted octahedral geometry by the four N atoms of the 1,12,15,26-tetraaza-5,8,19,22-tetra-oxa-3,4:9,10:17,18:23,24-tetrabenzocyclooctacosane ligand and two O atoms from the terephthalate dianions. The Ni^{II} atoms, which lie on inversion centres, are linked *via* terephthalate ligands to form a chain structure along [101]. The structure is stabilized by three intramolecular and one intermolecular N—H···O hydrogen bonds.

Related literature

For general background, see: Choi & Suh (1999); Massoud *et al.* (2006); Ray *et al.* (2006). For a related structure, see: Jiang *et al.* (2005).



Experimental

Crystal data

$[\text{Ni}(\text{C}_8\text{H}_4\text{O}_4)(\text{C}_{36}\text{H}_{44}\text{N}_4\text{O}_4)]$
 $M_r = 819.57$
Monoclinic, $P2_1/n$
 $a = 11.407$ (3) Å
 $b = 16.575$ (3) Å
 $c = 21.675$ (6) Å
 $\beta = 101.758$ (10)°

$V = 4012.2$ (17) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.54$ mm⁻¹
 $T = 293$ (2) K
 $0.35 \times 0.28 \times 0.21$ mm

Data collection

Rigaku R-Axis RAPID diffractometer
Absorption correction: multi-scan (ABSCOR; Higashi, 1995)
 $T_{\text{min}} = 0.839$, $T_{\text{max}} = 0.910$

37289 measured reflections
9133 independent reflections
5535 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.095$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.063$
 $wR(F^2) = 0.155$
 $S = 1.05$
9133 reflections
514 parameters
4 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.92$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.79$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1N}\cdots\text{O1}$	0.83 (2)	2.58 (4)	3.101 (4)	122 (3)
$\text{N2}-\text{H2N}\cdots\text{O8}^i$	0.81 (2)	2.26 (2)	3.003 (4)	152 (4)
$\text{N3}-\text{H3N}\cdots\text{O3}$	0.84 (2)	2.47 (3)	3.044 (4)	127 (3)
$\text{N4}-\text{H4N}\cdots\text{O5}$	0.83 (2)	2.14 (3)	2.862 (4)	144 (4)

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $x - \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (iii) $x + 1, y, z$.

Data collection: *PROCESS-AUTO* (Rigaku, 1998); cell refinement: *PROCESS-AUTO*; data reduction: *PROCESS-AUTO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL-Plus* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

We thank the Analysis and Testing Foundation of Northeast Normal University for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2108).

References

- Choi, H. J. & Suh, M. P. (1999). *Inorg. Chem.* **38**, 6309–6312.
Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
Jiang, L., Feng, X.-F. & Lu, T.-B. (2005). *Cryst. Growth Des.* **5**, 1469–1475.
Massoud, S. S., Mautner, F. A., Vicente, R. & Rodrigue, B. M. (2006). *Inorg. Chem. Acta*, **359**, 3321–3329.
Ray, A., Mijanuddin, M., Chatterjee, R., Marek, J., Mondal, S. & Ali, M. (2006). *Inorg. Chem. Commun.* **9**, 167–170.
Rigaku (1998). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, m1431 [doi:10.1107/S160053680803314X]

***catena*-Poly[[*(1,12,15,26-tetraaza-5,8,19,22-tetraoxa-3,4:9,10:17,18:23,24-tetrabenzocyclooctacosane-κ⁴N¹,N¹²,N¹⁵,N²⁶)nickel(II)*]-*μ*-terephthalato-κ²O¹:O⁴]**

S.-M. Meng, Y.-Q. Fan and Y. Guo

Comment

In recent years, intense research activity has been directed toward the assembly of carboxylato-bridged macrocyclic polymers due to their intriguing multidimensional networks (Choi & Suh, 1999; Massoud *et al.* 2006; Ray *et al.* 2006). As an extension of the research on the macrocyclic complexes, we have prepared the title compound, (I). In this paper the crystal structure of (I) is reported.

In the title compound, the Ni^{II} atom, which lies on an inversion centre, displays a distorted octahedral coordination geometry provided by four nitrogen atoms from the ligand, 3,4:9,10:17,18:23,24-tetrabenzocyclooctacosane (*L*) and two oxygen atoms from two distinct terephthalate (tp) dianion ligands (Fig. 1). The bond distances and angles show normal values (Jiang *et al.* 2005). The Ni^{II} atoms are linked *via* tp ligands to form a one-dimensional chain structure (Fig. 2). The constituent of the title compound are linked through hydrogen bonds to form a complicated three-dimensional network (Table 1); the N and C atoms play a role as donors, while carboxylate-O atoms function as acceptors in these hydrogen bonds.

Experimental

A mixture of NiCO₃ (0.119 mg, 1 mmol), terephthalic acid (0.162 mg, 1 mmol) and 3,4:9,10:17,18:23,24-tetrabenzocyclooctacosane (0.596 mg, 1 mmol) in EtOH (10 ml) was placed in a Teflon reactor and heated at 393 K for 3 days, and then it was gradually cooled to room temperature at a rate of 10 K.h⁻¹. Green crystals were obtained.

Refinement

All H atoms bound to C atoms were positioned geometrically and refined as riding, with C—H = 0.93 Å (CH) and 0.97 Å (CH₂) and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The H atoms bound to N atoms were located in a difference Fourier map and refined with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$.

Figures




Fig. 1. Molecular structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level and H atoms have been omitted for clarity. [Symmetry code: (i) $x + 1/2, 1/2 - y, 1/2 + z$]




Fig. 2. View of the one-dimensional polymeric chain of the title compound in the unit cell; H atoms have been omitted for clarity.

catena-Poly[[([1,12,15,26-tetraaza-5,8,19,22-tetraoxa-3,4:9,10:17,18:23,24-tetrabenzocyclooctacosane- $\kappa^4 N^1, N^{12}, N^{15}, N^{26}$)nickel(II)]- μ -terephthalato- $\kappa^2 O^1:O^4$]

Crystal data

[Ni(C₈H₄O₄)(C₃₆H₄₄N₄O₄)]

$M_r = 819.57$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 11.407 (3) \text{ \AA}$

$b = 16.575 (3) \text{ \AA}$

$c = 21.675 (6) \text{ \AA}$

$\beta = 101.758 (10)^\circ$

$V = 4012.2 (17) \text{ \AA}^3$

$Z = 4$

$F_{000} = 1728$

$D_x = 1.357 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 9133 reflections

$\theta = 3.1\text{--}27.5^\circ$

$\mu = 0.54 \text{ mm}^{-1}$

$T = 293 (2) \text{ K}$

Block, green

$0.35 \times 0.28 \times 0.21 \text{ mm}$

Data collection

Rigaku R-Axis RAPID
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 10.0 pixels mm^{-1}

$T = 293(2) \text{ K}$

ω scans

Absorption correction: multi-scan
(ABSCOR; Higashi, 1995)

$T_{\min} = 0.839, T_{\max} = 0.910$

37289 measured reflections

9133 independent reflections

5535 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.095$

$\theta_{\text{max}} = 27.5^\circ$

$\theta_{\text{min}} = 3.1^\circ$

$h = -14 \rightarrow 14$

$k = -21 \rightarrow 21$

$l = -27 \rightarrow 28$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.063$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.155$	$w = 1/[\sigma^2(F_o^2) + (0.0632P)^2 + 2.1456P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
9133 reflections	$(\Delta/\sigma)_{\max} = 0.001$
514 parameters	$\Delta\rho_{\max} = 0.92 \text{ e } \text{\AA}^{-3}$
4 restraints	$\Delta\rho_{\min} = -0.79 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.76937 (4)	0.21301 (2)	0.898586 (19)	0.02536 (13)
C1	0.5251 (3)	0.0013 (2)	0.84876 (18)	0.0422 (9)
C2	0.5463 (4)	-0.0627 (3)	0.8114 (2)	0.0660 (14)
H2	0.6008	-0.0565	0.7852	0.079*
C3	0.4876 (5)	-0.1363 (3)	0.8124 (3)	0.0747 (16)
H3	0.5032	-0.1786	0.7871	0.090*
C4	0.4075 (4)	-0.1459 (3)	0.8502 (2)	0.0628 (13)
H4	0.3674	-0.1947	0.8505	0.075*
C5	0.3859 (4)	-0.0837 (3)	0.8880 (2)	0.0573 (12)
H5	0.3307	-0.0902	0.9138	0.069*
C6	0.4457 (4)	-0.0118 (2)	0.8878 (2)	0.0479 (10)
C9	0.3299 (4)	0.0644 (3)	0.9479 (2)	0.0519 (11)
H9A	0.3091	0.0169	0.9696	0.062*
H9B	0.2668	0.0734	0.9112	0.062*
C10	0.3383 (4)	0.1346 (3)	0.9901 (2)	0.0623 (13)
H10A	0.2583	0.1526	0.9920	0.075*

supplementary materials

H10B	0.3779	0.1185	1.0322	0.075*
C11	0.4113 (3)	0.2674 (2)	1.00719 (17)	0.0414 (9)
C12	0.3125 (4)	0.3033 (3)	1.0236 (2)	0.0643 (13)
H12	0.2367	0.2809	1.0106	0.077*
C13	0.3278 (5)	0.3727 (4)	1.0595 (3)	0.0872 (18)
H13	0.2621	0.3977	1.0706	0.105*
C14	0.4397 (6)	0.4046 (4)	1.0786 (3)	0.0852 (18)
H14	0.4500	0.4506	1.1037	0.102*
C15	0.5377 (5)	0.3695 (3)	1.0612 (2)	0.0641 (13)
H15	0.6131	0.3923	1.0742	0.077*
C16	0.5245 (3)	0.3001 (2)	1.02442 (17)	0.0395 (9)
C17	0.6303 (3)	0.2610 (2)	1.00410 (16)	0.0389 (9)
H17A	0.7025	0.2758	1.0340	0.047*
H17B	0.6216	0.2029	1.0062	0.047*
C18	0.6834 (4)	0.3678 (2)	0.93574 (19)	0.0457 (10)
H18A	0.6499	0.4007	0.9649	0.055*
H18B	0.6541	0.3882	0.8935	0.055*
C19	0.8183 (4)	0.3724 (2)	0.95182 (19)	0.0476 (10)
H19A	0.8440	0.4281	0.9509	0.057*
H19B	0.8479	0.3512	0.9938	0.057*
C20	0.8686 (4)	0.3707 (2)	0.84692 (18)	0.0435 (9)
H20A	0.7875	0.3871	0.8282	0.052*
H20B	0.8962	0.3354	0.8171	0.052*
C21	0.9472 (4)	0.4448 (2)	0.85675 (18)	0.0429 (9)
C22	0.8995 (5)	0.5222 (2)	0.8504 (2)	0.0600 (12)
H22	0.8169	0.5296	0.8406	0.072*
C23	0.9760 (6)	0.5892 (3)	0.8586 (3)	0.0780 (17)
H23	0.9442	0.6410	0.8549	0.094*
C24	1.0953 (6)	0.5787 (3)	0.8720 (3)	0.0761 (16)
H24	1.1452	0.6236	0.8768	0.091*
C25	1.1447 (5)	0.5027 (3)	0.8786 (2)	0.0646 (13)
H25	1.2274	0.4961	0.8883	0.077*
C26	1.0705 (4)	0.4364 (2)	0.8707 (2)	0.0487 (10)
C27	1.2162 (4)	0.3402 (3)	0.8544 (3)	0.0679 (14)
H27A	1.2857	0.3572	0.8853	0.081*
H27B	1.2160	0.3694	0.8156	0.081*
C28	1.2233 (4)	0.2533 (3)	0.8431 (3)	0.0801 (17)
H28A	1.2944	0.2420	0.8265	0.096*
H28B	1.2305	0.2247	0.8827	0.096*
C29	1.1211 (4)	0.1467 (2)	0.78328 (19)	0.0444 (9)
C30	1.2200 (4)	0.1073 (3)	0.7688 (2)	0.0617 (13)
H30	1.2932	0.1336	0.7733	0.074*
C31	1.2076 (5)	0.0288 (3)	0.7478 (2)	0.0674 (14)
H31	1.2738	0.0016	0.7393	0.081*
C32	1.0996 (5)	-0.0101 (3)	0.7394 (2)	0.0665 (13)
H32	1.0920	-0.0628	0.7243	0.080*
C33	1.0018 (4)	0.0298 (3)	0.7534 (2)	0.0561 (11)
H33	0.9281	0.0038	0.7470	0.067*
C34	1.0114 (4)	0.1088 (2)	0.77716 (18)	0.0423 (9)

C35	0.9067 (3)	0.1515 (2)	0.79519 (17)	0.0388 (9)
H35A	0.9145	0.2089	0.7883	0.047*
H35B	0.8338	0.1334	0.7674	0.047*
C36	0.8639 (4)	0.0525 (3)	0.8715 (3)	0.0643 (9)
H36A	0.8305	0.0284	0.8310	0.077*
H36B	0.9374	0.0239	0.8890	0.077*
C37	0.7792 (4)	0.0404 (3)	0.9134 (3)	0.0643 (9)
H37A	0.7457	-0.0135	0.9075	0.077*
H37B	0.8211	0.0453	0.9569	0.077*
C38	0.5845 (4)	0.0823 (2)	0.84676 (19)	0.0474 (10)
H38A	0.6153	0.0855	0.8083	0.057*
H38B	0.5243	0.1240	0.8449	0.057*
C39	0.5795 (3)	0.2704 (2)	0.78663 (15)	0.0322 (8)
C40	0.5548 (3)	0.2906 (2)	0.71679 (15)	0.0295 (7)
C41	0.4375 (3)	0.2997 (2)	0.68456 (16)	0.0364 (8)
H41	0.3755	0.2989	0.7066	0.044*
C42	0.4119 (3)	0.3100 (2)	0.61992 (16)	0.0350 (8)
H42	0.3326	0.3154	0.5990	0.042*
C43	0.5016 (3)	0.3125 (2)	0.58600 (15)	0.0293 (7)
C44	0.4709 (3)	0.3139 (2)	0.51426 (16)	0.0335 (8)
C45	0.6199 (3)	0.3074 (2)	0.61860 (16)	0.0371 (9)
H45	0.6820	0.3117	0.5969	0.044*
C46	0.6456 (3)	0.2961 (2)	0.68337 (16)	0.0360 (8)
H46	0.7249	0.2921	0.7046	0.043*
O1	0.4368 (3)	0.05093 (19)	0.92886 (16)	0.0669 (7)
O2	0.4008 (3)	0.19793 (19)	0.97062 (16)	0.0669 (7)
O3	1.1119 (3)	0.35817 (18)	0.87622 (17)	0.0655 (9)
O4	1.1251 (3)	0.22632 (19)	0.80180 (18)	0.0706 (10)
O5	0.4986 (2)	0.2814 (2)	0.81597 (12)	0.0555 (8)
O6	0.6840 (2)	0.24327 (14)	0.80900 (10)	0.0333 (6)
O7	0.3600 (2)	0.31285 (15)	0.49024 (10)	0.0349 (6)
O8	0.5516 (2)	0.3137 (2)	0.48444 (13)	0.0710 (10)
N1	0.6832 (3)	0.09923 (18)	0.90061 (15)	0.0374 (7)
H1N	0.653 (3)	0.100 (2)	0.9325 (13)	0.045*
N2	0.8942 (3)	0.13883 (19)	0.86124 (13)	0.0343 (7)
H2N	0.954 (2)	0.153 (2)	0.8863 (15)	0.041*
N3	0.8662 (3)	0.32447 (18)	0.90502 (14)	0.0353 (7)
H3N	0.9388 (19)	0.317 (2)	0.9209 (17)	0.042*
N4	0.6460 (2)	0.28272 (18)	0.93980 (13)	0.0327 (6)
H4N	0.582 (2)	0.275 (2)	0.9139 (15)	0.039*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.0219 (2)	0.0338 (2)	0.0203 (2)	-0.00261 (19)	0.00421 (14)	0.00109 (18)
C1	0.041 (2)	0.044 (2)	0.038 (2)	-0.0139 (18)	0.0019 (17)	0.0005 (17)
C2	0.071 (3)	0.079 (3)	0.053 (3)	-0.038 (3)	0.025 (2)	-0.024 (2)
C3	0.086 (4)	0.068 (3)	0.074 (4)	-0.034 (3)	0.024 (3)	-0.037 (3)

supplementary materials

C4	0.067 (3)	0.050 (3)	0.069 (3)	-0.027 (2)	0.008 (3)	-0.007 (2)
C5	0.052 (3)	0.058 (3)	0.061 (3)	-0.022 (2)	0.011 (2)	0.000 (2)
C6	0.041 (2)	0.048 (2)	0.055 (3)	-0.0132 (19)	0.0094 (19)	-0.007 (2)
C9	0.037 (2)	0.062 (3)	0.060 (3)	-0.013 (2)	0.017 (2)	0.003 (2)
C10	0.058 (3)	0.063 (3)	0.076 (3)	-0.007 (2)	0.037 (3)	-0.001 (3)
C11	0.041 (2)	0.055 (2)	0.032 (2)	-0.0009 (19)	0.0169 (16)	-0.0058 (17)
C12	0.042 (2)	0.092 (4)	0.066 (3)	0.002 (2)	0.028 (2)	-0.016 (3)
C13	0.080 (4)	0.093 (4)	0.105 (5)	0.010 (3)	0.058 (4)	-0.029 (4)
C14	0.098 (5)	0.088 (4)	0.083 (4)	-0.008 (3)	0.051 (4)	-0.038 (3)
C15	0.065 (3)	0.073 (3)	0.062 (3)	-0.015 (3)	0.029 (2)	-0.024 (3)
C16	0.039 (2)	0.054 (2)	0.0284 (18)	-0.0033 (18)	0.0130 (15)	-0.0051 (17)
C17	0.0337 (19)	0.057 (2)	0.0275 (19)	0.0029 (18)	0.0086 (15)	0.0014 (16)
C18	0.060 (3)	0.038 (2)	0.045 (2)	0.003 (2)	0.024 (2)	-0.0016 (18)
C19	0.062 (3)	0.043 (2)	0.042 (2)	-0.013 (2)	0.022 (2)	-0.0072 (18)
C20	0.055 (2)	0.046 (2)	0.034 (2)	-0.008 (2)	0.0196 (18)	0.0013 (17)
C21	0.059 (3)	0.037 (2)	0.037 (2)	-0.0095 (19)	0.0198 (19)	-0.0019 (16)
C22	0.076 (3)	0.041 (2)	0.072 (3)	0.001 (2)	0.036 (3)	0.001 (2)
C23	0.110 (5)	0.034 (2)	0.104 (5)	0.000 (3)	0.054 (4)	-0.002 (3)
C24	0.093 (4)	0.049 (3)	0.095 (4)	-0.024 (3)	0.041 (3)	-0.011 (3)
C25	0.064 (3)	0.056 (3)	0.079 (3)	-0.023 (2)	0.028 (3)	-0.009 (2)
C26	0.060 (3)	0.038 (2)	0.053 (3)	-0.012 (2)	0.022 (2)	0.0032 (18)
C27	0.041 (3)	0.061 (3)	0.103 (4)	-0.012 (2)	0.017 (3)	0.005 (3)
C28	0.041 (3)	0.078 (4)	0.112 (5)	0.004 (3)	-0.007 (3)	-0.016 (3)
C29	0.043 (2)	0.047 (2)	0.044 (2)	0.0136 (19)	0.0114 (18)	0.0012 (18)
C30	0.046 (3)	0.079 (3)	0.064 (3)	0.017 (2)	0.020 (2)	0.008 (3)
C31	0.071 (3)	0.079 (3)	0.059 (3)	0.035 (3)	0.029 (3)	0.000 (3)
C32	0.079 (4)	0.061 (3)	0.068 (3)	0.018 (3)	0.034 (3)	-0.007 (2)
C33	0.065 (3)	0.052 (2)	0.057 (3)	0.007 (2)	0.026 (2)	-0.004 (2)
C34	0.047 (2)	0.046 (2)	0.038 (2)	0.0129 (19)	0.0186 (18)	0.0065 (17)
C35	0.040 (2)	0.045 (2)	0.034 (2)	0.0050 (17)	0.0133 (16)	0.0007 (16)
C36	0.068 (2)	0.0466 (17)	0.085 (3)	0.0102 (17)	0.0316 (19)	0.0148 (17)
C37	0.068 (2)	0.0466 (17)	0.085 (3)	0.0102 (17)	0.0316 (19)	0.0148 (17)
C38	0.045 (2)	0.052 (2)	0.041 (2)	-0.020 (2)	0.0005 (18)	0.0051 (18)
C39	0.0295 (18)	0.042 (2)	0.0235 (17)	0.0001 (16)	0.0031 (14)	0.0026 (14)
C40	0.0281 (16)	0.0335 (17)	0.0256 (16)	0.0029 (16)	0.0026 (13)	-0.0014 (14)
C41	0.0271 (17)	0.057 (2)	0.0259 (18)	0.0063 (17)	0.0065 (14)	-0.0033 (16)
C42	0.0227 (16)	0.053 (2)	0.0271 (18)	0.0100 (16)	-0.0004 (14)	-0.0020 (16)
C43	0.0266 (17)	0.0344 (17)	0.0252 (17)	0.0023 (15)	0.0013 (13)	0.0025 (14)
C44	0.0266 (18)	0.049 (2)	0.0247 (18)	0.0046 (16)	0.0034 (14)	0.0023 (15)
C45	0.0240 (17)	0.059 (2)	0.0280 (18)	-0.0045 (17)	0.0051 (14)	0.0066 (16)
C46	0.0234 (16)	0.051 (2)	0.0304 (18)	-0.0007 (16)	-0.0030 (14)	0.0055 (16)
O1	0.0562 (14)	0.0712 (15)	0.0840 (17)	-0.0255 (12)	0.0396 (12)	-0.0315 (13)
O2	0.0562 (14)	0.0712 (15)	0.0840 (17)	-0.0255 (12)	0.0396 (12)	-0.0315 (13)
O3	0.059 (2)	0.0482 (17)	0.096 (3)	-0.0060 (16)	0.0329 (18)	0.0088 (17)
O4	0.0445 (18)	0.061 (2)	0.099 (3)	0.0058 (16)	-0.0041 (17)	-0.0072 (18)
O5	0.0341 (14)	0.103 (2)	0.0301 (14)	0.0175 (16)	0.0081 (11)	0.0096 (15)
O6	0.0264 (12)	0.0468 (14)	0.0250 (12)	0.0031 (11)	0.0017 (10)	0.0051 (10)
O7	0.0254 (12)	0.0561 (15)	0.0222 (12)	0.0011 (11)	0.0022 (9)	-0.0025 (11)
O8	0.0278 (14)	0.157 (3)	0.0285 (15)	0.0042 (18)	0.0065 (12)	0.0067 (18)

N1	0.0338 (17)	0.0397 (17)	0.0370 (18)	-0.0087 (14)	0.0029 (14)	0.0030 (14)
N2	0.0317 (16)	0.0443 (17)	0.0275 (16)	0.0010 (15)	0.0077 (12)	0.0011 (13)
N3	0.0359 (17)	0.0403 (16)	0.0317 (17)	-0.0076 (15)	0.0119 (13)	-0.0028 (13)
N4	0.0274 (15)	0.0459 (17)	0.0261 (15)	0.0005 (15)	0.0084 (11)	0.0014 (14)

Geometric parameters (Å, °)

Ni1—O6	2.049 (2)	C24—H24	0.9300
Ni1—O7 ⁱ	2.089 (2)	C25—C26	1.376 (6)
Ni1—N1	2.131 (3)	C25—H25	0.9300
Ni1—N3	2.142 (3)	C26—O3	1.377 (5)
Ni1—N4	2.151 (3)	C27—O3	1.399 (5)
Ni1—N2	2.160 (3)	C27—C28	1.465 (7)
C1—C6	1.377 (6)	C27—H27A	0.9700
C1—C2	1.384 (6)	C27—H27B	0.9700
C1—C38	1.508 (5)	C28—O4	1.359 (5)
C2—C3	1.393 (6)	C28—H28A	0.9700
C2—H2	0.9300	C28—H28B	0.9700
C3—C4	1.356 (7)	C29—O4	1.377 (5)
C3—H3	0.9300	C29—C34	1.382 (6)
C4—C5	1.370 (6)	C29—C30	1.394 (6)
C4—H4	0.9300	C30—C31	1.376 (7)
C5—C6	1.374 (5)	C30—H30	0.9300
C5—H5	0.9300	C31—C32	1.369 (7)
C6—O1	1.385 (5)	C31—H31	0.9300
C9—O1	1.383 (5)	C32—C33	1.383 (6)
C9—C10	1.471 (6)	C32—H32	0.9300
C9—H9A	0.9700	C33—C34	1.402 (6)
C9—H9B	0.9700	C33—H33	0.9300
C10—O2	1.382 (5)	C34—C35	1.506 (5)
C10—H10A	0.9700	C35—N2	1.482 (4)
C10—H10B	0.9700	C35—H35A	0.9700
C11—C16	1.381 (5)	C35—H35B	0.9700
C11—C12	1.383 (6)	C36—C37	1.469 (6)
C11—O2	1.389 (5)	C36—N2	1.500 (5)
C12—C13	1.380 (7)	C36—H36A	0.9700
C12—H12	0.9300	C36—H36B	0.9700
C13—C14	1.366 (8)	C37—N1	1.450 (5)
C13—H13	0.9300	C37—H37A	0.9700
C14—C15	1.380 (7)	C37—H37B	0.9700
C14—H14	0.9300	C38—N1	1.474 (5)
C15—C16	1.389 (6)	C38—H38A	0.9700
C15—H15	0.9300	C38—H38B	0.9700
C16—C17	1.512 (5)	C39—O5	1.236 (4)
C17—N4	1.485 (4)	C39—O6	1.274 (4)
C17—H17A	0.9700	C39—C40	1.519 (4)
C17—H17B	0.9700	C40—C46	1.382 (5)
C18—N4	1.481 (5)	C40—C41	1.386 (4)
C18—C19	1.508 (6)	C41—C42	1.382 (5)

supplementary materials

C18—H18A	0.9700	C41—H41	0.9300
C18—H18B	0.9700	C42—C43	1.377 (5)
C19—N3	1.478 (5)	C42—H42	0.9300
C19—H19A	0.9700	C43—C45	1.393 (4)
C19—H19B	0.9700	C43—C44	1.523 (5)
C20—N3	1.479 (5)	C44—O8	1.228 (4)
C20—C21	1.510 (5)	C44—O7	1.267 (4)
C20—H20A	0.9700	C45—C46	1.387 (5)
C20—H20B	0.9700	C45—H45	0.9300
C21—C26	1.384 (6)	C46—H46	0.9300
C21—C22	1.390 (6)	O7—Ni1 ⁱⁱ	2.089 (2)
C22—C23	1.401 (7)	N1—H1N	0.83 (2)
C22—H22	0.9300	N2—H2N	0.81 (2)
C23—C24	1.343 (7)	N3—H3N	0.84 (2)
C23—H23	0.9300	N4—H4N	0.83 (2)
C24—C25	1.376 (7)		
O6—Ni1—O7 ⁱ	177.49 (10)	C28—C27—H27A	109.6
O6—Ni1—N1	95.97 (11)	O3—C27—H27B	109.6
O7 ⁱ —Ni1—N1	86.54 (11)	C28—C27—H27B	109.6
O6—Ni1—N3	89.42 (11)	H27A—C27—H27B	108.1
O7 ⁱ —Ni1—N3	88.08 (11)	O4—C28—C27	111.5 (4)
N1—Ni1—N3	174.59 (12)	O4—C28—H28A	109.3
O6—Ni1—N4	92.13 (10)	C27—C28—H28A	109.3
O7 ⁱ —Ni1—N4	87.37 (10)	O4—C28—H28B	109.3
N1—Ni1—N4	97.09 (12)	C27—C28—H28B	109.3
N3—Ni1—N4	83.23 (12)	H28A—C28—H28B	108.0
O6—Ni1—N2	90.38 (10)	O4—C29—C34	116.1 (3)
O7 ⁱ —Ni1—N2	90.20 (10)	O4—C29—C30	122.3 (4)
N1—Ni1—N2	81.13 (12)	C34—C29—C30	121.5 (4)
N3—Ni1—N2	98.32 (12)	C31—C30—C29	119.0 (5)
N4—Ni1—N2	177.07 (11)	C31—C30—H30	120.5
C6—C1—C2	117.0 (4)	C29—C30—H30	120.5
C6—C1—C38	120.8 (4)	C32—C31—C30	121.2 (4)
C2—C1—C38	122.2 (4)	C32—C31—H31	119.4
C1—C2—C3	121.4 (4)	C30—C31—H31	119.4
C1—C2—H2	119.3	C31—C32—C33	119.3 (5)
C3—C2—H2	119.3	C31—C32—H32	120.4
C4—C3—C2	119.8 (5)	C33—C32—H32	120.4
C4—C3—H3	120.1	C32—C33—C34	121.4 (5)
C2—C3—H3	120.1	C32—C33—H33	119.3
C3—C4—C5	119.9 (4)	C34—C33—H33	119.3
C3—C4—H4	120.1	C29—C34—C33	117.5 (4)
C5—C4—H4	120.1	C29—C34—C35	120.7 (4)
C4—C5—C6	120.1 (5)	C33—C34—C35	121.8 (4)
C4—C5—H5	119.9	N2—C35—C34	115.0 (3)
C6—C5—H5	119.9	N2—C35—H35A	108.5
C5—C6—C1	121.8 (4)	C34—C35—H35A	108.5

C5—C6—O1	123.3 (4)	N2—C35—H35B	108.5
C1—C6—O1	114.9 (3)	C34—C35—H35B	108.5
O1—C9—C10	111.3 (3)	H35A—C35—H35B	107.5
O1—C9—H9A	109.4	C37—C36—N2	115.0 (4)
C10—C9—H9A	109.4	C37—C36—H36A	108.5
O1—C9—H9B	109.4	N2—C36—H36A	108.5
C10—C9—H9B	109.4	C37—C36—H36B	108.5
H9A—C9—H9B	108.0	N2—C36—H36B	108.5
O2—C10—C9	112.6 (4)	H36A—C36—H36B	107.5
O2—C10—H10A	109.1	N1—C37—C36	110.8 (4)
C9—C10—H10A	109.1	N1—C37—H37A	109.5
O2—C10—H10B	109.1	C36—C37—H37A	109.5
C9—C10—H10B	109.1	N1—C37—H37B	109.5
H10A—C10—H10B	107.8	C36—C37—H37B	109.5
C16—C11—C12	121.8 (4)	H37A—C37—H37B	108.1
C16—C11—O2	116.6 (3)	N1—C38—C1	115.0 (3)
C12—C11—O2	121.6 (4)	N1—C38—H38A	108.5
C13—C12—C11	119.2 (5)	C1—C38—H38A	108.5
C13—C12—H12	120.4	N1—C38—H38B	108.5
C11—C12—H12	120.4	C1—C38—H38B	108.5
C14—C13—C12	119.8 (5)	H38A—C38—H38B	107.5
C14—C13—H13	120.1	O5—C39—O6	126.5 (3)
C12—C13—H13	120.1	O5—C39—C40	118.3 (3)
C13—C14—C15	120.8 (5)	O6—C39—C40	115.2 (3)
C13—C14—H14	119.6	C46—C40—C41	118.6 (3)
C15—C14—H14	119.6	C46—C40—C39	122.0 (3)
C14—C15—C16	120.5 (5)	C41—C40—C39	119.3 (3)
C14—C15—H15	119.8	C42—C41—C40	120.5 (3)
C16—C15—H15	119.8	C42—C41—H41	119.8
C11—C16—C15	117.8 (4)	C40—C41—H41	119.8
C11—C16—C17	120.7 (3)	C43—C42—C41	121.2 (3)
C15—C16—C17	121.5 (4)	C43—C42—H42	119.4
N4—C17—C16	115.2 (3)	C41—C42—H42	119.4
N4—C17—H17A	108.5	C42—C43—C45	118.5 (3)
C16—C17—H17A	108.5	C42—C43—C44	120.3 (3)
N4—C17—H17B	108.5	C45—C43—C44	121.1 (3)
C16—C17—H17B	108.5	O8—C44—O7	125.2 (3)
H17A—C17—H17B	107.5	O8—C44—C43	119.8 (3)
N4—C18—C19	109.1 (3)	O7—C44—C43	114.9 (3)
N4—C18—H18A	109.9	C46—C45—C43	120.3 (3)
C19—C18—H18A	109.9	C46—C45—H45	119.9
N4—C18—H18B	109.9	C43—C45—H45	119.9
C19—C18—H18B	109.9	C40—C46—C45	120.9 (3)
H18A—C18—H18B	108.3	C40—C46—H46	119.6
N3—C19—C18	108.5 (3)	C45—C46—H46	119.6
N3—C19—H19A	110.0	C9—O1—C6	119.8 (3)
C18—C19—H19A	110.0	C10—O2—C11	116.4 (3)
N3—C19—H19B	110.0	C26—O3—C27	118.1 (3)
C18—C19—H19B	110.0	C28—O4—C29	118.8 (4)

supplementary materials

H19A—C19—H19B	108.4	C39—O6—Ni1	132.4 (2)
N3—C20—C21	114.5 (3)	C44—O7—Ni1 ⁱⁱ	130.8 (2)
N3—C20—H20A	108.6	C37—N1—C38	116.6 (4)
C21—C20—H20A	108.6	C37—N1—Ni1	105.4 (3)
N3—C20—H20B	108.6	C38—N1—Ni1	115.7 (2)
C21—C20—H20B	108.6	C37—N1—H1N	106 (3)
H20A—C20—H20B	107.6	C38—N1—H1N	106 (3)
C26—C21—C22	118.3 (4)	Ni1—N1—H1N	106 (3)
C26—C21—C20	119.8 (4)	C35—N2—C36	110.4 (3)
C22—C21—C20	121.8 (4)	C35—N2—Ni1	118.8 (2)
C21—C22—C23	119.8 (5)	C36—N2—Ni1	107.4 (2)
C21—C22—H22	120.1	C35—N2—H2N	112 (3)
C23—C22—H22	120.1	C36—N2—H2N	112 (3)
C24—C23—C22	120.2 (5)	Ni1—N2—H2N	96 (3)
C24—C23—H23	119.9	C19—N3—C20	112.3 (3)
C22—C23—H23	119.9	C19—N3—Ni1	104.5 (2)
C23—C24—C25	121.1 (5)	C20—N3—Ni1	119.3 (2)
C23—C24—H24	119.5	C19—N3—H3N	106 (3)
C25—C24—H24	119.5	C20—N3—H3N	104 (3)
C26—C25—C24	119.2 (5)	Ni1—N3—H3N	110 (3)
C26—C25—H25	120.4	C18—N4—C17	112.2 (3)
C24—C25—H25	120.4	C18—N4—Ni1	105.4 (2)
C25—C26—O3	123.3 (4)	C17—N4—Ni1	118.3 (2)
C25—C26—C21	121.3 (4)	C18—N4—H4N	109 (3)
O3—C26—C21	115.4 (3)	C17—N4—H4N	110 (3)
O3—C27—C28	110.2 (4)	Ni1—N4—H4N	101 (3)
O3—C27—H27A	109.6		
C6—C1—C2—C3	1.5 (7)	C41—C40—C46—C45	-2.4 (5)
C38—C1—C2—C3	-177.8 (4)	C39—C40—C46—C45	174.2 (3)
C1—C2—C3—C4	0.3 (8)	C43—C45—C46—C40	-0.8 (6)
C2—C3—C4—C5	-0.9 (8)	C10—C9—O1—C6	-178.7 (4)
C3—C4—C5—C6	-0.4 (8)	C5—C6—O1—C9	-34.0 (6)
C4—C5—C6—C1	2.3 (7)	C1—C6—O1—C9	149.7 (4)
C4—C5—C6—O1	-173.8 (4)	C9—C10—O2—C11	178.9 (4)
C2—C1—C6—C5	-2.8 (6)	C16—C11—O2—C10	131.3 (4)
C38—C1—C6—C5	176.5 (4)	C12—C11—O2—C10	-50.8 (6)
C2—C1—C6—O1	173.6 (4)	C25—C26—O3—C27	-36.7 (7)
C38—C1—C6—O1	-7.0 (6)	C21—C26—O3—C27	143.7 (4)
O1—C9—C10—O2	41.1 (6)	C28—C27—O3—C26	-160.0 (4)
C16—C11—C12—C13	-1.6 (7)	C27—C28—O4—C29	176.2 (4)
O2—C11—C12—C13	-179.3 (5)	C34—C29—O4—C28	140.7 (5)
C11—C12—C13—C14	-0.5 (9)	C30—C29—O4—C28	-41.9 (7)
C12—C13—C14—C15	1.7 (10)	O5—C39—O6—Ni1	-4.2 (6)
C13—C14—C15—C16	-0.9 (9)	C40—C39—O6—Ni1	175.8 (2)
C12—C11—C16—C15	2.4 (6)	N1—Ni1—O6—C39	78.3 (3)
O2—C11—C16—C15	-179.8 (4)	N3—Ni1—O6—C39	-102.3 (3)
C12—C11—C16—C17	-178.1 (4)	N4—Ni1—O6—C39	-19.1 (3)
O2—C11—C16—C17	-0.2 (6)	N2—Ni1—O6—C39	159.4 (3)

C14—C15—C16—C11	-1.1 (7)	O8—C44—O7—Ni1 ⁱⁱ	14.3 (6)
C14—C15—C16—C17	179.3 (5)	C43—C44—O7—Ni1 ⁱⁱ	-163.6 (2)
C11—C16—C17—N4	83.3 (5)	C36—C37—N1—C38	81.0 (5)
C15—C16—C17—N4	-97.2 (5)	C36—C37—N1—Ni1	-48.9 (5)
N4—C18—C19—N3	61.8 (4)	C1—C38—N1—C37	54.7 (5)
N3—C20—C21—C26	71.6 (5)	C1—C38—N1—Ni1	179.6 (3)
N3—C20—C21—C22	-110.4 (4)	O6—Ni1—N1—C37	122.4 (3)
C26—C21—C22—C23	-0.6 (7)	O7 ⁱ —Ni1—N1—C37	-57.8 (3)
C20—C21—C22—C23	-178.6 (4)	N4—Ni1—N1—C37	-144.7 (3)
C21—C22—C23—C24	0.8 (8)	N2—Ni1—N1—C37	32.9 (3)
C22—C23—C24—C25	-0.9 (9)	O6—Ni1—N1—C38	-8.1 (3)
C23—C24—C25—C26	0.8 (8)	O7 ⁱ —Ni1—N1—C38	171.7 (3)
C24—C25—C26—O3	179.8 (5)	N4—Ni1—N1—C38	84.8 (3)
C24—C25—C26—C21	-0.6 (7)	N2—Ni1—N1—C38	-97.6 (3)
C22—C21—C26—C25	0.5 (7)	C34—C35—N2—C36	-66.6 (4)
C20—C21—C26—C25	178.6 (4)	C34—C35—N2—Ni1	168.8 (3)
C22—C21—C26—O3	-179.9 (4)	C37—C36—N2—C35	-142.5 (4)
C20—C21—C26—O3	-1.8 (6)	C37—C36—N2—Ni1	-11.6 (5)
O3—C27—C28—O4	56.8 (7)	O6—Ni1—N2—C35	18.2 (3)
O4—C29—C30—C31	-177.0 (4)	O7 ⁱ —Ni1—N2—C35	-159.3 (3)
C34—C29—C30—C31	0.2 (7)	N1—Ni1—N2—C35	114.2 (3)
C29—C30—C31—C32	1.8 (7)	N3—Ni1—N2—C35	-71.2 (3)
C30—C31—C32—C33	-1.4 (8)	O6—Ni1—N2—C36	-107.9 (3)
C31—C32—C33—C34	-1.1 (7)	O7 ⁱ —Ni1—N2—C36	74.6 (3)
O4—C29—C34—C33	174.9 (4)	N1—Ni1—N2—C36	-11.9 (3)
C30—C29—C34—C33	-2.4 (6)	N3—Ni1—N2—C36	162.6 (3)
O4—C29—C34—C35	-4.8 (6)	C18—C19—N3—C20	83.7 (4)
C30—C29—C34—C35	177.8 (4)	C18—C19—N3—Ni1	-47.0 (3)
C32—C33—C34—C29	2.9 (6)	C21—C20—N3—C19	61.7 (4)
C32—C33—C34—C35	-177.4 (4)	C21—C20—N3—Ni1	-175.6 (3)
C29—C34—C35—N2	-91.4 (4)	O6—Ni1—N3—C19	111.2 (2)
C33—C34—C35—N2	88.8 (5)	O7 ⁱ —Ni1—N3—C19	-68.6 (2)
N2—C36—C37—N1	41.6 (6)	N4—Ni1—N3—C19	19.0 (2)
C6—C1—C38—N1	76.5 (5)	N2—Ni1—N3—C19	-158.5 (2)
C2—C1—C38—N1	-104.2 (5)	O6—Ni1—N3—C20	-15.3 (3)
O5—C39—C40—C46	166.5 (4)	O7 ⁱ —Ni1—N3—C20	164.9 (3)
O6—C39—C40—C46	-13.6 (5)	N4—Ni1—N3—C20	-107.5 (3)
O5—C39—C40—C41	-16.9 (5)	N2—Ni1—N3—C20	75.0 (3)
O6—C39—C40—C41	163.0 (3)	C19—C18—N4—C17	89.0 (4)
C46—C40—C41—C42	3.2 (5)	C19—C18—N4—Ni1	-41.1 (3)
C39—C40—C41—C42	-173.5 (3)	C16—C17—N4—C18	69.4 (4)
C40—C41—C42—C43	-0.8 (6)	C16—C17—N4—Ni1	-167.5 (2)
C41—C42—C43—C45	-2.5 (5)	O6—Ni1—N4—C18	-77.3 (2)
C41—C42—C43—C44	172.8 (3)	O7 ⁱ —Ni1—N4—C18	100.2 (2)
C42—C43—C44—O8	-177.9 (4)	N1—Ni1—N4—C18	-173.6 (2)
C45—C43—C44—O8	-2.8 (6)	N3—Ni1—N4—C18	11.9 (2)
C42—C43—C44—O7	0.2 (5)	O6—Ni1—N4—C17	156.3 (2)

supplementary materials

C45—C43—C44—O7	175.3 (3)	O7 ⁱ —Ni1—N4—C17	-26.1 (2)
C42—C43—C45—C46	3.2 (5)	N1—Ni1—N4—C17	60.0 (3)
C44—C43—C45—C46	-172.0 (3)	N3—Ni1—N4—C17	-114.5 (3)

Symmetry codes: (i) $x+1/2, -y+1/2, z+1/2$; (ii) $x-1/2, -y+1/2, z-1/2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1N \cdots O1	0.83 (2)	2.58 (4)	3.101 (4)	122 (3)
N2—H2N \cdots O8 ⁱ	0.81 (2)	2.26 (2)	3.003 (4)	152 (4)
N3—H3N \cdots O3	0.84 (2)	2.47 (3)	3.044 (4)	127 (3)
N4—H4N \cdots O5	0.83 (2)	2.14 (3)	2.862 (4)	144 (4)
C10—H10A \cdots O8 ⁱⁱⁱ	0.97	2.40	3.359 (6)	172
C12—H12 \cdots O8 ⁱⁱⁱ	0.93	2.60	3.508 (6)	166
C28—H28B \cdots O2 ^{iv}	0.97	2.47	3.210 (7)	133
C28—H28A \cdots O5 ^{iv}	0.97	2.47	3.342 (6)	149
C35—H35A \cdots O4	0.97	2.38	2.761 (5)	103
C37—H37B \cdots O7 ⁱ	0.97	2.47	2.985 (6)	113
C42—H42 \cdots O7	0.93	2.44	2.752 (4)	100

Symmetry codes: (i) $x+1/2, -y+1/2, z+1/2$; (iii) $x-1/2, -y+1/2, z+1/2$; (iv) $x+1, y, z$.

Fig. 1




Fig. 2

