

Tris(tetramethylammonium) tetra- μ_2 -sulfido-tetrasulfidocopper(I)dimolybdenum(VI) *N,N*-dimethylformamide solvate

Yuan Cao,^a Yulin Chen^b and Guosheng Cheng^{a*}

^aInstitute of Transition Metallic Compounds, Nanjing University of Information Science and Technology, 219 Ninliu Road, Nanjing 210044, Jiangsu, People's Republic of China, and ^bSchool of Mathematics and Physics, Nanjing University of Information Science and Technology, 219 Ninliu Road, Nanjing 210044, Jiangsu, People's Republic of China

Correspondence e-mail: caoyuannust@yahoo.com.cn


Received 26 September 2008; accepted 6 November 2008

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{O}-\text{C}) = 0.009$ Å; R factor = 0.042; wR factor = 0.136; data-to-parameter ratio = 24.2.

The title compound, $(\text{C}_4\text{H}_{12}\text{N})_3[\text{CuMo}_2\text{S}_8]\cdot\text{C}_3\text{H}_7\text{NO}$, was obtained from the self-assembly of tetrathiomolybdate, tetramethylammonium nitrate and cuprous sulfide in dimethylformamide (DMF). The asymmetric unit contains three $(\text{NMe}_4)^+$ cations, one $[\text{Mo}_2\text{S}_8\text{Cu}]^{3-}$ anion and one DMF solvent molecule, and no obvious interactions are observed between these species. The trinuclear anion can be viewed as fused $[\text{MoS}_4\text{Cu}]^-$ units sharing a copper center. The geometric parameters of the trivalent anion are comparable to those reported for other related salts including isomorphous anions, namely $(\text{NEt}_4)_2(\text{PPh}_4)[\text{Mo}_2\text{S}_8\text{Cu}]$ (*a*) and $(\text{Ph}_3\text{P}=\text{N}=\text{PPh}_3)_2(\text{NEt}_4)[\text{W}_2\text{S}_8\text{Cu}]\cdot 2\text{CH}_3\text{CN}$ (*b*). However, the Mo—Cu—Mo angle is found to be $160.24(3)^\circ$ for the title salt, while this angle is $162.97(2)^\circ$ in (*a*) and the W—Cu—W angle is $170.3(2)^\circ$ in (*b*), indicating that the largest deviation from linearity is in the title compound.

Related literature

For related $\text{Mo}^{\text{VI}}/\text{Cu}^{\text{I}}$ and $\text{W}^{\text{VI}}/\text{Cu}^{\text{I}}$ complexes, see: Niu *et al.* (2002); Maiti *et al.* (2004); Müller *et al.* (1989).



Experimental

Crystal data

$(\text{C}_4\text{H}_{12}\text{N})_3[\text{CuMo}_2\text{S}_8]\cdot\text{C}_3\text{H}_7\text{NO}$
 $M_r = 807.43$
 Monoclinic, $P2_1/n$
 $a = 9.4380(19)$ Å
 $b = 20.336(4)$ Å
 $c = 17.718(4)$ Å
 $\beta = 98.60(3)^\circ$

$V = 3362.4(12)$ Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.87$ mm⁻¹
 $T = 293(2)$ K
 $0.40 \times 0.30 \times 0.25$ mm

Data collection

Bruker APEX CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 1998)
 $T_{\text{min}} = 0.527$, $T_{\text{max}} = 0.618$

17155 measured reflections
 6625 independent reflections
 5061 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.039$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.136$
 $S = 1.05$
 6625 reflections

274 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.93$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.85$ e Å⁻³

Data collection: SMART (Bruker, 1998); cell refinement: SAINT (Bruker, 1998); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL.

Financial support from NUIST is acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2199).

References

- Bruker (1998). SMART, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Maiti, B., Pal, K. & Sarkar, S. (2004). *Inorg. Chem. Commun.* **7**, 1027–1029.
- Müller, A., Bögge, H., Schimanski, U., Penk, M., Nieradzik, K., Dartmann, M., Krickemeyer, E., Schimanski, J., Römer, C., Römer, M., Dornfeld, H., Wienböcker, U., Hellmann, W. & Zimmermann, M. (1989). *Monatsh. Chem.* **120**, 367–391.
- Niu, Y.-Y., Chen, T.-N., Liu, S.-X., Song, Y.-L., Wang, Y.-X., Xue, Z.-L. & Xin, X.-Q. (2002). *J. Chem. Soc. Dalton Trans.* pp. 1980–1984.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, m1546 [doi:10.1107/S1600536808036532]

Tris(tetramethylammonium) tetra- μ_2 -sulfido-tetrakisulfidocopper(I)dimolybdenum(VI) *N,N*-dimethylformamide solvate

Y. Cao, Y. Chen and G. Cheng

Comment

The title compound, $(\text{NMe}_4)_3[\text{Mo}_2\text{S}_8\text{Cu}]\cdot\text{C}_3\text{H}_7\text{ON}$, was obtained from the self-assembly of tetrathiomolybdate, tetramethylammonium nitrate and cuprous sulfide in *N,N*-dimethylformamide (DMF). The asymmetric unit contains three tetramethylammonium cations, one $[\text{Mo}_2\text{S}_8\text{Cu}]^{3-}$ anion and one DMF solvent molecule, with no obvious interactions observed among these species. The trinuclear anion can be viewed as fused by two binuclear units $[\text{MoS}_4\text{Cu}]^-$ through one shared copper. The two binuclear units are perpendicular to each other, as found in other linear trinuclear clusters, such as $\{\text{MoS}_4\text{Cu}_2[\text{PPh}_2(\text{py})_4]\}$ (py = pyridyl) (Niu *et al.*, 2002). However, in the latter, $[\text{MoS}_4]^{2-}$ is a tetradentate ligand, while in the former, it serves as a bidentate ligand. In the title cluster, one Mo center is close to retain the original tetrahedral configuration of free $[\text{MoS}_4]^{2-}$, with six S—Mo—S bond angles varying from 108.88 (6) to 111.20 (6)°. The other Mo center distorts slightly more, the corresponding angles ranging from 107.24 (6) to 111.79 (6)°. The geometric parameters of the trivalent anion are comparable to those reported for the compounds bis(tetraethylammonium) tetraphenylphosphonium bis(di- μ_2 -sulfido-dithioxomolybdenum)copper (*a*) (Maiti *et al.*, 2004) and bis[bis(triphenylphosphine)iminium] tetraethylammonium bis(di- μ_2 -sulfido-dithioxotungsten)copper acetonitrile disolvate (*b*) (Müller *et al.*, 1989), except a notable difference among the Mo—Cu—Mo angles: 160.24 (3)° for the title compound, 162.97 (2)° for (*a*), while W—Cu—W angle is 170.3 (2)° in (*b*), indicating a largest deviation from ideal linear array in the title salt.

Experimental

0.75 mmol of Cu_2S powder was added to a solution of $[\text{NH}_4]_2\text{MoS}_4$ (1 mmol in 10 ml DMF) and stirred for 4 h. at room temperature. After filtration, the filtrate was carefully laid on the surface with DMF (3 ml) and saturated $(\text{Me}_4\text{N})\text{NO}_3$ solution (5 ml in methanol), successively. Black red polyhedral crystals of the title compound were obtained after two weeks.

Refinement

All H atoms were positioned geometrically and refined using a riding model, with C—H = 0.93 Å for aldehydic H atom and C—H = 0.96 Å for methyl groups. Isotropic displacement parameters for H atoms were calculated as $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{carrier C})$, except for H11A: $U_{\text{iso}}(\text{H11A}) = 1.2U_{\text{eq}}(\text{C11})$.

Figures




Fig. 1. The molecular structure of the title compound, with atom labels and 50% probability displacement ellipsoids. All H atoms have been omitted.

Tris(tetramethylammonium) tetra- μ_2 -sulfido-tetrasulfidocopper(I)dimolybdenum(VI) *N,N*-dimethylformamide solvate

Crystal data

$(C_4H_{12}N)_3[CuMo_2S_8] \cdot C_3H_7NO$

$M_r = 807.43$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2yn$

$a = 9.4380$ (19) Å

$b = 20.336$ (4) Å

$c = 17.718$ (4) Å

$\beta = 98.60$ (3)°

$V = 3362.4$ (12) Å³

$Z = 4$

$F_{000} = 1640$

$D_x = 1.595$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 7032 reflections

$\theta = 1.2$ – 26.1 °

$\mu = 1.87$ mm⁻¹

$T = 293$ (2) K

Polyhedron, black red

$0.40 \times 0.30 \times 0.25$ mm

Data collection

Bruker APEX CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293$ (2) K

φ and ω scans

Absorption correction: multi-scan (SADABS; Bruker, 1998)

$T_{\min} = 0.527$, $T_{\max} = 0.618$

17155 measured reflections

6625 independent reflections

5061 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.039$

$\theta_{\max} = 26.0$ °

$\theta_{\min} = 1.5$ °

$h = -11 \rightarrow 11$

$k = -24 \rightarrow 25$

$l = -21 \rightarrow 21$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.042$

$wR(F^2) = 0.136$

$S = 1.05$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0646P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

6625 reflections $\Delta\rho_{\max} = 0.93 \text{ e } \text{\AA}^{-3}$
 274 parameters $\Delta\rho_{\min} = -0.85 \text{ e } \text{\AA}^{-3}$
 Primary atom site location: structure-invariant direct methods Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Mo1	0.25496 (5)	0.15816 (2)	0.93899 (3)	0.04775 (14)
Mo2	0.31519 (5)	0.05342 (2)	0.66754 (3)	0.04773 (14)
Cu1	0.24065 (7)	0.09541 (3)	0.80223 (4)	0.04912 (18)
S1	0.11495 (15)	0.18682 (7)	0.83342 (8)	0.0494 (3)
S2	0.31587 (15)	0.05454 (7)	0.92589 (8)	0.0489 (3)
S3	0.12487 (15)	0.02298 (7)	0.71839 (8)	0.0483 (3)
S4	0.43472 (16)	0.12836 (7)	0.74081 (8)	0.0514 (3)
S5	0.45325 (16)	-0.03117 (8)	0.65994 (8)	0.0532 (3)
S6	0.24512 (16)	0.09507 (7)	0.55703 (9)	0.0543 (3)
S7	0.44848 (16)	0.21739 (7)	0.95638 (8)	0.0516 (3)
S8	0.13280 (16)	0.16860 (7)	1.03204 (8)	0.0522 (3)
O1	0.1656 (5)	0.1069 (2)	1.2555 (3)	0.0610 (11)
N1	0.2487 (5)	0.1922 (2)	1.3279 (3)	0.0500 (11)
C11	0.2068 (7)	0.1295 (3)	1.3196 (4)	0.0553 (14)
H11A	0.2088	0.1028	1.3624	0.066*
C12	0.2912 (7)	0.2217 (3)	1.4013 (4)	0.0605 (15)
H12A	0.3093	0.1879	1.4393	0.091*
H12B	0.2160	0.2499	1.4133	0.091*
H12C	0.3767	0.2471	1.4006	0.091*
C13	0.2171 (7)	0.2367 (3)	1.2639 (4)	0.0579 (15)
H13A	0.1901	0.2118	1.2179	0.087*
H13B	0.3006	0.2625	1.2592	0.087*
H13C	0.1398	0.2653	1.2719	0.087*
N2	0.7731 (5)	0.1115 (2)	0.5453 (3)	0.0507 (11)
C21	0.8726 (7)	0.1130 (3)	0.4883 (4)	0.0616 (16)
H21A	0.9607	0.0916	0.5089	0.092*
H21B	0.8918	0.1578	0.4761	0.092*
H21C	0.8301	0.0905	0.4429	0.092*
C22	0.8482 (7)	0.1530 (3)	0.6068 (3)	0.0549 (14)
H22A	0.9399	0.1340	0.6254	0.082*
H22B	0.7922	0.1557	0.6476	0.082*
H22C	0.8613	0.1963	0.5873	0.082*
C23	0.6306 (7)	0.1439 (3)	0.5187 (4)	0.0600 (15)
H23A	0.5717	0.1411	0.5583	0.090*
H23B	0.5839	0.1219	0.4738	0.090*
H23C	0.6457	0.1892	0.5070	0.090*
C24	0.7503 (7)	0.0474 (3)	0.5762 (4)	0.0557 (14)
H24A	0.8411	0.0269	0.5931	0.084*
H24B	0.6960	0.0207	0.5376	0.084*
H24C	0.6985	0.0519	0.6186	0.084*

supplementary materials

N3	0.7749 (5)	0.0309 (2)	0.8680 (3)	0.0525 (11)
C31	0.9019 (7)	-0.0048 (3)	0.9017 (3)	0.0554 (14)
H31A	0.9818	0.0248	0.9108	0.083*
H31B	0.9224	-0.0389	0.8674	0.083*
H31C	0.8857	-0.0241	0.9491	0.083*
C32	0.7374 (7)	0.0836 (3)	0.9250 (4)	0.0574 (14)
H32A	0.8146	0.1146	0.9350	0.086*
H32B	0.7231	0.0626	0.9719	0.086*
H32C	0.6513	0.1061	0.9036	0.086*
C33	0.7972 (6)	0.0578 (3)	0.7979 (4)	0.0577 (14)
H33A	0.8760	0.0881	0.8061	0.087*
H33B	0.7123	0.0805	0.7751	0.087*
H33C	0.8187	0.0232	0.7645	0.087*
C34	0.6474 (4)	-0.01784 (16)	0.85381 (19)	0.0594 (14)
H34A	0.6675	-0.0512	0.8185	0.089*
H34B	0.5620	0.0054	0.8329	0.089*
H34C	0.6339	-0.0380	0.9012	0.089*
N4	0.2179 (4)	0.32703 (16)	0.67898 (19)	0.0504 (10)
C41	0.2608 (4)	0.27407 (16)	0.63794 (19)	0.0593 (15)
H41A	0.2482	0.2339	0.6647	0.089*
H41B	0.3599	0.2791	0.6325	0.089*
H41C	0.2037	0.2727	0.5883	0.089*
C42	0.3149 (6)	0.3291 (3)	0.7528 (3)	0.0522 (13)
H42A	0.3032	0.2897	0.7811	0.078*
H42B	0.2922	0.3666	0.7816	0.078*
H42C	0.4124	0.3322	0.7436	0.078*
C43	0.2388 (6)	0.3897 (3)	0.6357 (3)	0.0512 (13)
H43A	0.3368	0.3929	0.6276	0.077*
H43B	0.2152	0.4269	0.6647	0.077*
H43C	0.1774	0.3890	0.5873	0.077*
C44	0.0667 (6)	0.3221 (3)	0.6915 (4)	0.0560 (14)
H44A	0.0536	0.2824	0.7190	0.084*
H44B	0.0053	0.3214	0.6432	0.084*
H44C	0.0431	0.3593	0.7206	0.084*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mo1	0.0462 (3)	0.0487 (3)	0.0477 (3)	0.00610 (19)	0.00487 (19)	-0.00193 (19)
Mo2	0.0459 (3)	0.0490 (3)	0.0484 (3)	0.00548 (19)	0.00763 (19)	0.00304 (19)
Cu1	0.0491 (4)	0.0505 (4)	0.0476 (4)	0.0071 (3)	0.0065 (3)	-0.0035 (3)
S1	0.0479 (7)	0.0533 (7)	0.0479 (7)	0.0088 (6)	0.0104 (6)	-0.0037 (6)
S2	0.0476 (7)	0.0524 (7)	0.0464 (7)	0.0074 (6)	0.0060 (5)	-0.0030 (6)
S3	0.0467 (7)	0.0504 (7)	0.0484 (7)	0.0073 (5)	0.0092 (6)	0.0045 (6)
S4	0.0545 (8)	0.0472 (7)	0.0531 (7)	0.0107 (6)	0.0103 (6)	0.0061 (6)
S5	0.0503 (8)	0.0580 (8)	0.0526 (7)	0.0073 (6)	0.0124 (6)	0.0045 (6)
S6	0.0538 (8)	0.0518 (8)	0.0567 (8)	0.0112 (6)	0.0069 (6)	-0.0029 (6)
S7	0.0513 (7)	0.0521 (7)	0.0526 (7)	0.0119 (6)	0.0118 (6)	-0.0056 (6)

S8	0.0511 (7)	0.0520 (7)	0.0540 (8)	0.0122 (6)	0.0101 (6)	0.0052 (6)
O1	0.058 (2)	0.069 (3)	0.056 (2)	0.014 (2)	0.0089 (19)	0.022 (2)
N1	0.046 (2)	0.050 (3)	0.055 (3)	-0.0102 (19)	0.011 (2)	0.012 (2)
C11	0.053 (3)	0.053 (3)	0.061 (4)	-0.015 (3)	0.010 (3)	0.015 (3)
C12	0.062 (4)	0.053 (3)	0.068 (4)	-0.014 (3)	0.014 (3)	0.010 (3)
C13	0.063 (3)	0.044 (3)	0.061 (3)	0.009 (3)	-0.006 (3)	-0.015 (3)
N2	0.051 (3)	0.051 (3)	0.051 (3)	-0.005 (2)	0.011 (2)	0.011 (2)
C21	0.060 (4)	0.056 (3)	0.061 (3)	0.017 (3)	-0.015 (3)	0.008 (3)
C22	0.051 (3)	0.067 (4)	0.050 (3)	0.005 (3)	0.017 (2)	0.010 (3)
C23	0.067 (4)	0.051 (3)	0.054 (3)	0.020 (3)	-0.015 (3)	0.008 (3)
C24	0.054 (3)	0.059 (3)	0.055 (3)	0.013 (3)	0.008 (3)	0.008 (3)
N3	0.056 (3)	0.047 (2)	0.054 (3)	-0.004 (2)	0.009 (2)	-0.004 (2)
C31	0.067 (4)	0.049 (3)	0.051 (3)	0.009 (3)	0.011 (3)	0.009 (2)
C32	0.056 (3)	0.063 (4)	0.054 (3)	-0.006 (3)	0.009 (3)	0.015 (3)
C33	0.040 (3)	0.064 (4)	0.070 (4)	-0.005 (2)	0.009 (3)	0.013 (3)
C34	0.059 (4)	0.057 (3)	0.062 (4)	0.004 (3)	0.011 (3)	0.010 (3)
N4	0.049 (3)	0.045 (2)	0.057 (3)	-0.010 (2)	0.006 (2)	0.006 (2)
C41	0.066 (4)	0.063 (4)	0.049 (3)	0.010 (3)	0.009 (3)	0.015 (3)
C42	0.047 (3)	0.057 (3)	0.054 (3)	-0.006 (2)	0.011 (2)	0.011 (3)
C43	0.050 (3)	0.056 (3)	0.050 (3)	-0.015 (2)	0.015 (2)	-0.011 (2)
C44	0.053 (3)	0.051 (3)	0.059 (3)	0.012 (3)	-0.005 (3)	0.009 (3)

Geometric parameters (Å, °)

Mo1—S8	2.1589 (16)	C23—H23C	0.9600
Mo1—S7	2.1710 (17)	C24—H24A	0.9600
Mo1—S1	2.2013 (16)	C24—H24B	0.9600
Mo1—S2	2.2057 (15)	C24—H24C	0.9600
Mo1—Cu1	2.7241 (9)	N3—C33	1.401 (8)
Mo2—S6	2.1451 (17)	N3—C31	1.453 (8)
Mo2—S5	2.1742 (16)	N3—C32	1.549 (9)
Mo2—S4	2.2006 (17)	N3—C34	1.550 (6)
Mo2—S3	2.2148 (16)	C31—H31A	0.9600
Mo2—Cu1	2.7241 (10)	C31—H31B	0.9600
Cu1—S3	2.2549 (17)	C31—H31C	0.9600
Cu1—S1	2.3161 (15)	C32—H32A	0.9600
Cu1—S2	2.3518 (16)	C32—H32B	0.9600
Cu1—S4	2.3630 (17)	C32—H32C	0.9600
O1—C11	1.232 (8)	C33—H33A	0.9600
N1—C11	1.338 (8)	C33—H33B	0.9600
N1—C12	1.434 (9)	C33—H33C	0.9600
N1—C13	1.447 (8)	C34—H34A	0.9600
C11—H11A	0.9300	C34—H34B	0.9600
C12—H12A	0.9600	C34—H34C	0.9600
C12—H12B	0.9600	N4—C41	1.392 (9)
C12—H12C	0.9600	N4—C42	1.481 (7)
C13—H13A	0.9600	N4—C44	1.481 (7)
C13—H13B	0.9600	N4—C43	1.515 (7)
C13—H13C	0.9600	C41—H41A	0.9600

supplementary materials

N2—C24	1.443 (8)	C41—H41B	0.9600
N2—C22	1.473 (8)	C41—H41C	0.9600
N2—C21	1.478 (9)	C42—H42A	0.9600
N2—C23	1.508 (7)	C42—H42B	0.9600
C21—H21A	0.9600	C42—H42C	0.9600
C21—H21B	0.9600	C43—H43A	0.9600
C21—H21C	0.9600	C43—H43B	0.9600
C22—H22A	0.9600	C43—H43C	0.9600
C22—H22B	0.9600	C44—H44A	0.9600
C22—H22C	0.9600	C44—H44B	0.9600
C23—H23A	0.9600	C44—H44C	0.9600
C23—H23B	0.9600		
S8—Mo1—S7	111.79 (6)	H22B—C22—H22C	109.5
S8—Mo1—S1	107.55 (6)	N2—C23—H23A	109.5
S7—Mo1—S1	111.04 (6)	N2—C23—H23B	109.5
S8—Mo1—S2	110.59 (6)	H23A—C23—H23B	109.5
S7—Mo1—S2	108.54 (6)	N2—C23—H23C	109.5
S1—Mo1—S2	107.24 (6)	H23A—C23—H23C	109.5
S8—Mo1—Cu1	139.84 (5)	H23B—C23—H23C	109.5
S7—Mo1—Cu1	108.35 (5)	N2—C24—H24A	109.5
S1—Mo1—Cu1	54.87 (4)	N2—C24—H24B	109.5
S2—Mo1—Cu1	55.80 (4)	H24A—C24—H24B	109.5
S6—Mo2—S5	111.20 (6)	N2—C24—H24C	109.5
S6—Mo2—S4	109.02 (6)	H24A—C24—H24C	109.5
S5—Mo2—S4	108.98 (6)	H24B—C24—H24C	109.5
S6—Mo2—S3	108.88 (6)	C33—N3—C31	109.6 (5)
S5—Mo2—S3	109.52 (6)	C33—N3—C32	112.8 (5)
S4—Mo2—S3	109.21 (6)	C31—N3—C32	109.4 (5)
S6—Mo2—Cu1	126.23 (5)	C33—N3—C34	108.8 (4)
S5—Mo2—Cu1	122.54 (5)	C31—N3—C34	108.6 (4)
S4—Mo2—Cu1	56.14 (4)	C32—N3—C34	107.5 (4)
S3—Mo2—Cu1	53.12 (5)	N3—C31—H31A	109.5
S3—Cu1—S1	117.80 (6)	N3—C31—H31B	109.5
S3—Cu1—S2	115.59 (6)	H31A—C31—H31B	109.5
S1—Cu1—S2	98.94 (6)	N3—C31—H31C	109.5
S3—Cu1—S4	102.39 (6)	H31A—C31—H31C	109.5
S1—Cu1—S4	110.01 (6)	H31B—C31—H31C	109.5
S2—Cu1—S4	112.49 (6)	N3—C32—H32A	109.5
S3—Cu1—Mo2	51.78 (4)	N3—C32—H32B	109.5
S1—Cu1—Mo2	132.00 (5)	H32A—C32—H32B	109.5
S2—Cu1—Mo2	128.60 (4)	N3—C32—H32C	109.5
S4—Cu1—Mo2	50.66 (4)	H32A—C32—H32C	109.5
S3—Cu1—Mo1	147.85 (5)	H32B—C32—H32C	109.5
S1—Cu1—Mo1	51.01 (4)	N3—C33—H33A	109.5
S2—Cu1—Mo1	50.87 (4)	N3—C33—H33B	109.5
S4—Cu1—Mo1	109.76 (5)	H33A—C33—H33B	109.5
Mo2—Cu1—Mo1	160.24 (3)	N3—C33—H33C	109.5
Mo1—S1—Cu1	74.13 (5)	H33A—C33—H33C	109.5
Mo1—S2—Cu1	73.34 (5)	H33B—C33—H33C	109.5

Mo2—S3—Cu1	75.09 (5)	N3—C34—H34A	109.5
Mo2—S4—Cu1	73.20 (5)	N3—C34—H34B	109.5
C11—N1—C12	122.4 (5)	H34A—C34—H34B	109.5
C11—N1—C13	119.2 (5)	N3—C34—H34C	109.5
C12—N1—C13	116.6 (5)	H34A—C34—H34C	109.5
O1—C11—N1	120.3 (5)	H34B—C34—H34C	109.5
O1—C11—H11A	119.9	C41—N4—C42	107.0 (3)
N1—C11—H11A	119.9	C41—N4—C44	112.8 (3)
N1—C12—H12A	109.5	C42—N4—C44	110.6 (4)
N1—C12—H12B	109.5	C41—N4—C43	108.4 (3)
H12A—C12—H12B	109.5	C42—N4—C43	108.3 (4)
N1—C12—H12C	109.5	C44—N4—C43	109.5 (4)
H12A—C12—H12C	109.5	N4—C41—H41A	109.5
H12B—C12—H12C	109.5	N4—C41—H41B	109.5
N1—C13—H13A	109.5	H41A—C41—H41B	109.5
N1—C13—H13B	109.5	N4—C41—H41C	109.5
H13A—C13—H13B	109.5	H41A—C41—H41C	109.5
N1—C13—H13C	109.5	H41B—C41—H41C	109.5
H13A—C13—H13C	109.5	N4—C42—H42A	109.5
H13B—C13—H13C	109.5	N4—C42—H42B	109.5
C24—N2—C22	108.6 (5)	H42A—C42—H42B	109.5
C24—N2—C21	115.1 (5)	N4—C42—H42C	109.5
C22—N2—C21	102.0 (4)	H42A—C42—H42C	109.5
C24—N2—C23	109.7 (5)	H42B—C42—H42C	109.5
C22—N2—C23	106.8 (5)	N4—C43—H43A	109.5
C21—N2—C23	113.9 (5)	N4—C43—H43B	109.5
N2—C21—H21A	109.5	H43A—C43—H43B	109.5
N2—C21—H21B	109.5	N4—C43—H43C	109.5
H21A—C21—H21B	109.5	H43A—C43—H43C	109.5
N2—C21—H21C	109.5	H43B—C43—H43C	109.5
H21A—C21—H21C	109.5	N4—C44—H44A	109.5
H21B—C21—H21C	109.5	N4—C44—H44B	109.5
N2—C22—H22A	109.5	H44A—C44—H44B	109.5
N2—C22—H22B	109.5	N4—C44—H44C	109.5
H22A—C22—H22B	109.5	H44A—C44—H44C	109.5
N2—C22—H22C	109.5	H44B—C44—H44C	109.5
H22A—C22—H22C	109.5		
S6—Mo2—Cu1—S3	87.24 (8)	S7—Mo1—Cu1—Mo2	3.80 (11)
S5—Mo2—Cu1—S3	-90.96 (7)	S1—Mo1—Cu1—Mo2	107.05 (11)
S4—Mo2—Cu1—S3	176.95 (6)	S2—Mo1—Cu1—Mo2	-96.54 (10)
S6—Mo2—Cu1—S1	-7.91 (9)	S8—Mo1—S1—Cu1	-139.25 (6)
S5—Mo2—Cu1—S1	173.89 (8)	S7—Mo1—S1—Cu1	98.15 (6)
S4—Mo2—Cu1—S1	81.80 (8)	S2—Mo1—S1—Cu1	-20.28 (6)
S3—Mo2—Cu1—S1	-95.15 (8)	S3—Cu1—S1—Mo1	143.57 (6)
S6—Mo2—Cu1—S2	-178.47 (8)	S2—Cu1—S1—Mo1	18.32 (6)
S5—Mo2—Cu1—S2	3.34 (9)	S4—Cu1—S1—Mo1	-99.67 (6)
S4—Mo2—Cu1—S2	-88.76 (8)	Mo2—Cu1—S1—Mo1	-154.22 (5)
S3—Mo2—Cu1—S2	94.29 (8)	S8—Mo1—S2—Cu1	137.04 (6)
S6—Mo2—Cu1—S4	-89.71 (8)	S7—Mo1—S2—Cu1	-99.99 (6)

supplementary materials

S5—Mo2—Cu1—S4	92.10 (7)	S1—Mo1—S2—Cu1	20.04 (6)
S3—Mo2—Cu1—S4	-176.95 (6)	S3—Cu1—S2—Mo1	-145.14 (5)
S6—Mo2—Cu1—Mo1	-98.05 (11)	S1—Cu1—S2—Mo1	-18.36 (6)
S5—Mo2—Cu1—Mo1	83.75 (11)	S4—Cu1—S2—Mo1	97.75 (6)
S4—Mo2—Cu1—Mo1	-8.35 (10)	Mo2—Cu1—S2—Mo1	154.55 (4)
S3—Mo2—Cu1—Mo1	174.71 (11)	S6—Mo2—S3—Cu1	-121.63 (6)
S8—Mo1—Cu1—S3	-5.97 (12)	S5—Mo2—S3—Cu1	116.59 (6)
S7—Mo1—Cu1—S3	175.98 (9)	S4—Mo2—S3—Cu1	-2.69 (6)
S1—Mo1—Cu1—S3	-80.77 (10)	S1—Cu1—S3—Mo2	123.21 (6)
S2—Mo1—Cu1—S3	75.63 (10)	S2—Cu1—S3—Mo2	-120.22 (5)
S8—Mo1—Cu1—S1	74.81 (8)	S4—Cu1—S3—Mo2	2.42 (5)
S7—Mo1—Cu1—S1	-103.25 (7)	Mo1—Cu1—S3—Mo2	-176.64 (7)
S2—Mo1—Cu1—S1	156.40 (7)	S6—Mo2—S4—Cu1	121.44 (5)
S8—Mo1—Cu1—S2	-81.60 (8)	S5—Mo2—S4—Cu1	-117.02 (5)
S7—Mo1—Cu1—S2	100.35 (7)	S3—Mo2—S4—Cu1	2.59 (5)
S1—Mo1—Cu1—S2	-156.40 (7)	S3—Cu1—S4—Mo2	-2.46 (5)
S8—Mo1—Cu1—S4	175.01 (7)	S1—Cu1—S4—Mo2	-128.49 (5)
S7—Mo1—Cu1—S4	-3.04 (6)	S2—Cu1—S4—Mo2	122.26 (6)
S1—Mo1—Cu1—S4	100.20 (7)	Mo1—Cu1—S4—Mo2	177.01 (3)
S2—Mo1—Cu1—S4	-103.39 (6)	C12—N1—C11—O1	177.0 (6)
S8—Mo1—Cu1—Mo2	-178.14 (9)	C13—N1—C11—O1	13.0 (9)

Fig. 1

