

catena-Poly[[[N'-(4-cyanobenzylidene)-nicotinohydrazide]silver(I)]- μ -N'-(4-cyanobenzylidene)nicotinohydrazide] hexafluoridophosphate]

Xin-Sheng Wan, Yu-Li Dang, Chun-Hong Kou, Zhan-Fang Zhou and Cao-Yuan Niu*

College of Sciences, Henan Agricultural University, Zhengzhou 450002, People's Republic of China

Correspondence e-mail: niu_cy2000@yahoo.com.cn


Received 11 October 2008; accepted 20 November 2008

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; disorder in solvent or counterion; R factor = 0.037; wR factor = 0.099; data-to-parameter ratio = 14.8.

In the title polymer, $\{[\text{Ag}(\text{C}_{14}\text{H}_{10}\text{N}_4\text{O})_2]\text{PF}_6\}_n$, each Ag^{I} ion is coordinated by two N atoms from two pyridyl rings of independent N'-(4-cyanobenzylidene)nicotinohydrazide ligands, and one N atom from one carbonitrile group of a symmetry-related ligand in a distorted T-shaped geometry. The ligands exhibit two modes of coordination. One acts as a bridge connecting Ag atoms to form one-dimensional chains along $[\bar{1}01]$. The other acts as a terminal monodentate ligand, coordinating to Ag through its pyridyl N atom. Two neighbouring antiparallel chains in the crystal are connected through $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds. Other adjacent chains are packed via $\text{Ag}\cdots\text{O}$ interactions, with $\text{Ag}\cdots\text{O}$ separations of 2.876 (2) Å. In addition, PF_6^- counter-anions interact with the hydrazone groups through $\text{N}-\text{H}\cdots\text{F}$ hydrogen bonds. The PF_6^- anion is disordered over two sites, with occupancies of 0.773 (8) and 0.227 (8).

Related literature

For background on fluorescent silver coordination complexes, see: Dong *et al.* (2004); Sumbly & Hardie (2005). For related structures, see: Niu *et al.* (2007, 2008); Vatsadze *et al.* (2004); Zheng *et al.* (2003).



Experimental

Crystal data

$[\text{Ag}(\text{C}_{14}\text{H}_{10}\text{N}_4\text{O})_2]\text{PF}_6$
 $M_r = 753.36$
 Monoclinic, $C2/c$
 $a = 22.3252$ (17) Å
 $b = 13.6939$ (11) Å
 $c = 19.8523$ (16) Å
 $\beta = 99.9770$ (10)°

$V = 5977.4$ (8) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.81$ mm⁻¹
 $T = 173$ (2) K
 $0.44 \times 0.32 \times 0.29$ mm

Data collection

Siemens SMART CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Siemens, 1996)
 $T_{\text{min}} = 0.718$, $T_{\text{max}} = 0.799$

19020 measured reflections
 6823 independent reflections
 5105 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.099$
 $S = 1.03$
 6823 reflections
 460 parameters
 96 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.92$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.67$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Ag1—N1	2.172 (2)	Ag1—N8 ⁱ	2.456 (3)
Ag1—N2	2.199 (2)		
N1—Ag1—N2	156.22 (8)	N2—Ag1—N8 ⁱ	92.22 (9)
N1—Ag1—N8 ⁱ	109.53 (9)		

Symmetry code: (i) $x - \frac{1}{2}, -y + \frac{3}{2}, z + \frac{1}{2}$.

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N3—H28 ⁱⁱ ···O2 ⁱⁱ	0.865 (18)	2.15 (2)	2.990 (3)	162 (3)
N6—H29···F5	0.853 (18)	2.21 (2)	3.001 (4)	155 (3)

Symmetry code: (ii) $-x + \frac{1}{2}, -y + \frac{3}{2}, -z$.

Data collection: SMART (Siemens, 1996); cell refinement: SAINT (Siemens, 1996); data reduction: SAINT; program(s) used to solve structure: SHELXL97 (Sheldrick, 2008); program(s) used to refine

structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXL97* and *DIAMOND* (Brandenburg, 2005); software used to prepare material for publication: *SHELXL97*.

We are grateful to Mrs Li (Wuhan University) for her assistance with the X-ray crystallographic analysis. We also gratefully acknowledge financial support from the Natural Science Foundation of Henan Province (2008B150008) and the Science and Technology Key Task of Henan Province (0624040011).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2203).

References

- Brandenburg, K. (2005). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Dong, Y.-B., Zhao, X., Huang, R.-Q., Smith, M. D. & zur Loye, H.-C. (2004). *Inorg. Chem.* **43**, 5603–5612.
- Niu, C.-Y., Wu, B.-L., Zheng, X.-F., Zhang, H.-Y., Li, Z.-J. & Hou, H.-W. (2007). *Dalton Trans.* pp. 5710–5713.
- Niu, C.-Y., Zheng, X.-F., Bai, L.-L., Wu, X.-L. & Kou, C.-H. (2008). *Acta Cryst. C* **64**, m305–m307.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Siemens (1996). *SMART, SAINT and SADABS*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Sumby, C. J. & Hardie, M. J. (2005). *Angew. Chem. Int. Ed.* **44**, 6395–6399.
- Vatsadze, S. Z., Kovalkina, M. A., Sviridenkova, N. V., Zyk, N. V., Churakov, A. V., Kuz'mina, L. G. & Howard, J. A. K. (2004). *CrystEngComm*, **6**, 112–115.
- Zheng, Y., Du, M., Li, J.-R., Zhang, R.-H. & Bu, X.-H. (2003). *Dalton Trans.* pp. 1509–1514.

supplementary materials

Acta Cryst. (2008). E64, m1618-m1619 [doi:10.1107/S1600536808038907]

***catena*-Poly[[[*N'*-(4-cyanobenzylidene)nicotinohydrazide]silver(I)]- μ -*N'*-(4-cyanobenzylidene)nicotinohydrazide] hexafluoridophosphate]**

X.-S. Wan, Y.-L. Dang, C.-H. Kou, Z.-F. Zhou and C.-Y. Niu

Comment

Pyridyl organic ligands with carbonitrile groups can be used to construct silver coordination complexes with fascinating structures and good fluorescent properties (Sumbly & Hardie, 2005; Dong *et al.*, 2004). We also synthesized one-dimensional and two-dimensional silver coordination polymers using this kind of ligands (Niu *et al.*, 2007, 2008). Herein, a one-dimensional silver coordination polymer constructed with a new bridging ligand of this type, 4-cyanobenzylidene nicotinohydrazide, is reported.

In the title compound, (I), the central Ag^I ion is coordinated by two N atoms from two pyridyl rings of two different ligands (N1, N2) and one N atom from one carbonitrile group of another ligand [N8ⁱ, symmetry code: (i) $x - 1/2, -y + 3/2, z + 1/2$], forming a slightly distorted T-shaped coordination environment (Fig. 1). The N1—Ag1—N2 bond angle is 156.22 (8)°, indicating these three atoms are not exactly in one line. Bond angles N1—Ag1—N8ⁱ and N2—Ag1—N8ⁱ are larger than 90° (Table 1). The Ag—N bond distances for pyridyl rings are in the range 2.172 (2)–2.199 (2) Å, which is smaller than N—Ag bond distance for the carbonitrile group (Table 1).

One 4-cyanobenzylidene nicotinohydrazide molecule acts as a μ_2 -bridging ligand, by coordinating pyridyl and carbonitrile N atoms. Each bridging ligand connects two silver atoms together by one pyridyl N atom (N1) and one carbonitrile N atom (N8) to form a one-dimensional chain along the [-1 0 1] direction. The separation between two neighbouring Ag atoms in one chain is about 16 Å. Meanwhile, the other independent ligand is acting as a terminal ligand, being coordinated to Ag only through a pyridyl N atom. Two terminal ligands connected to two adjacent Ag atoms in one chain are located at the opposite positions away from the chain (Fig. 2).

There are hydrogen bonds between uncoordinating groups, including pyridyl rings of terminal ligands and all hydrazone groups, as well as other groups like counteranions. On one hand, counteranions PF₆⁻ interact with the ligands in the polymer through N—H...F hydrogen bonds (Table 2). Four F atoms (F1 to F4) of the PF₆⁻ anion are disordered over two sites, with occupancies 0.773 (8) and 0.227 (8). On the other hand, there are also N—H...O hydrogen bonds between two neighbouring antiparallel chains in the crystal (Fig. 3). In addition to these intermolecular contacts, there are weak Ag...O interactions between one O atom (O1) of the terminal ligand and one Ag atom in the neighbouring chain, with Ag...O separations of 2.876 (2) Å. These noncovalent interactions have large contributions to the supramolecular three-dimensional framework.

One-dimensional Ag^I coordination polymers with T-shaped coordination geometry were previously described in a few compounds: {[Ag(2,6-di(3-pyridylmethylidene) cyclohexanone)](NO₃)}_n (Vatsadze *et al.*, 2004) and {[Ag(2,2'-(methylenebis(thio)) bis(pyrimidine))](NO₃)}_n (Zheng *et al.*, 2003). The N—Ag—N bond angles in these two compounds deviate from 180° (*ca.* 158 and 131°, respectively), which are close to that observed in (I).

Experimental

A solution of AgPF_6 (0.025 g, 0.1 mmol) in CH_3OH (10 ml) was carefully layered on a $\text{CH}_3\text{OH}/\text{CHCl}_3$ solution (5 ml/10 ml) of 4-cyanobenzylidene nicotinohydrazide (0.025 g, 0.1 mmol) in a straight glass tube. About ten days later, colourless single crystals suitable for X-ray analysis were obtained (yield: *ca.* 50%). Analysis, calculated for $\text{C}_{28}\text{H}_{20}\text{AgN}_8\text{O}_2\text{F}_6\text{P}$: C 44.64, H 2.68, N 14.87%; found: C 44.79, H 2.69, N 14.99%.

Refinement

C-bound H atoms were placed in calculated positions and refined using a riding model [$\text{C}-\text{H} = 0.95 \text{ \AA}$ and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$]. The N-bound H atoms were first introduced in calculated positions, and then their positions and displacement parameters were refined with the N—H bond lengths restrained to $0.88 (2) \text{ \AA}$. Four F atoms (F1/F2/F3/F4) of the hexafluorophosphate anion are disordered over two positions (F1'/F2'/F3'/F4'), and all P—F bond lengths were restrained to a target value of $1.58 (2) \text{ \AA}$. Displacement parameters for disordered F atoms were also subjected to restraints. The final difference map had a highest peak at 0.88 \AA from Ag1 and a deepest hole at 0.71 \AA from Ag1, but was otherwise featureless.

Figures




Fig. 1. A view of the Ag^{I} coordination environment in the polymeric structure of (I), showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level and H atoms are shown as small spheres of arbitrary radii [Symmetry code: (i) $x - 1/2, -y + 3/2, z + 1/2$].



Fig. 2. A ball and stick diagram showing the one-dimensional chain. All counteranions and H atoms have been omitted for clarity.



Fig. 3. A diagram showing the intermolecular hydrogen bonds, indicated by dashed lines.

catena-Poly[[[N^I-(4-cyanobenzylidene)nicotinohydrazide)silver(I)]- μ -N^I-(4-cyanobenzylidene)nicotinohydrazide] hexafluorophosphate]

Crystal data

$[\text{Ag}(\text{C}_{14}\text{H}_{10}\text{N}_4\text{O})_2]\text{PF}_6$

$M_r = 753.36$

$F_{000} = 3008$

$D_x = 1.674 \text{ Mg m}^{-3}$

Monoclinic, $C2/c$
Hall symbol: $-C\ 2yc$
 $a = 22.3252\ (17)\ \text{\AA}$
 $b = 13.6939\ (11)\ \text{\AA}$
 $c = 19.8523\ (16)\ \text{\AA}$
 $\beta = 99.9770\ (10)^\circ$
 $V = 5977.4\ (8)\ \text{\AA}^3$
 $Z = 8$

Mo $K\alpha$ radiation
 $\lambda = 0.71073\ \text{\AA}$
Cell parameters from 5745 reflections
 $\theta = 2.1\text{--}27.5^\circ$
 $\mu = 0.81\ \text{mm}^{-1}$
 $T = 173\ (2)\ \text{K}$
Prism, colourless
 $0.44 \times 0.32 \times 0.29\ \text{mm}$

Data collection

Siemens SMART CCD area-detector diffractometer
Radiation source: fine-focus sealed tube
Monochromator: graphite
 $T = 173(2)\ \text{K}$
 φ and ω scans
Absorption correction: multi-scan (SADABS; Siemens, 1996)
 $T_{\min} = 0.718$, $T_{\max} = 0.800$
19020 measured reflections

6823 independent reflections
5105 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$
 $\theta_{\max} = 27.5^\circ$
 $\theta_{\min} = 2.1^\circ$
 $h = -28 \rightarrow 25$
 $k = -16 \rightarrow 17$
 $l = -25 \rightarrow 25$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.099$
 $S = 1.03$
6823 reflections
460 parameters
96 restraints
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0457P)^2 + 5.9377P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.92\ \text{e}\ \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.67\ \text{e}\ \text{\AA}^{-3}$
Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
N8	0.61351 (13)	0.7668 (2)	-0.23514 (14)	0.0680 (8)	
C28	0.57801 (14)	0.7655 (3)	-0.20064 (15)	0.0538 (7)	
C25	0.53188 (13)	0.7592 (2)	-0.15834 (14)	0.0478 (7)	
C24	0.50609 (14)	0.6686 (2)	-0.14987 (15)	0.0531 (7)	
H24	0.5184	0.6128	-0.1724	0.064*	
C26	0.51433 (15)	0.8411 (3)	-0.12646 (17)	0.0601 (8)	

supplementary materials

H26	0.5315	0.9031	-0.1330	0.072*
C27	0.47114 (15)	0.8310 (2)	-0.08462 (18)	0.0601 (8)
H27	0.4592	0.8868	-0.0618	0.072*
C23	0.46282 (14)	0.6602 (2)	-0.10881 (15)	0.0510 (7)
H23	0.4449	0.5985	-0.1032	0.061*
C22	0.44513 (13)	0.7418 (2)	-0.07544 (14)	0.0477 (7)
C21	0.40138 (14)	0.7334 (2)	-0.02860 (15)	0.0520 (7)
H21	0.3960	0.7864	0.0006	0.062*
N7	0.37061 (11)	0.6556 (2)	-0.02684 (12)	0.0494 (6)
N6	0.33253 (11)	0.6554 (2)	0.02109 (12)	0.0486 (6)
Ag1	0.182725 (12)	0.812121 (16)	0.197931 (13)	0.05723 (10)
C12	0.44862 (16)	1.2299 (3)	-0.09411 (17)	0.0643 (9)
H12	0.4699	1.1967	-0.1248	0.077*
C19	0.24057 (13)	0.66761 (18)	0.10608 (13)	0.0401 (6)
H19	0.2509	0.7270	0.0862	0.048*
C15	0.19185 (13)	0.5876 (2)	0.18284 (14)	0.0429 (6)
H15	0.1673	0.5897	0.2174	0.052*
C6	0.24333 (13)	1.19804 (19)	0.14358 (14)	0.0416 (6)
C18	0.26109 (12)	0.58116 (18)	0.08291 (12)	0.0367 (5)
C9	0.38638 (14)	1.3271 (2)	-0.00520 (16)	0.0532 (7)
H9	0.3651	1.3608	0.0253	0.064*
C7	0.33364 (14)	1.1743 (2)	0.01789 (15)	0.0502 (7)
H7	0.3256	1.1073	0.0076	0.060*
C4	0.20356 (12)	1.13009 (19)	0.17502 (13)	0.0380 (6)
C17	0.24587 (13)	0.49446 (19)	0.11196 (14)	0.0449 (6)
H17	0.2594	0.4336	0.0971	0.054*
C16	0.21082 (14)	0.4979 (2)	0.16270 (14)	0.0486 (7)
H16	0.1999	0.4396	0.1834	0.058*
C1	0.13502 (14)	1.0108 (2)	0.24051 (14)	0.0507 (7)
H1	0.1108	0.9695	0.2634	0.061*
C3	0.16615 (14)	1.1703 (2)	0.21661 (15)	0.0482 (7)
H3	0.1647	1.2390	0.2226	0.058*
C5	0.20473 (13)	1.02969 (19)	0.16908 (13)	0.0410 (6)
H5	0.2304	1.0018	0.1408	0.049*
C13	0.40755 (16)	1.1802 (2)	-0.06192 (17)	0.0606 (8)
H13	0.4006	1.1126	-0.0708	0.073*
C11	0.45847 (15)	1.3286 (3)	-0.08112 (17)	0.0575 (8)
C8	0.37610 (13)	1.2281 (2)	-0.01660 (14)	0.0485 (7)
C14	0.50138 (17)	1.3823 (3)	-0.1134 (2)	0.0713 (10)
C2	0.13130 (14)	1.1097 (2)	0.24891 (16)	0.0535 (8)
H2	0.1048	1.1361	0.2769	0.064*
C10	0.42664 (15)	1.3771 (2)	-0.03712 (17)	0.0576 (8)
H10	0.4328	1.4450	-0.0291	0.069*
N3	0.26781 (11)	1.16107 (18)	0.09093 (12)	0.0452 (5)
N2	0.20658 (11)	0.67179 (15)	0.15560 (11)	0.0417 (5)
N4	0.30732 (11)	1.21732 (18)	0.06194 (12)	0.0474 (6)
N1	0.17155 (11)	0.96955 (16)	0.20120 (11)	0.0443 (5)
N5	0.53463 (17)	1.4260 (3)	-0.1381 (2)	0.1005 (13)
P1	0.37466 (4)	0.89969 (6)	0.14663 (5)	0.0599 (2)

O2	0.29542 (10)	0.50513 (15)	-0.01052 (10)	0.0563 (5)	
O1	0.25284 (10)	1.28063 (15)	0.16533 (11)	0.0584 (6)	
C20	0.29779 (12)	0.57604 (19)	0.02673 (13)	0.0409 (6)	
F5	0.38656 (13)	0.78661 (16)	0.13688 (14)	0.1030 (8)	
F6	0.35972 (14)	1.01041 (16)	0.15169 (18)	0.1254 (11)	
F1	0.3229 (3)	0.8764 (3)	0.1891 (3)	0.1111 (19)	0.773 (8)
F2	0.4206 (3)	0.9026 (5)	0.2114 (4)	0.182 (3)	0.773 (8)
F3	0.4237 (2)	0.9205 (3)	0.1009 (4)	0.129 (2)	0.773 (8)
F4	0.3249 (2)	0.8894 (4)	0.0790 (2)	0.1269 (19)	0.773 (8)
F1'	0.3816 (8)	0.8806 (8)	0.2243 (5)	0.096 (5)	0.227 (8)
F2'	0.4440 (4)	0.9236 (7)	0.1569 (9)	0.082 (4)	0.227 (8)
F3'	0.3702 (9)	0.9217 (9)	0.0687 (5)	0.112 (5)	0.227 (8)
F4'	0.3055 (4)	0.8774 (9)	0.1384 (11)	0.109 (5)	0.227 (8)
H28	0.2574 (13)	1.1069 (16)	0.0696 (14)	0.050 (9)*	
H29	0.3373 (14)	0.7020 (18)	0.0501 (13)	0.048 (9)*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N8	0.0642 (18)	0.082 (2)	0.0669 (17)	-0.0083 (15)	0.0370 (15)	-0.0052 (15)
C28	0.0488 (17)	0.067 (2)	0.0496 (16)	-0.0054 (15)	0.0193 (14)	-0.0023 (15)
C25	0.0400 (15)	0.0643 (19)	0.0425 (15)	-0.0035 (13)	0.0168 (12)	-0.0011 (13)
C24	0.0556 (18)	0.0592 (19)	0.0508 (16)	0.0032 (14)	0.0264 (14)	0.0007 (14)
C26	0.060 (2)	0.0591 (19)	0.068 (2)	-0.0163 (15)	0.0323 (17)	-0.0065 (16)
C27	0.061 (2)	0.057 (2)	0.070 (2)	-0.0081 (15)	0.0337 (17)	-0.0143 (16)
C23	0.0533 (18)	0.0538 (17)	0.0513 (16)	-0.0046 (13)	0.0241 (14)	0.0040 (13)
C22	0.0434 (16)	0.0615 (19)	0.0422 (14)	-0.0043 (13)	0.0185 (13)	0.0001 (13)
C21	0.0528 (18)	0.0588 (19)	0.0508 (16)	-0.0022 (14)	0.0266 (14)	-0.0057 (14)
N7	0.0492 (14)	0.0615 (15)	0.0438 (13)	-0.0014 (12)	0.0254 (11)	-0.0007 (11)
N6	0.0535 (15)	0.0522 (15)	0.0478 (13)	-0.0051 (11)	0.0304 (12)	-0.0071 (11)
Ag1	0.07322 (19)	0.03326 (13)	0.07451 (18)	0.00004 (10)	0.03887 (14)	-0.00591 (10)
C12	0.064 (2)	0.075 (2)	0.062 (2)	0.0135 (18)	0.0322 (17)	0.0016 (18)
C19	0.0487 (16)	0.0332 (13)	0.0431 (14)	-0.0006 (11)	0.0210 (12)	0.0034 (11)
C15	0.0494 (16)	0.0379 (14)	0.0468 (14)	-0.0020 (11)	0.0230 (13)	-0.0003 (11)
C6	0.0461 (15)	0.0332 (14)	0.0469 (15)	0.0052 (11)	0.0122 (12)	0.0029 (11)
C18	0.0411 (14)	0.0351 (13)	0.0366 (13)	0.0005 (10)	0.0145 (11)	-0.0020 (10)
C9	0.0540 (18)	0.0563 (19)	0.0534 (17)	0.0099 (14)	0.0210 (14)	0.0056 (14)
C7	0.0539 (18)	0.0480 (17)	0.0508 (16)	-0.0006 (13)	0.0151 (14)	0.0028 (13)
C4	0.0431 (15)	0.0345 (13)	0.0377 (13)	0.0032 (11)	0.0107 (11)	-0.0017 (10)
C17	0.0579 (18)	0.0308 (13)	0.0502 (15)	0.0018 (12)	0.0210 (13)	-0.0012 (11)
C16	0.0652 (19)	0.0339 (14)	0.0519 (16)	-0.0052 (13)	0.0246 (14)	0.0049 (12)
C1	0.0609 (19)	0.0464 (16)	0.0517 (16)	-0.0040 (13)	0.0294 (15)	-0.0037 (13)
C3	0.0550 (17)	0.0375 (15)	0.0556 (16)	0.0041 (12)	0.0194 (14)	-0.0086 (12)
C5	0.0520 (16)	0.0349 (14)	0.0396 (13)	0.0029 (11)	0.0176 (12)	-0.0028 (11)
C13	0.069 (2)	0.054 (2)	0.063 (2)	0.0052 (15)	0.0251 (17)	-0.0041 (15)
C11	0.0524 (18)	0.067 (2)	0.0578 (18)	0.0096 (15)	0.0222 (15)	0.0139 (16)
C8	0.0455 (16)	0.0573 (18)	0.0444 (15)	0.0042 (13)	0.0123 (13)	0.0069 (13)
C14	0.072 (2)	0.070 (2)	0.082 (2)	0.0147 (18)	0.041 (2)	0.0162 (19)

supplementary materials

C2	0.0587 (19)	0.0492 (17)	0.0604 (18)	0.0035 (14)	0.0321 (15)	-0.0082 (14)
C10	0.0587 (19)	0.0539 (19)	0.0648 (19)	0.0017 (14)	0.0238 (16)	0.0081 (15)
N3	0.0534 (15)	0.0370 (13)	0.0491 (13)	-0.0059 (10)	0.0197 (11)	-0.0014 (10)
N2	0.0493 (13)	0.0349 (12)	0.0466 (12)	0.0005 (9)	0.0247 (11)	0.0001 (9)
N4	0.0512 (14)	0.0456 (13)	0.0482 (13)	-0.0036 (11)	0.0163 (11)	0.0063 (11)
N1	0.0579 (15)	0.0331 (12)	0.0463 (12)	-0.0004 (10)	0.0215 (11)	-0.0037 (9)
N5	0.105 (3)	0.084 (3)	0.134 (3)	0.009 (2)	0.083 (3)	0.021 (2)
P1	0.0625 (5)	0.0379 (4)	0.0841 (6)	0.0022 (4)	0.0261 (5)	-0.0005 (4)
O2	0.0716 (14)	0.0488 (12)	0.0551 (12)	0.0005 (10)	0.0296 (11)	-0.0130 (10)
O1	0.0766 (15)	0.0328 (10)	0.0723 (14)	-0.0048 (10)	0.0310 (12)	-0.0021 (10)
C20	0.0463 (15)	0.0397 (14)	0.0403 (14)	0.0051 (11)	0.0179 (12)	0.0000 (11)
F5	0.139 (2)	0.0483 (12)	0.125 (2)	0.0164 (13)	0.0331 (17)	-0.0072 (13)
F6	0.142 (2)	0.0447 (13)	0.208 (3)	0.0124 (13)	0.083 (2)	0.0042 (16)
F1	0.141 (4)	0.087 (2)	0.128 (4)	0.010 (2)	0.085 (3)	0.026 (2)
F2	0.143 (5)	0.219 (6)	0.156 (5)	0.022 (4)	-0.051 (4)	-0.062 (4)
F3	0.112 (4)	0.094 (3)	0.207 (6)	-0.018 (2)	0.105 (4)	-0.012 (3)
F4	0.110 (3)	0.157 (4)	0.101 (3)	0.006 (3)	-0.017 (3)	0.017 (3)
F1'	0.149 (10)	0.085 (7)	0.064 (6)	-0.010 (7)	0.041 (7)	0.010 (5)
F2'	0.056 (5)	0.065 (6)	0.125 (9)	-0.004 (4)	0.020 (6)	-0.001 (6)
F3'	0.146 (11)	0.099 (8)	0.085 (7)	0.012 (8)	0.000 (7)	0.021 (6)
F4'	0.075 (7)	0.097 (8)	0.156 (11)	-0.012 (5)	0.020 (7)	-0.009 (8)

Geometric parameters (\AA , $^\circ$)

N8—C28	1.134 (4)	C9—C8	1.387 (4)
N8—Ag1 ⁱ	2.456 (3)	C9—H9	0.9500
C28—C25	1.440 (4)	C7—N4	1.279 (4)
C25—C26	1.378 (4)	C7—C8	1.461 (4)
C25—C24	1.389 (4)	C7—H7	0.9500
C24—C23	1.372 (4)	C4—C5	1.381 (4)
C24—H24	0.9500	C4—C3	1.386 (4)
C26—C27	1.384 (4)	C17—C16	1.379 (4)
C26—H26	0.9500	C17—H17	0.9500
C27—C22	1.379 (4)	C16—H16	0.9500
C27—H27	0.9500	C1—N1	1.347 (3)
C23—C22	1.390 (4)	C1—C2	1.369 (4)
C23—H23	0.9500	C1—H1	0.9500
C22—C21	1.465 (4)	C3—C2	1.370 (4)
C21—N7	1.271 (4)	C3—H3	0.9500
C21—H21	0.9500	C5—N1	1.341 (3)
N7—N6	1.382 (3)	C5—H5	0.9500
N6—C20	1.351 (4)	C13—C8	1.397 (4)
N6—H29	0.853 (18)	C13—H13	0.9500
Ag1—N1	2.172 (2)	C11—C10	1.387 (4)
Ag1—N2	2.199 (2)	C11—C14	1.443 (5)
Ag1—N8 ⁱⁱ	2.456 (3)	C14—N5	1.130 (4)
C12—C13	1.384 (5)	C2—H2	0.9500
C12—C11	1.387 (5)	C10—H10	0.9500
C12—H12	0.9500	N3—N4	1.371 (3)

C19—N2	1.343 (3)	N3—H28	0.865 (18)
C19—C18	1.377 (3)	P1—F2	1.500 (5)
C19—H19	0.9500	P1—F1'	1.545 (9)
C15—N2	1.338 (3)	P1—F4'	1.555 (10)
C15—C16	1.381 (4)	P1—F6	1.559 (2)
C15—H15	0.9500	P1—F2'	1.560 (8)
C6—O1	1.216 (3)	P1—F3'	1.562 (9)
C6—N3	1.358 (3)	P1—F3	1.564 (4)
C6—C4	1.495 (4)	P1—F1	1.577 (3)
C18—C17	1.388 (4)	P1—F5	1.589 (2)
C18—C20	1.496 (3)	P1—F4	1.594 (4)
C9—C10	1.369 (4)	O2—C20	1.216 (3)
C28—N8—Ag1 ⁱ	153.0 (3)	C12—C13—H13	119.6
N8—C28—C25	177.1 (4)	C8—C13—H13	119.6
C26—C25—C24	120.9 (3)	C10—C11—C12	120.1 (3)
C26—C25—C28	120.5 (3)	C10—C11—C14	119.2 (3)
C24—C25—C28	118.5 (3)	C12—C11—C14	120.7 (3)
C23—C24—C25	119.7 (3)	C9—C8—C13	118.5 (3)
C23—C24—H24	120.1	C9—C8—C7	121.3 (3)
C25—C24—H24	120.1	C13—C8—C7	120.3 (3)
C25—C26—C27	118.5 (3)	N5—C14—C11	178.6 (4)
C25—C26—H26	120.7	C1—C2—C3	119.4 (3)
C27—C26—H26	120.7	C1—C2—H2	120.3
C22—C27—C26	121.4 (3)	C3—C2—H2	120.3
C22—C27—H27	119.3	C9—C10—C11	120.1 (3)
C26—C27—H27	119.3	C9—C10—H10	119.9
C24—C23—C22	120.2 (3)	C11—C10—H10	119.9
C24—C23—H23	119.9	C6—N3—N4	119.2 (2)
C22—C23—H23	119.9	C6—N3—H28	126 (2)
C27—C22—C23	119.2 (3)	N4—N3—H28	114 (2)
C27—C22—C21	119.7 (3)	C15—N2—C19	117.9 (2)
C23—C22—C21	121.0 (3)	C15—N2—Ag1	120.36 (17)
N7—C21—C22	120.4 (3)	C19—N2—Ag1	121.45 (17)
N7—C21—H21	119.8	C7—N4—N3	115.7 (2)
C22—C21—H21	119.8	C5—N1—C1	117.3 (2)
C21—N7—N6	114.8 (3)	C5—N1—Ag1	121.31 (17)
C20—N6—N7	119.3 (2)	C1—N1—Ag1	121.18 (18)
C20—N6—H29	123 (2)	F2—P1—F4'	127.9 (7)
N7—N6—H29	116 (2)	F1'—P1—F4'	89.8 (7)
N1—Ag1—N2	156.22 (8)	F2—P1—F6	92.4 (3)
N1—Ag1—N8 ⁱⁱ	109.53 (9)	F1'—P1—F6	94.9 (5)
N2—Ag1—N8 ⁱⁱ	92.22 (9)	F4'—P1—F6	88.7 (5)
C13—C12—C11	119.3 (3)	F2—P1—F2'	50.8 (5)
C13—C12—H12	120.3	F1'—P1—F2'	88.9 (7)
C11—C12—H12	120.3	F4'—P1—F2'	178.3 (8)
N2—C19—C18	123.0 (2)	F6—P1—F2'	90.3 (4)
N2—C19—H19	118.5	F2—P1—F3'	139.5 (7)
C18—C19—H19	118.5	F1'—P1—F3'	177.5 (8)

supplementary materials

N2—C15—C16	122.7 (2)	F4'—P1—F3'	92.5 (8)
N2—C15—H15	118.7	F6—P1—F3'	84.2 (4)
C16—C15—H15	118.7	F2'—P1—F3'	88.8 (7)
O1—C6—N3	123.4 (3)	F2—P1—F3	92.8 (4)
O1—C6—C4	120.7 (2)	F1'—P1—F3	130.6 (6)
N3—C6—C4	115.9 (2)	F4'—P1—F3	139.2 (7)
C19—C18—C17	118.5 (2)	F6—P1—F3	92.24 (19)
C19—C18—C20	123.1 (2)	F3'—P1—F3	47.2 (6)
C17—C18—C20	118.4 (2)	F2—P1—F1	90.2 (4)
C10—C9—C8	121.1 (3)	F1'—P1—F1	52.3 (6)
C10—C9—H9	119.5	F6—P1—F1	88.72 (18)
C8—C9—H9	119.5	F2'—P1—F1	140.8 (6)
N4—C7—C8	120.0 (3)	F3'—P1—F1	129.9 (7)
N4—C7—H7	120.0	F3—P1—F1	176.8 (3)
C8—C7—H7	120.0	F2—P1—F5	91.5 (3)
C5—C4—C3	117.9 (2)	F1'—P1—F5	88.1 (4)
C5—C4—C6	124.2 (2)	F4'—P1—F5	88.8 (5)
C3—C4—C6	117.7 (2)	F6—P1—F5	176.14 (19)
C16—C17—C18	119.0 (2)	F2'—P1—F5	92.2 (4)
C16—C17—H17	120.5	F3'—P1—F5	92.9 (4)
C18—C17—H17	120.5	F3—P1—F5	87.63 (19)
C17—C16—C15	118.9 (2)	F1—P1—F5	91.21 (18)
C17—C16—H16	120.5	F2—P1—F4	176.3 (4)
C15—C16—H16	120.5	F1'—P1—F4	139.5 (6)
N1—C1—C2	122.8 (3)	F4'—P1—F4	50.2 (7)
N1—C1—H1	118.6	F6—P1—F4	90.8 (2)
C2—C1—H1	118.6	F2'—P1—F4	131.2 (6)
C2—C3—C4	119.2 (3)	F3—P1—F4	89.0 (3)
C2—C3—H3	120.4	F1—P1—F4	87.9 (3)
C4—C3—H3	120.4	F5—P1—F4	85.3 (2)
N1—C5—C4	123.4 (2)	O2—C20—N6	123.8 (2)
N1—C5—H5	118.3	O2—C20—C18	121.3 (2)
C4—C5—H5	118.3	N6—C20—C18	114.9 (2)
C12—C13—C8	120.9 (3)		
C26—C25—C24—C23	-0.5 (5)	C12—C13—C8—C7	-179.0 (3)
C28—C25—C24—C23	178.8 (3)	N4—C7—C8—C9	-3.3 (5)
C24—C25—C26—C27	1.2 (5)	N4—C7—C8—C13	176.4 (3)
C28—C25—C26—C27	-178.1 (3)	N1—C1—C2—C3	-0.4 (5)
C25—C26—C27—C22	-1.0 (5)	C4—C3—C2—C1	1.2 (5)
C25—C24—C23—C22	-0.4 (5)	C8—C9—C10—C11	-0.7 (5)
C26—C27—C22—C23	0.2 (5)	C12—C11—C10—C9	1.4 (5)
C26—C27—C22—C21	177.6 (3)	C14—C11—C10—C9	-179.0 (3)
C24—C23—C22—C27	0.6 (5)	O1—C6—N3—N4	2.7 (4)
C24—C23—C22—C21	-176.8 (3)	C4—C6—N3—N4	-176.6 (2)
C27—C22—C21—N7	169.9 (3)	C16—C15—N2—C19	0.9 (4)
C23—C22—C21—N7	-12.7 (5)	C16—C15—N2—Ag1	-173.5 (2)
C22—C21—N7—N6	177.8 (3)	C18—C19—N2—C15	-0.6 (4)
C21—N7—N6—C20	179.8 (3)	C18—C19—N2—Ag1	173.8 (2)
N2—C19—C18—C17	0.1 (4)	N1—Ag1—N2—C15	-169.9 (2)

N2—C19—C18—C20	178.6 (3)	N8 ⁱⁱ —Ag1—N2—C15	-13.3 (2)
O1—C6—C4—C5	-159.2 (3)	N1—Ag1—N2—C19	15.9 (4)
N3—C6—C4—C5	20.1 (4)	N8 ⁱⁱ —Ag1—N2—C19	172.5 (2)
O1—C6—C4—C3	16.2 (4)	C8—C7—N4—N3	179.3 (3)
N3—C6—C4—C3	-164.5 (3)	C6—N3—N4—C7	172.6 (3)
C19—C18—C17—C16	0.1 (4)	C4—C5—N1—C1	0.7 (4)
C20—C18—C17—C16	-178.5 (3)	C4—C5—N1—Ag1	-173.5 (2)
C18—C17—C16—C15	0.2 (4)	C2—C1—N1—C5	-0.6 (4)
N2—C15—C16—C17	-0.7 (5)	C2—C1—N1—Ag1	173.6 (2)
C5—C4—C3—C2	-1.1 (4)	N2—Ag1—N1—C5	-25.6 (4)
C6—C4—C3—C2	-176.9 (3)	N8 ⁱⁱ —Ag1—N1—C5	179.3 (2)
C3—C4—C5—N1	0.2 (4)	N2—Ag1—N1—C1	160.4 (2)
C6—C4—C5—N1	175.6 (3)	N8 ⁱⁱ —Ag1—N1—C1	5.3 (2)
C11—C12—C13—C8	-0.1 (5)	N7—N6—C20—O2	-3.9 (4)
C13—C12—C11—C10	-1.0 (5)	N7—N6—C20—C18	176.5 (2)
C13—C12—C11—C14	179.5 (3)	C19—C18—C20—O2	-150.7 (3)
C10—C9—C8—C13	-0.4 (5)	C17—C18—C20—O2	27.8 (4)
C10—C9—C8—C7	179.4 (3)	C19—C18—C20—N6	29.0 (4)
C12—C13—C8—C9	0.8 (5)	C17—C18—C20—N6	-152.5 (3)

Symmetry codes: (i) $x+1/2, -y+3/2, z-1/2$; (ii) $x-1/2, -y+3/2, z+1/2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N3—H28 \cdots O2 ⁱⁱⁱ	0.865 (18)	2.15 (2)	2.990 (3)	162 (3)
N6—H29 \cdots F5	0.853 (18)	2.21 (2)	3.001 (4)	155 (3)

Symmetry codes: (iii) $-x+1/2, -y+3/2, -z$.

Fig. 1




Fig. 2




Fig. 3

