

## Di- $\mu_{1,1}$ -azido-bis[azido(5,5'-dimethyl-2,2'-bipyridine)nickel(II)]

Jin Hou

Department of Chemistry and Chemical Engineering, South-East University, Nanjing 211189, and Nantong Entry–Exit Inspection and Quarantine Bureau, Nantong 226005, People's Republic of China

Correspondence e-mail: jinhou\_jinhou@yahoo.com.cn


Received 7 November 2008; accepted 14 November 2008

 Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.008$  Å;  $R$  factor = 0.056;  $wR$  factor = 0.149; data-to-parameter ratio = 15.9.

In the title azide-bridged dinuclear centrosymmetric nickel(II) complex,  $[\text{Ni}_2(\text{N}_3)_4(\text{C}_{12}\text{H}_{12}\text{N}_2)_2]$ , the  $\text{Ni}^{\text{II}}$  atom is five-coordinated by two N atoms of the 5,5'-dimethyl-2,2'-bipyridine ligand and three N atoms from three azide ligands in a distorted trigonal-bipyramidal geometry. The  $\text{Ni}\cdots\text{Ni}$  distance is 3.2398 (12) Å.

### Related literature

For general background, see: Abramo *et al.* (2002); Dey *et al.* (2007); Jiang *et al.* (2005). For related structures, see: Fu *et al.* (2005); Song *et al.* (2007).



### Experimental

#### Crystal data

$[\text{Ni}_2(\text{N}_3)_4(\text{C}_{12}\text{H}_{12}\text{N}_2)_2]$   
 $M_r = 654.01$   
 Monoclinic,  $P2_1/n$   
 $a = 7.938$  (2) Å  
 $b = 15.067$  (3) Å  
 $c = 11.755$  (2) Å  
 $\beta = 91.650$  (2)°

$V = 1405.3$  (5) Å<sup>3</sup>  
 $Z = 2$   
 Mo  $K\alpha$  radiation  
 $\mu = 1.39$  mm<sup>-1</sup>  
 $T = 298$  (2) K  
 $0.13 \times 0.10 \times 0.08$  mm

#### Data collection

Bruker SMART CCD area-detector diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 2001)  
 $T_{\text{min}} = 0.840$ ,  $T_{\text{max}} = 0.897$   
 11588 measured reflections  
 3060 independent reflections  
 2165 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.063$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$   
 $wR(F^2) = 0.149$   
 $S = 1.03$   
 3060 reflections  
 192 parameters

14 restraints  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.67$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.77$  e Å<sup>-3</sup>

**Table 1**

Selected bond lengths (Å).

Ni1—N1	2.145 (4)	Ni1—N6	2.041 (4)
Ni1—N2	2.081 (4)	Ni1—N6 <sup>i</sup>	2.175 (4)
Ni1—N3	2.064 (5)		

 Symmetry code: (i)  $-x, -y + 1, -z$ .

Data collection: SMART (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2712).

### References

- Abramo, G. P., Li, L. & Marks, T. J. (2002). *J. Am. Chem. Soc.* **124**, 13966–13967.  
 Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Bruker (2007). *SMART* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Dey, S. K., Abedin, T. S. M., Dawe, L. N., Tandon, S. S., Collins, J. L., Thompson, L. K., Postnikov, A. V., Alam, M. S. & Muller, P. (2007). *Inorg. Chem.* **46**, 7767–7781.  
 Fu, Y.-L., Xu, Z.-W., Ren, J.-L. & Ng, S. W. (2005). *Acta Cryst.* **E61**, m1897–m1899.  
 Jiang, Y.-B., Kou, H.-Z., Wang, R.-J., Cui, A.-L. & Ribas, J. (2005). *Inorg. Chem.* **44**, 709–715.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Song, W.-C., Zhang, M.-J., Tao, Y. & Li, J.-R. (2007). *Acta Cryst.* **E63**, m3062.

**supplementary materials**

*Acta Cryst.* (2008). E64, m1571 [ doi:10.1107/S1600536808037902 ]

## Di- $\mu_{1,1}$ -azido-bis[azido(5,5'-dimethyl-2,2'-bipyridine)nickel(II)]

J. Hou

### Comment

Polynuclear complexes play an important role in many fields, such as catalysis and magnetism (Dey *et al.*, 2007; Jiang *et al.*, 2005; Abramo *et al.*, 2002). The main strategy for the design of the polynuclear complexes is to use suitable bridging ligands. In this paper, we report the synthesis and molecular structure of the title azide-bridged dinuclear nickel(II) complex derived from 5,5'-dimethyl-[2,2']bipyridine.

The molecule of the title complex is located on a crystallographic centre of inversion (Fig. 1). The complex contains two NiL (L is 5,5'-dimethyl-[2,2']bipyridine) units connected to each other by two bridging azide ligands. The Ni<sup>II</sup> atom in the complex is five-coordinated by two N atoms of 5,5'-dimethyl-[2,2']bipyridine ligand and by three N atoms from three azide ligands in a trigonal-bipyramidal geometry. The bond lengths subtended at the metal center are within normal ranges (Song *et al.*, 2007; Fu *et al.*, 2005). The Ni $\cdots$ Ni distance is 3.2398 (12) Å.

### Experimental

5,5'-Dimethyl-[2,2']bipyridine (2 mmol, 368.3 mg), sodium azide (4 mmol, 261.2 mg) and nickel acetate tetrahydrate (2 mmol, 497.8 mg) were dissolved in methanol (100 ml). The mixture was stirred for 30 min at room temperature to give a green solution. The solution was kept still in air for a week, green block-shaped crystals of the title complex were formed.

### Refinement

H atoms were positioned geometrically (C–H = 0.93–0.96 Å) and refined using a riding model, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  and  $1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$ . The displacement ellipsoids of atoms N4 and N5 are extremely elongated and hence the  $U^{\text{ij}}$  parameters of these atoms were restrained to an approximate isotropic behaviour. The distance between atoms N3 and N4 was restrained to 1.23 (1) Å and that between atoms N4 and N5 was restrained to 1.13 (1) Å.

### Figures




Fig. 1. The molecular structure of the title compound, showing 30% probability displacement ellipsoids and the atom-numbering scheme.

## Di- $\mu$ ,1-azido-bis[azido(5,5'-dimethyl-2,2'-bipyridine)nickel(II)]

### Crystal data

[Ni<sub>2</sub>(N<sub>3</sub>)<sub>4</sub>(C<sub>12</sub>H<sub>12</sub>N<sub>2</sub>)<sub>2</sub>]

$M_r = 654.01$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 7.938$  (2) Å

$b = 15.067$  (3) Å

$c = 11.755$  (2) Å

$\beta = 91.650$  (2)°

$V = 1405.3$  (5) Å<sup>3</sup>

$Z = 2$

$F_{000} = 672$

$D_x = 1.546$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation

$\lambda = 0.71073$  Å

Cell parameters from 1576 reflections

$\theta = 2.3$ – $25.1$ °

$\mu = 1.39$  mm<sup>-1</sup>

$T = 298$  (2) K

Block, green

$0.13 \times 0.10 \times 0.08$  mm

### Data collection

Bruker SMART CCD area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 298$ (2) K

$\varphi$  and  $\omega$  scan

Absorption correction: multi-scan  
(SADABS; Bruker, 2001)

$T_{\min} = 0.840$ ,  $T_{\max} = 0.897$

11588 measured reflections

3060 independent reflections

2165 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.063$

$\theta_{\text{max}} = 27.0$ °

$\theta_{\text{min}} = 2.2$ °

$h = -10 \rightarrow 10$

$k = -19 \rightarrow 19$

$l = -15 \rightarrow 15$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.056$

$wR(F^2) = 0.149$

$S = 1.03$

3060 reflections

192 parameters

14 restraints

Primary atom site location: structure-invariant direct  
methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring  
sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0643P)^2 + 1.1062P]$$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} = 0.001$

$\Delta\rho_{\text{max}} = 0.67$  e Å<sup>-3</sup>

$\Delta\rho_{\text{min}} = -0.77$  e Å<sup>-3</sup>

Extinction correction: none

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor wR and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.03235 (7)	0.44079 (4)	0.11356 (5)	0.0386 (2)
N1	0.2280 (5)	0.3549 (3)	0.1785 (4)	0.0491 (10)
N2	-0.0777 (5)	0.3158 (3)	0.0994 (3)	0.0416 (9)
N3	0.0029 (5)	0.5108 (3)	0.2626 (4)	0.0459 (11)
N4	0.0958 (8)	0.5106 (4)	0.3252 (5)	0.0857 (18)
N5	0.1983 (12)	0.5160 (6)	0.4040 (7)	0.144 (3)
N6	0.1668 (5)	0.4913 (3)	-0.0170 (4)	0.0583 (12)
N7	0.2901 (6)	0.4579 (3)	-0.0571 (4)	0.0594 (12)
N8	0.4096 (8)	0.4282 (4)	-0.0950 (6)	0.094 (2)
C1	0.1794 (6)	0.2706 (3)	0.1926 (4)	0.0445 (11)
C2	0.2820 (7)	0.2113 (4)	0.2525 (5)	0.0573 (14)
H2	0.2475	0.1529	0.2628	0.069*
C3	0.4356 (7)	0.2398 (4)	0.2967 (5)	0.0632 (16)
H3	0.5057	0.2000	0.3357	0.076*
C4	0.4857 (6)	0.3259 (4)	0.2836 (5)	0.0575 (14)
C5	0.3771 (6)	0.3811 (4)	0.2219 (5)	0.0570 (14)
H5	0.4100	0.4396	0.2102	0.068*
C6	0.6485 (7)	0.3623 (5)	0.3348 (5)	0.085 (2)
H6A	0.6289	0.3847	0.4097	0.128*
H6B	0.6886	0.4095	0.2878	0.128*
H6C	0.7312	0.3159	0.3393	0.128*
C7	0.0139 (6)	0.2474 (3)	0.1428 (4)	0.0442 (12)
C8	-0.0506 (7)	0.1625 (4)	0.1383 (5)	0.0626 (15)
H8	0.0127	0.1155	0.1680	0.075*
C9	-0.2070 (8)	0.1468 (4)	0.0904 (5)	0.0652 (16)
H9	-0.2501	0.0894	0.0893	0.078*
C10	-0.3023 (6)	0.2160 (4)	0.0433 (5)	0.0530 (13)
C11	-0.2308 (6)	0.2993 (3)	0.0520 (4)	0.0477 (12)
H11	-0.2925	0.3472	0.0233	0.057*
C12	-0.4711 (7)	0.2026 (4)	-0.0126 (5)	0.0657 (16)
H12A	-0.5490	0.1824	0.0427	0.098*
H12B	-0.4629	0.1591	-0.0719	0.098*
H12C	-0.5103	0.2577	-0.0448	0.098*

## supplementary materials

---

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Ni1	0.0339 (3)	0.0328 (3)	0.0487 (4)	-0.0009 (3)	-0.0026 (2)	0.0057 (3)
N1	0.039 (2)	0.048 (3)	0.060 (3)	0.0018 (19)	-0.003 (2)	0.004 (2)
N2	0.039 (2)	0.040 (2)	0.046 (2)	0.0000 (17)	-0.0016 (18)	-0.0013 (17)
N3	0.035 (2)	0.036 (2)	0.067 (3)	0.0025 (18)	-0.006 (2)	-0.012 (2)
N4	0.119 (5)	0.043 (3)	0.097 (5)	0.001 (4)	0.045 (4)	-0.006 (3)
N5	0.165 (7)	0.149 (6)	0.115 (6)	-0.004 (6)	-0.029 (5)	0.002 (5)
N6	0.038 (2)	0.066 (3)	0.072 (3)	0.010 (2)	0.005 (2)	0.021 (2)
N7	0.051 (3)	0.056 (3)	0.070 (3)	0.005 (2)	-0.002 (2)	0.016 (2)
N8	0.071 (4)	0.105 (5)	0.108 (5)	0.037 (3)	0.020 (3)	0.011 (4)
C1	0.048 (3)	0.043 (3)	0.043 (3)	0.009 (2)	0.004 (2)	0.001 (2)
C2	0.065 (4)	0.048 (3)	0.058 (3)	0.010 (3)	0.002 (3)	0.002 (3)
C3	0.058 (4)	0.084 (4)	0.047 (3)	0.025 (3)	-0.009 (3)	0.004 (3)
C4	0.041 (3)	0.080 (4)	0.051 (3)	0.008 (3)	0.001 (2)	-0.001 (3)
C5	0.044 (3)	0.064 (4)	0.063 (4)	-0.003 (3)	-0.001 (3)	0.002 (3)
C6	0.049 (3)	0.131 (7)	0.074 (4)	-0.001 (4)	-0.014 (3)	0.003 (4)
C7	0.049 (3)	0.038 (3)	0.046 (3)	0.003 (2)	0.004 (2)	0.001 (2)
C8	0.060 (4)	0.047 (3)	0.080 (4)	0.003 (3)	-0.002 (3)	0.013 (3)
C9	0.067 (4)	0.047 (3)	0.081 (4)	-0.016 (3)	-0.001 (3)	0.004 (3)
C10	0.049 (3)	0.056 (3)	0.054 (3)	-0.008 (3)	0.006 (2)	-0.004 (3)
C11	0.045 (3)	0.047 (3)	0.050 (3)	0.001 (2)	0.001 (2)	0.002 (2)
C12	0.054 (3)	0.075 (4)	0.068 (4)	-0.020 (3)	-0.002 (3)	-0.009 (3)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Ni1—N1	2.145 (4)	C3—H3	0.93
Ni1—N2	2.081 (4)	C4—C5	1.387 (7)
Ni1—N3	2.064 (5)	C4—C6	1.512 (8)
Ni1—N6	2.041 (4)	C5—H5	0.93
Ni1—N6 <sup>i</sup>	2.175 (4)	C6—H6A	0.96
N1—C5	1.336 (6)	C6—H6B	0.96
N1—C1	1.339 (6)	C6—H6C	0.96
N2—C11	1.345 (6)	C7—C8	1.379 (7)
N2—C7	1.352 (6)	C8—C9	1.369 (8)
N3—N4	1.027 (6)	C8—H8	0.93
N4—N5	1.217 (7)	C9—C10	1.394 (8)
N6—N7	1.209 (6)	C9—H9	0.93
N6—Ni1 <sup>i</sup>	2.175 (4)	C10—C11	1.380 (7)
N7—N8	1.149 (7)	C10—C12	1.489 (7)
C1—C2	1.387 (7)	C11—H11	0.93
C1—C7	1.465 (7)	C12—H12A	0.96
C2—C3	1.380 (8)	C12—H12B	0.96
C2—H2	0.93	C12—H12C	0.96
C3—C4	1.367 (8)		
N6—Ni1—N3	121.5 (2)	C3—C4—C6	123.2 (5)

N6—Ni1—N2	120.32 (18)	C5—C4—C6	120.1 (6)
N3—Ni1—N2	118.23 (17)	N1—C5—C4	123.5 (5)
N6—Ni1—N1	95.97 (17)	N1—C5—H5	118.2
N3—Ni1—N1	96.02 (17)	C4—C5—H5	118.2
N2—Ni1—N1	77.30 (15)	C4—C6—H6A	109.5
N6—Ni1—N6 <sup>i</sup>	79.63 (18)	C4—C6—H6B	109.5
N3—Ni1—N6 <sup>i</sup>	95.99 (18)	H6A—C6—H6B	109.5
N2—Ni1—N6 <sup>i</sup>	94.98 (16)	C4—C6—H6C	109.5
N1—Ni1—N6 <sup>i</sup>	167.78 (18)	H6A—C6—H6C	109.5
C5—N1—C1	119.3 (5)	H6B—C6—H6C	109.5
C5—N1—Ni1	125.6 (4)	N2—C7—C8	119.8 (5)
C1—N1—Ni1	114.1 (3)	N2—C7—C1	115.8 (4)
C11—N2—C7	119.0 (4)	C8—C7—C1	124.3 (5)
C11—N2—Ni1	124.9 (3)	C9—C8—C7	120.5 (5)
C7—N2—Ni1	116.1 (3)	C9—C8—H8	119.8
N4—N3—Ni1	120.7 (5)	C7—C8—H8	119.8
N3—N4—N5	174.4 (8)	C8—C9—C10	120.6 (5)
N7—N6—Ni1	125.8 (4)	C8—C9—H9	119.7
N7—N6—Ni1 <sup>i</sup>	125.2 (4)	C10—C9—H9	119.7
Ni1—N6—Ni1 <sup>i</sup>	100.37 (18)	C11—C10—C9	115.7 (5)
N8—N7—N6	178.1 (6)	C11—C10—C12	121.3 (5)
N1—C1—C2	120.5 (5)	C9—C10—C12	123.0 (5)
N1—C1—C7	115.8 (4)	N2—C11—C10	124.3 (5)
C2—C1—C7	123.8 (5)	N2—C11—H11	117.8
C3—C2—C1	119.3 (5)	C10—C11—H11	117.8
C3—C2—H2	120.3	C10—C12—H12A	109.5
C1—C2—H2	120.3	C10—C12—H12B	109.5
C4—C3—C2	120.7 (5)	H12A—C12—H12B	109.5
C4—C3—H3	119.7	C10—C12—H12C	109.5
C2—C3—H3	119.7	H12A—C12—H12C	109.5
C3—C4—C5	116.7 (5)	H12B—C12—H12C	109.5

Symmetry codes: (i)  $-x, -y+1, -z$ .

Fig. 1

