

# [ $\mu$ -2,2,4,4,6,6-Hexakis(3,5-dimethylpyrazol-1-yl)-2 $\lambda^5$ ,4 $\lambda^5$ ,6 $\lambda^5$ -1,3,5,2,4,6-triazatriphosphinine]bis[bis(nitrato- $\kappa^2$ O,O')cadmium(II)]

Sung Yol Yun and Soon W. Lee\*

Department of Chemistry (BK21), Sungkyunkwan University, Natural Science Campus, Suwon 440-746, South Korea

Correspondence e-mail: soonwlee@skku.edu


Received 15 September 2008; accepted 1 November 2008

 Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.008$  Å;  $R$  factor = 0.044;  $wR$  factor = 0.112; data-to-parameter ratio = 20.9.

The complete title complex,  $[\text{Cd}_2(\text{NO}_3)_4(\text{C}_{30}\text{H}_{42}\text{N}_{15}\text{P}_3)]$ , is generated by crystallographic twofold symmetry, with one P and one N atom of the cyclotriphosphazene ligand located on the rotation axis. The non-planar cyclotriphosphazene ring accommodates two Cd ions, and only four out of six exocyclic pyrazolyl ligands are bound to the Cd metal atoms. Each of these two symmetry-related Cd atoms is coordinated by two bidentate nitrato ligands, two exocyclic pyrazolyl N atoms, and one cyclotriphosphazene N atom.

## Related literature

For background, see: Allen (1991); Byun *et al.* (1996); Chandrasekhar & Nagendran (2001); Mark *et al.* (2005); Thomas *et al.* (1997 and references therein). For the synthesis of the ligand, see: Thomas *et al.* (1993). For related structures, see: Yun & Lee (2008).



## Experimental

### Crystal data

$[\text{Cd}_2(\text{NO}_3)_4(\text{C}_{30}\text{H}_{42}\text{N}_{15}\text{P}_3)]$   
 $M_r = 1178.54$   
 Orthorhombic, *Fdd2*  
 $a = 28.2418$  (5) Å  
 $b = 36.2033$  (6) Å  
 $c = 10.3673$  (2) Å

$V = 10600.0$  (3) Å<sup>3</sup>  
 $Z = 8$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.96$  mm<sup>-1</sup>  
 $T = 296$  (2) K  
 $0.30 \times 0.14 \times 0.10$  mm

### Data collection

Bruker SMART CCD diffractometer  
 Absorption correction: multi-scan (*SADABS*; Bruker, 1997)  
 $T_{\min} = 0.851$ ,  $T_{\max} = 0.908$

32426 measured reflections  
 6260 independent reflections  
 5122 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.033$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$   
 $wR(F^2) = 0.112$   
 $S = 1.04$   
 6260 reflections  
 299 parameters  
 1 restraint

H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.85$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.48$  e Å<sup>-3</sup>  
 Absolute structure: Flack (1983),  
 2758 Friedel pairs  
 Flack parameter:  $-0.02$  (2)

**Table 1**

Selected bond lengths (Å).

Cd1—N1	2.546 (3)	Cd1—O2	2.365 (6)
Cd1—N4	2.265 (3)	Cd1—O5	2.367 (8)
Cd1—N8	2.311 (4)	Cd1—O4	2.373 (9)
Cd1—O1	2.509 (5)		

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINTE* (Bruker, 1997); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

This paper was supported by Samsung Research Fund, Sungkyunkwan University, 2008.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2796).

## References

- Allen, C. W. (1991). *Chem. Rev.* **91**, 119–135.  
 Bruker (1997). *SMART*, *SAINTE* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Byun, Y. G., Min, D. W., Do, J. H., Yun, H. S. & Do, Y. K. (1996). *Inorg. Chem.* **35**, 3981–3989.  
 Chandrasekhar, V. & Nagendran, S. (2001). *Chem. Soc. Rev.* **30**, 193–203.  
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.  
 Mark, J. E., Allcock, H. R. & West, R. (2005). *Inorganic Polymers*, 2nd ed., pp. 62–153 New York: Oxford University Press.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Thomas, K. R. J., Chandrasekhar, V., Pal, P., Scott, S. R., Hallford, R. & Cordes, A. W. (1993). *Inorg. Chem.* **32**, 606–611.  
 Thomas, K. R. J., Chandrasekhar, V., Zanello, P. & Laschi, F. (1997). *Polyhedron*, **16**, 1003–1011.  
 Yun, S. Y. & Lee, S. W. (2008). *Acta Cryst.* **E64**, m1084.

**supplementary materials**

*Acta Cryst.* (2008). E64, m1513 [ doi:10.1107/S160053680803585X ]

**[ $\mu$ -2,2,4,4,6,6-Hexakis(3,5-dimethylpyrazol-1-yl)-2 $\lambda^5$ ,4 $\lambda^5$ ,6 $\lambda^5$ -1,3,5,2,4,6-triazatriphosphinine]bis[bis(nitrato-  $\kappa^2O,O'$ )cadmium(II)]**

**S. Y. Yun and S. W. Lee**

**Comment**

Polyphosphazenes, linear or cyclic, are an important class of inorganic macromolecules (Mark *et al.*, 2005), and various cyclotriphosphazene derivatives are frequently used as ligands for the preparation of their intriguing coordination and organometallic complexes (Allen, 1991; Chandrasekhar & Nagendran, 2001). Particular attention has been paid to the six-membered cyclotriphosphazene  $N_3P_3(3,5\text{-Me}_2\text{pz})_6$  ( $3,5\text{-Me}_2\text{pz} = 3,5\text{-dimethylpyrazolyl}$ ), due to its several potential donor sites such as the exocyclic pyrazolyl nitrogen atoms and the central cyclotriphosphazene ring nitrogen and phosphorus atoms. This ligand binds to transition metals *via* (1) two non-geminal pyrazolyl N atoms (non-geminal  $N_2$  coordination), (2) two non-geminal pyrazolyl N atoms and one cyclotriphosphazene ring nitrogen (non-geminal  $N_3$  coordination), (3) two geminal pyrazolyl N atoms (geminal  $N_2$  coordination), or (4) two geminal pyrazolyl N atoms and one ring nitrogen (geminal  $N_3$  coordination) (Thomas *et al.*, 1997). We recently reported the structure of a  $C_3$ -symmetric tripalladium–cyclotriphosphazene complex, in which the cyclotriphosphazene exhibits the geminal  $N_2$  coordination mode (Yun & Lee, 2008). In this paper, we describe the preparation and structure of the title compound, (I), a dicadmium–cyclotriphosphazene complex [ $Cd_2(NO_3)_4(N_3P_3(3,5\text{-Me}_2\text{pz})_6)$ ].

The molecular structure of (I) is given in Fig. 1, which demonstrates the non-geminal  $N_3$  coordination mode of the cyclotriphosphazene ligand. This molecule possesses a crystallographic 2-fold axis passing through the P2 and N2 atoms, which explains the  $Z$  value of 8 instead of 16. The cyclotriphosphazene ring is severely distorted from planarity with an average atomic displacement of 0.146 Å. Each Cd(II) metal is seven-coordinate and bonded to four O atoms from two nitrates, two N atoms from two imidazole rings, and one nitrogen atom from the cyclotriphosphazene ring (Table 1). Four imidazole N atoms coordinate to the two Cd metals to form four 5-membered ( $PdPN_3$ ) chelate rings. The fact that the cyclotriphosphazene ring accommodates only two rather than three  $Cd(NO_3)_2$  units may be attributed to the steric bulk of the 7-coordinate Cd metals.

The Cd— $N_{\text{pyz}}$  bond lengths [2.265 (3)–2.311 (4) Å] are significantly shorter than the Cd— $N_{\text{ring}}$  bond length [2.546 (3) Å], indicating that the Cd ions interact more strongly with the imidazole N atoms than with the cyclotriphosphazene ring N atoms. Consistent with our expectation, the average P— $N_{\text{ring}}$  bond length [1.583 (3) Å] is considerably shorter than the average P— $N_{\text{pyz}}$  bond length [1.677 (3) Å]. The Cd···Cd separation is 7.0177 (6) Å, which is shorter than the corresponding separation (7.195 Å) observed in the chloro analogue [ $Cd_2Cl_4(N_3P_3(3,5\text{-Me}_2\text{pz})_6)$ ] (Byun *et al.*, 1996).

**Experimental**

The ligand was prepared by the literature method (Thomas *et al.*, 1993). An acetone (30 ml) solution containing  $Cd(NO_3)_2 \cdot 4H_2O$  (0.144 g, 0.75 mmol) and ligand  $N_3P_3(3,5\text{-Me}_2\text{pz})_6$  (0.176 g, 0.25 mmol) was stirred for 24 h at room temperature. The resulting white solution was filtered off, washed with diethyl ether (6 ml  $\times$  2) and then hexane (5 ml  $\times$  2)

## supplementary materials

to give a white solid, which was crystallized from acetone/hexane (1:1 v/v) to yield colorless blocks of (I). IR (KBr,  $\text{cm}^{-1}$ ): 2964 (s), 2362 (m), 1576 (m), 1383 (s), 1260 (s), 1095 (s), 1028 (s), 805 (s). mp: 411–413 K.

### Refinement

The hydrogen atoms were generated in ideal positions ( $\text{C}-\text{H} = 0.93\text{--}0.96\text{\AA}$ ) and refined in a riding model. The nitrate ligands are slightly disordered, but the disorder was not resolved and anisotropic refinement applying several possible site occupation factors was unstable. Site occupancy refinements of the nitrate atoms all yielded values close to unity.

### Figures




Fig. 1. The molecular structure of (I) showing 50% probability displacement ellipsoids. H atoms are omitted for clarity. Atoms with the suffix A are generated by the symmetry operation  $(-x, -y, z)$ .

**$[\mu\text{-}2,2,4,4,6,6\text{-Hexakis}(3,5\text{-dimethylpyrazol-}1\text{-yl})\text{-}2\lambda^5,4\lambda^5,6\lambda^5\text{-}1,3,5,2,4,6\text{-triazatriphosphinine}]_{\text{bis}}[\text{bis}(\text{nitrato-}\kappa^2\text{O},\text{O}')\text{cadmium(II)}]$**

### Crystal data

$[\text{Cd}_2(\text{NO}_3)_4(\text{C}_{30}\text{H}_{42}\text{N}_{15}\text{P}_3)]$

$M_r = 1178.54$

Orthorhombic,  $Fdd2$

Hall symbol:  $F\ 2\ -2d$

$a = 28.2418\ (5)\ \text{\AA}$

$b = 36.2033\ (6)\ \text{\AA}$

$c = 10.3673\ (2)\ \text{\AA}$

$V = 10600.0\ (3)\ \text{\AA}^3$

$Z = 8$

$F_{000} = 4736$

$D_x = 1.477\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 9939 reflections

$\theta = 2.2\text{--}25.5^\circ$

$\mu = 0.96\ \text{mm}^{-1}$

$T = 296\ (2)\ \text{K}$

BLOCK, colourless

$0.30 \times 0.14 \times 0.10\ \text{mm}$

### Data collection

Bruker SMART CCD  
diffractometer

Radiation source: sealed tube

Monochromator: graphite

$T = 296(2)\ \text{K}$

$\omega$  scans

Absorption correction: multi-scan  
(SADABS; Bruker, 1997)

$T_{\min} = 0.851$ ,  $T_{\max} = 0.908$

32426 measured reflections

6260 independent reflections

5122 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.033$

$\theta_{\max} = 28.4^\circ$

$\theta_{\min} = 2.2^\circ$

$h = -36 \rightarrow 28$

$k = -48 \rightarrow 40$

$l = -13 \rightarrow 13$

Refinement

Refinement on $F^2$	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.044$	$w = 1/[\sigma^2(F_o^2) + (0.0676P)^2 + 6.1853P]$
$wR(F^2) = 0.112$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.04$	$(\Delta/\sigma)_{\max} = 0.001$
6260 reflections	$\Delta\rho_{\max} = 0.85 \text{ e } \text{\AA}^{-3}$
299 parameters	$\Delta\rho_{\min} = -0.48 \text{ e } \text{\AA}^{-3}$
1 restraint	Extinction correction: none
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 2758 Friedel pairs
Secondary atom site location: difference Fourier map	Flack parameter: $-0.02 (2)$

Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	-0.083094 (12)	-0.072056 (9)	0.52427 (3)	0.05498 (12)
P1	-0.04843 (3)	-0.00189 (2)	0.74616 (9)	0.0339 (2)
P2	0.0000	0.0000	0.51827 (12)	0.0311 (2)
O1	-0.0099 (2)	-0.0965 (2)	0.6318 (6)	0.1100 (19)
O2	-0.0638 (3)	-0.13446 (15)	0.5667 (8)	0.144 (3)
O3	-0.0064 (4)	-0.1554 (3)	0.6716 (11)	0.251 (7)
O4	-0.1372 (4)	-0.0530 (3)	0.3622 (15)	0.258 (8)
O5	-0.1455 (4)	-0.1014 (3)	0.4113 (10)	0.195 (5)
O6	-0.1902 (3)	-0.0824 (2)	0.2697 (10)	0.170 (4)
N1	-0.04555 (11)	-0.01150 (8)	0.5958 (3)	0.0357 (7)
N2	0.0000	0.0000	0.8214 (4)	0.0408 (10)
N3	-0.08391 (12)	-0.03495 (9)	0.8082 (3)	0.0429 (8)
N4	-0.10447 (12)	-0.06131 (10)	0.7314 (3)	0.0450 (8)
N5	-0.07821 (10)	0.03652 (9)	0.7738 (4)	0.0431 (7)
N6	-0.12225 (13)	0.03751 (11)	0.7149 (4)	0.0512 (9)
N7	0.01317 (12)	-0.03608 (9)	0.4208 (3)	0.0398 (7)

## supplementary materials

---

N8	-0.02456 (14)	-0.05705 (10)	0.3775 (4)	0.0490 (8)
N9	-0.0244 (3)	-0.1266 (3)	0.6295 (8)	0.114 (2)
N10	-0.1568 (2)	-0.07751 (19)	0.3548 (8)	0.0857 (18)
C1	-0.10083 (18)	-0.03802 (13)	0.9323 (4)	0.0536 (11)
C2	-0.1308 (2)	-0.06616 (15)	0.9360 (5)	0.0703 (15)
H2	-0.1472	-0.0748	1.0078	0.084*
C3	-0.13292 (19)	-0.08072 (14)	0.8077 (5)	0.0593 (12)
C4	-0.0847 (3)	-0.01309 (18)	1.0404 (5)	0.0837 (18)
H4A	-0.0628	0.0048	1.0073	0.126*
H4B	-0.0695	-0.0276	1.1059	0.126*
H4C	-0.1116	-0.0007	1.0768	0.126*
C5	-0.1598 (2)	-0.11282 (18)	0.7556 (7)	0.0834 (18)
H5A	-0.1533	-0.1154	0.6652	0.125*
H5B	-0.1931	-0.1088	0.7681	0.125*
H5C	-0.1504	-0.1349	0.8001	0.125*
C6	-0.07031 (19)	0.06865 (11)	0.8451 (4)	0.0486 (10)
C7	-0.11000 (19)	0.08900 (14)	0.8317 (5)	0.0606 (12)
H7	-0.1160	0.1119	0.8692	0.073*
C8	-0.14106 (18)	0.06886 (13)	0.7490 (5)	0.0607 (12)
C9	-0.02660 (19)	0.07712 (13)	0.9190 (5)	0.0615 (13)
H9A	-0.0046	0.0571	0.9103	0.092*
H9B	-0.0344	0.0804	1.0084	0.092*
H9C	-0.0126	0.0994	0.8862	0.092*
C10	-0.1886 (3)	0.0808 (2)	0.7004 (8)	0.097 (2)
H10A	-0.2016	0.0618	0.6464	0.146*
H10B	-0.1853	0.1031	0.6515	0.146*
H10C	-0.2094	0.0850	0.7722	0.146*
C11	0.05422 (16)	-0.04550 (12)	0.3580 (4)	0.0465 (10)
C12	0.0421 (2)	-0.07391 (11)	0.2775 (5)	0.0578 (12)
H12	0.0625	-0.0868	0.2233	0.069*
C13	-0.0061 (2)	-0.07985 (12)	0.2915 (4)	0.0569 (12)
C14	0.10103 (19)	-0.02914 (17)	0.3787 (5)	0.0659 (14)
H14A	0.0987	-0.0098	0.4416	0.099*
H14B	0.1126	-0.0191	0.2989	0.099*
H14C	0.1225	-0.0478	0.4091	0.099*
C15	-0.0358 (3)	-0.10831 (18)	0.2264 (6)	0.092 (2)
H15A	-0.0680	-0.1059	0.2546	0.138*
H15B	-0.0242	-0.1325	0.2481	0.138*
H15C	-0.0342	-0.1049	0.1347	0.138*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cd1	0.0606 (2)	0.05662 (19)	0.04776 (17)	-0.02047 (15)	0.00994 (15)	-0.01297 (14)
P1	0.0381 (5)	0.0323 (4)	0.0314 (4)	0.0014 (4)	0.0036 (4)	-0.0026 (3)
P2	0.0365 (6)	0.0296 (5)	0.0272 (5)	0.0004 (5)	0.000	0.000
O1	0.095 (4)	0.137 (5)	0.098 (4)	0.043 (4)	0.016 (3)	0.033 (4)
O2	0.189 (6)	0.068 (3)	0.176 (8)	-0.022 (4)	0.092 (6)	-0.016 (4)

O3	0.258 (10)	0.203 (8)	0.292 (12)	0.150 (8)	0.168 (9)	0.191 (9)
O4	0.165 (7)	0.185 (8)	0.422 (19)	-0.087 (7)	-0.175 (11)	0.126 (11)
O5	0.226 (9)	0.213 (10)	0.146 (7)	-0.137 (9)	-0.036 (7)	-0.002 (7)
O6	0.143 (6)	0.135 (5)	0.231 (9)	-0.028 (4)	-0.085 (7)	-0.035 (6)
N1	0.0366 (16)	0.0361 (15)	0.0343 (15)	-0.0021 (13)	0.0017 (12)	-0.0047 (12)
N2	0.046 (3)	0.045 (2)	0.031 (2)	0.000 (2)	0.000	0.000
N3	0.0471 (19)	0.0413 (18)	0.0403 (18)	-0.0063 (14)	0.0082 (14)	-0.0033 (14)
N4	0.0414 (18)	0.0480 (18)	0.0455 (18)	-0.0091 (16)	0.0084 (15)	-0.0042 (15)
N5	0.0419 (16)	0.0422 (16)	0.0452 (17)	0.0065 (13)	-0.0021 (16)	-0.0110 (15)
N6	0.0417 (19)	0.057 (2)	0.055 (2)	0.0120 (16)	-0.0053 (16)	-0.0146 (17)
N7	0.0434 (18)	0.0391 (17)	0.0368 (15)	0.0047 (14)	0.0064 (14)	-0.0042 (13)
N8	0.063 (2)	0.0416 (18)	0.0430 (18)	-0.0095 (17)	0.0077 (16)	-0.0141 (15)
N9	0.114 (6)	0.130 (7)	0.098 (5)	0.027 (6)	0.063 (4)	0.026 (5)
N10	0.072 (3)	0.069 (4)	0.116 (5)	-0.033 (3)	0.000 (3)	-0.008 (3)
C1	0.065 (3)	0.051 (2)	0.045 (2)	-0.009 (2)	0.017 (2)	-0.0054 (19)
C2	0.084 (4)	0.069 (3)	0.058 (3)	-0.013 (3)	0.033 (3)	0.006 (2)
C3	0.060 (3)	0.059 (3)	0.059 (3)	-0.018 (2)	0.016 (2)	0.002 (2)
C4	0.126 (5)	0.085 (4)	0.040 (3)	-0.028 (4)	0.021 (3)	-0.013 (3)
C5	0.084 (4)	0.087 (4)	0.080 (4)	-0.042 (3)	0.016 (3)	-0.003 (3)
C6	0.065 (3)	0.039 (2)	0.042 (2)	0.0061 (19)	0.002 (2)	-0.0057 (16)
C7	0.064 (3)	0.048 (3)	0.070 (3)	0.016 (2)	-0.008 (2)	-0.019 (2)
C8	0.059 (3)	0.065 (3)	0.058 (3)	0.026 (2)	0.003 (2)	-0.013 (2)
C9	0.067 (3)	0.051 (3)	0.066 (3)	-0.001 (2)	-0.004 (3)	-0.025 (2)
C10	0.078 (4)	0.099 (5)	0.114 (5)	0.039 (4)	-0.023 (4)	-0.028 (4)
C11	0.055 (3)	0.048 (2)	0.0358 (19)	0.0123 (19)	0.0070 (18)	0.0062 (17)
C12	0.079 (3)	0.053 (2)	0.042 (2)	0.013 (2)	0.012 (2)	-0.008 (2)
C13	0.086 (3)	0.044 (2)	0.040 (2)	-0.005 (2)	0.016 (2)	-0.0110 (19)
C14	0.049 (3)	0.097 (4)	0.052 (3)	0.013 (3)	0.010 (2)	-0.007 (3)
C15	0.136 (6)	0.072 (3)	0.069 (3)	-0.034 (4)	0.019 (4)	-0.036 (3)

*Geometric parameters (Å, °)*

Cd1—N1	2.546 (3)	C1—C4	1.509 (7)
Cd1—N4	2.265 (3)	C2—C3	1.432 (8)
Cd1—N8	2.311 (4)	C2—H2	0.9300
Cd1—O1	2.509 (5)	C3—C5	1.490 (8)
Cd1—O2	2.365 (6)	C4—H4A	0.9600
Cd1—O5	2.367 (8)	C4—H4B	0.9600
Cd1—O4	2.373 (9)	C4—H4C	0.9600
P1—N2	1.576 (2)	C5—H5A	0.9600
P1—N1	1.599 (3)	C5—H5B	0.9600
P1—N5	1.650 (3)	C5—H5C	0.9600
P1—N3	1.688 (3)	C6—C7	1.349 (7)
P2—N1	1.573 (3)	C6—C9	1.485 (7)
P2—N1 <sup>i</sup>	1.573 (3)	C7—C8	1.427 (7)
P2—N7 <sup>i</sup>	1.693 (3)	C7—H7	0.9300
P2—N7	1.693 (3)	C8—C10	1.498 (8)
O1—N9	1.165 (11)	C9—H9A	0.9600
O2—N9	1.323 (11)	C9—H9B	0.9600

## supplementary materials

---

O3—N9	1.239 (10)	C9—H9C	0.9600
O4—N10	1.049 (9)	C10—H10A	0.9600
O5—N10	1.094 (12)	C10—H10B	0.9600
O6—N10	1.302 (10)	C10—H10C	0.9600
N2—P1 <sup>i</sup>	1.576 (2)	C11—C12	1.368 (6)
N3—N4	1.372 (5)	C11—C14	1.464 (8)
N3—C1	1.376 (6)	C12—C13	1.386 (8)
N4—C3	1.328 (6)	C12—H12	0.9300
N5—N6	1.386 (5)	C13—C15	1.489 (7)
N5—C6	1.396 (5)	C14—H14A	0.9600
N6—C8	1.302 (5)	C14—H14B	0.9600
N7—C11	1.373 (5)	C14—H14C	0.9600
N7—N8	1.383 (5)	C15—H15A	0.9600
N8—C13	1.323 (6)	C15—H15B	0.9600
C1—C2	1.325 (7)	C15—H15C	0.9600
N4—Cd1—N8	140.76 (12)	C2—C1—N3	108.1 (4)
N4—Cd1—O2	92.8 (2)	C2—C1—C4	129.1 (5)
N8—Cd1—O2	100.51 (19)	N3—C1—C4	122.7 (4)
N4—Cd1—O5	110.4 (3)	C1—C2—C3	106.4 (4)
N8—Cd1—O5	108.2 (3)	C1—C2—H2	126.8
O2—Cd1—O5	80.4 (5)	C3—C2—H2	126.8
N4—Cd1—O4	116.7 (4)	N4—C3—C2	109.5 (4)
N8—Cd1—O4	85.8 (4)	N4—C3—C5	120.3 (5)
O2—Cd1—O4	123.8 (4)	C2—C3—C5	130.2 (5)
O5—Cd1—O4	45.7 (4)	C1—C4—H4A	109.5
N4—Cd1—O1	81.87 (16)	C1—C4—H4B	109.5
N8—Cd1—O1	77.65 (18)	H4A—C4—H4B	109.5
O2—Cd1—O1	52.5 (3)	C1—C4—H4C	109.5
O5—Cd1—O1	132.4 (4)	H4A—C4—H4C	109.5
O4—Cd1—O1	161.3 (4)	H4B—C4—H4C	109.5
N4—Cd1—N1	71.75 (11)	C3—C5—H5A	109.5
N8—Cd1—N1	72.03 (11)	C3—C5—H5B	109.5
O2—Cd1—N1	132.3 (3)	H5A—C5—H5B	109.5
O5—Cd1—N1	147.2 (4)	C3—C5—H5C	109.5
O4—Cd1—N1	103.0 (3)	H5A—C5—H5C	109.5
O1—Cd1—N1	80.3 (2)	H5B—C5—H5C	109.5
N2—P1—N1	116.61 (18)	C7—C6—N5	105.6 (4)
N2—P1—N5	108.63 (13)	C7—C6—C9	129.1 (4)
N1—P1—N5	112.25 (18)	N5—C6—C9	125.3 (4)
N2—P1—N3	110.93 (16)	C6—C7—C8	107.1 (4)
N1—P1—N3	104.34 (17)	C6—C7—H7	126.5
N5—P1—N3	103.21 (17)	C8—C7—H7	126.5
N1—P2—N1 <sup>i</sup>	118.5 (2)	N6—C8—C7	111.0 (4)
N1—P2—N7 <sup>i</sup>	109.24 (16)	N6—C8—C10	121.7 (5)
N1 <sup>i</sup> —P2—N7 <sup>i</sup>	106.30 (16)	C7—C8—C10	127.3 (4)
N1—P2—N7	106.30 (16)	C6—C9—H9A	109.5
N1 <sup>i</sup> —P2—N7	109.24 (16)	C6—C9—H9B	109.5

N7 <sup>i</sup> —P2—N7	106.7 (2)	H9A—C9—H9B	109.5
N9—O1—Cd1	91.9 (6)	C6—C9—H9C	109.5
N9—O2—Cd1	94.6 (5)	H9A—C9—H9C	109.5
N10—O4—Cd1	98.5 (7)	H9B—C9—H9C	109.5
N10—O5—Cd1	97.3 (6)	C8—C10—H10A	109.5
P2—N1—P1	118.8 (2)	C8—C10—H10B	109.5
P2—N1—Cd1	114.81 (15)	H10A—C10—H10B	109.5
P1—N1—Cd1	116.75 (16)	C8—C10—H10C	109.5
P1—N2—P1 <sup>i</sup>	120.7 (3)	H10A—C10—H10C	109.5
N4—N3—C1	109.8 (3)	H10B—C10—H10C	109.5
N4—N3—P1	121.5 (3)	C12—C11—N7	105.4 (4)
C1—N3—P1	128.3 (3)	C12—C11—C14	128.2 (4)
C3—N4—N3	106.2 (4)	N7—C11—C14	126.3 (4)
C3—N4—Cd1	129.4 (3)	C11—C12—C13	107.3 (4)
N3—N4—Cd1	123.9 (2)	C11—C12—H12	126.3
N6—N5—C6	110.8 (3)	C13—C12—H12	126.3
N6—N5—P1	113.7 (3)	N8—C13—C12	111.2 (4)
C6—N5—P1	135.5 (3)	N8—C13—C15	121.0 (5)
C8—N6—N5	105.6 (4)	C12—C13—C15	127.8 (5)
C11—N7—N8	111.1 (3)	C11—C14—H14A	109.5
C11—N7—P2	131.3 (3)	C11—C14—H14B	109.5
N8—N7—P2	116.6 (2)	H14A—C14—H14B	109.5
C13—N8—N7	104.9 (4)	C11—C14—H14C	109.5
C13—N8—Cd1	125.4 (3)	H14A—C14—H14C	109.5
N7—N8—Cd1	117.8 (2)	H14B—C14—H14C	109.5
O1—N9—O3	129.7 (12)	C13—C15—H15A	109.5
O1—N9—O2	120.5 (9)	C13—C15—H15B	109.5
O3—N9—O2	109.7 (12)	H15A—C15—H15B	109.5
O4—N10—O5	118.3 (10)	C13—C15—H15C	109.5
O4—N10—O6	123.1 (10)	H15A—C15—H15C	109.5
O5—N10—O6	117.9 (8)	H15B—C15—H15C	109.5

Symmetry codes: (i)  $-x, -y, z$ .

