

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-(3,4-Dihydroxyphenyl)-2-(4-fluorophenyl)ethanone

Xiao-Qing Song

School of Chemical and Material Engineering, Jiangnan University, Lihu Road No. 1800 Wuxi, Wuxi 214122, People's Republic of China
Correspondence e-mail: owengoal13@163.com


Received 28 October 2008; accepted 31 October 2008

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å;
 R factor = 0.050; wR factor = 0.145; data-to-parameter ratio = 12.6.

In the title compound, $\text{C}_{14}\text{H}_{11}\text{FO}_3$, the dihedral angle between the aromatic rings is $69.11(8)^\circ$. An intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond is present. Intermolecular $\text{O}-\text{H}\cdots\text{O}$ interactions help to establish the packing.

Related literature

For bond-length data, see: Allen *et al.* (1987). For background on deoxybenzoins, see: Li *et al.* (2007, 2008).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{11}\text{FO}_3$
 $M_r = 246.23$
Monoclinic, $P2_1/c$
 $a = 8.1640(16)$ Å
 $b = 5.9120(12)$ Å
 $c = 24.946(6)$ Å
 $\beta = 105.33(3)^\circ$

$V = 1161.2(4)$ Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.11$ mm⁻¹
 $T = 293(2)$ K
 $0.28 \times 0.25 \times 0.17$ mm

Data collection

Enraf-Nonius CAD-4
diffractometer
Absorption correction: ψ scan
(North *et al.*, 1968)
 $T_{\min} = 0.970$, $T_{\max} = 0.982$

2229 measured reflections
2072 independent reflections
1452 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.145$
 $S = 1.04$
2072 reflections

164 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.18$ e Å⁻³
 $\Delta\rho_{\min} = -0.17$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1A}\cdots\text{O2}$	0.82	2.28	2.690 (2)	111
$\text{O1}-\text{H1A}\cdots\text{O2}^{\text{i}}$	0.82	2.16	2.876 (3)	146
$\text{O2}-\text{H2B}\cdots\text{O3}^{\text{ii}}$	0.82	1.92	2.744 (2)	178

Symmetry codes: (i) $-x, -y, -z$; (ii) $x - 1, y, z$.

Data collection: *CAD-4 Software* (Enraf-Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

I gratefully acknowledge financial support from the Science Foundation for the Youth of Jiangnan University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2831).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Enraf-Nonius (1989). *CAD-4 Software*. Enraf-Nonius, Delft, The Netherlands.
Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
Li, H.-Q., Xu, C., Li, H.-S., Xiao, Z.-P., Shi, L. & Zhu, H.-L. (2007). *ChemMedChem*, **2**, 1361–1369.
Li, H.-Q., Xue, J.-Y., Shi, L., Gui, S.-Y. & Zhu, H.-L. (2008). *Eur. J. Med. Chem.* **43**, 662–667.
North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, o2282 [doi:10.1107/S1600536808035733]

1-(3,4-Dihydroxyphenyl)-2-(4-fluorophenyl)ethanone

X.-Q. Song

Comment

Doxybenzoin derivatives play an important role in organic chemistry (Li *et al.*, 2007; Li *et al.*, 2008). In the title compound, (I) (Fig. 1), the bond lengths and angles are within normal ranges (Allen *et al.*, 1987). The dihedral angle between the least-squares planes of the two benzene rings is 69.11 (8) °. In the crystal, O—H...O hydrogen bonds (Table 1) help to establish the packing.

Experimental

Pyrocatechol (0.050 mol) and 2-(4-fluorophenyl)acetic acid (0.050 mol) were dissolved into freshly distilled BF₃Et₂O under argon. The mixture was stirred at room temperature and then poured in an ice bath. The resulting mixture was extracted with ethyl acetate, and the organic layer was washed with aq. dried (Na₂S₁O₄), and evaporated. The white deposits precipitated were separated from the solvents by filtration. They were washed with aqueous saturated Na₁H₁C₁O₃ twice. The solid was dissolved in acetone (15 ml) and stirred for about 10 min to give a clear solution. After keeping the solution in air for 10 d, colorless blocks of (I) were formed at the bottom of the vessel on slow evaporation of the solvent. They were collected, washed three times with acetone and dried in a vacuum desiccator using CaCl₂. The compound was isolated in 90% yield.

Refinement

All the H atoms were positioned geometrically (C—H = 0.93–0.96 Å, O—H = 0.82 Å) and refined as riding, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. The molecular structure of (I) showing 30% probability displacement ellipsoids for the non-hydrogen atoms.

1-(3,4-Dihydroxyphenyl)-2-(4-fluorophenyl)ethanone

Crystal data

C₁₄H₁₁FO₃

$M_r = 246.23$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 8.1640(16)$ Å

$F_{000} = 512$

$D_x = 1.408$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 25 reflections

$\theta = 9\text{--}12^\circ$

supplementary materials

$b = 5.9120(12) \text{ \AA}$
 $c = 24.946(6) \text{ \AA}$
 $\beta = 105.33(3)^\circ$
 $V = 1161.2(4) \text{ \AA}^3$
 $Z = 4$

$\mu = 0.11 \text{ mm}^{-1}$
 $T = 293(2) \text{ K}$
Block, colorless
 $0.28 \times 0.25 \times 0.17 \text{ mm}$

Data collection

Enraf–Nonius CAD-4 diffractometer
Radiation source: fine-focus sealed tube
Monochromator: graphite
 $T = 293(2) \text{ K}$
 $\omega/2\theta$ scans
Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\min} = 0.970$, $T_{\max} = 0.982$
2229 measured reflections

2072 independent reflections
1452 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$
 $\theta_{\max} = 25.2^\circ$
 $\theta_{\min} = 1.7^\circ$
 $h = -9 \rightarrow 0$
 $k = -7 \rightarrow 0$
 $l = -28 \rightarrow 29$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.145$
 $S = 1.04$
2072 reflections
164 parameters

Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0627P)^2 + 0.5485P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.18 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.17 \text{ e \AA}^{-3}$
Extinction correction: SHELXL97 (Sheldrick, 2008),
 $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Extinction coefficient: 0.019 (3)

Primary atom site location: structure-invariant direct methods
Secondary atom site location: difference Fourier map

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.5159 (3)	0.4835 (4)	0.08518 (10)	0.0502 (6)
C2	0.4714 (3)	0.2980 (4)	0.04986 (10)	0.0536 (7)
H2A	0.5544	0.2229	0.0374	0.064*
C3	0.3067 (3)	0.2257 (4)	0.03341 (11)	0.0533 (6)
C4	0.1810 (3)	0.3396 (4)	0.05140 (10)	0.0496 (6)
C5	0.2243 (3)	0.5232 (5)	0.08571 (11)	0.0609 (7)
H5A	0.1406	0.5990	0.0977	0.073*
C6	0.3901 (3)	0.5969 (5)	0.10269 (12)	0.0622 (8)
H6A	0.4177	0.7224	0.1258	0.075*
C7	0.6960 (3)	0.5566 (4)	0.10307 (10)	0.0502 (6)
C8	0.7428 (3)	0.7480 (5)	0.14453 (12)	0.0636 (8)
H8A	0.6923	0.8862	0.1266	0.076*
H8B	0.6923	0.7179	0.1749	0.076*
C9	0.9297 (3)	0.7876 (4)	0.16843 (10)	0.0496 (6)
C10	1.0273 (3)	0.6323 (4)	0.20441 (11)	0.0543 (7)
H10A	0.9768	0.5008	0.2129	0.065*
C11	1.1974 (3)	0.6678 (5)	0.22796 (12)	0.0614 (7)
H11A	1.2620	0.5620	0.2522	0.074*
C12	1.2691 (3)	0.8598 (5)	0.21518 (13)	0.0640 (8)
C13	1.1793 (4)	1.0184 (5)	0.17977 (13)	0.0693 (8)
H13A	1.2317	1.1486	0.1715	0.083*
C14	1.0083 (4)	0.9806 (5)	0.15645 (12)	0.0624 (7)
H14A	0.9450	1.0872	0.1322	0.075*
F1	1.4375 (2)	0.8958 (4)	0.23816 (10)	0.1039 (7)
O1	0.2648 (2)	0.0477 (4)	−0.00250 (10)	0.0813 (7)
H1A	0.1726	−0.0035	−0.0012	0.122*
O2	0.0199 (2)	0.2558 (3)	0.03311 (8)	0.0615 (5)
H2B	−0.0421	0.3199	0.0493	0.092*
O3	0.8048 (2)	0.4626 (3)	0.08579 (8)	0.0650 (6)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0430 (13)	0.0574 (15)	0.0503 (14)	0.0101 (11)	0.0126 (10)	−0.0043 (12)
C2	0.0470 (14)	0.0553 (15)	0.0637 (16)	0.0095 (12)	0.0240 (12)	−0.0087 (13)
C3	0.0513 (14)	0.0511 (15)	0.0610 (15)	0.0044 (12)	0.0208 (12)	−0.0110 (13)
C4	0.0426 (13)	0.0552 (15)	0.0528 (14)	0.0084 (11)	0.0157 (11)	0.0012 (12)
C5	0.0429 (14)	0.0747 (19)	0.0676 (17)	0.0135 (13)	0.0190 (12)	−0.0202 (15)
C6	0.0477 (14)	0.0707 (18)	0.0689 (17)	0.0087 (13)	0.0169 (12)	−0.0199 (15)
C7	0.0424 (12)	0.0580 (16)	0.0503 (14)	0.0099 (12)	0.0127 (11)	−0.0033 (12)
C8	0.0495 (14)	0.0680 (18)	0.0720 (18)	0.0114 (13)	0.0139 (13)	−0.0164 (15)
C9	0.0475 (13)	0.0477 (14)	0.0550 (15)	0.0049 (11)	0.0163 (11)	−0.0086 (12)
C10	0.0524 (14)	0.0447 (14)	0.0690 (17)	−0.0029 (12)	0.0216 (12)	0.0032 (13)
C11	0.0513 (15)	0.0593 (17)	0.0721 (18)	0.0063 (13)	0.0137 (13)	0.0068 (14)

supplementary materials

C12	0.0458 (14)	0.0609 (17)	0.087 (2)	-0.0100 (13)	0.0207 (14)	-0.0153 (16)
C13	0.075 (2)	0.0485 (16)	0.093 (2)	-0.0114 (15)	0.0380 (17)	-0.0034 (16)
C14	0.0721 (18)	0.0514 (16)	0.0657 (18)	0.0113 (14)	0.0216 (14)	0.0065 (14)
F1	0.0526 (10)	0.0961 (14)	0.157 (2)	-0.0212 (10)	0.0171 (11)	-0.0227 (14)
O1	0.0564 (11)	0.0761 (14)	0.1190 (18)	-0.0081 (10)	0.0364 (11)	-0.0460 (13)
O2	0.0454 (9)	0.0655 (12)	0.0780 (12)	0.0031 (9)	0.0241 (8)	-0.0133 (10)
O3	0.0458 (10)	0.0736 (13)	0.0791 (13)	0.0107 (9)	0.0226 (9)	-0.0201 (10)

Geometric parameters (Å, °)

C1—C6	1.391 (3)	C8—H8A	0.9700
C1—C2	1.393 (3)	C8—H8B	0.9700
C1—C7	1.483 (3)	C9—C14	1.380 (4)
C2—C3	1.366 (3)	C9—C10	1.380 (3)
C2—H2A	0.9300	C10—C11	1.374 (3)
C3—O1	1.365 (3)	C10—H10A	0.9300
C3—C4	1.397 (3)	C11—C12	1.354 (4)
C4—O2	1.366 (3)	C11—H11A	0.9300
C4—C5	1.369 (4)	C12—F1	1.359 (3)
C5—C6	1.378 (3)	C12—C13	1.362 (4)
C5—H5A	0.9300	C13—C14	1.381 (4)
C6—H6A	0.9300	C13—H13A	0.9300
C7—O3	1.219 (3)	C14—H14A	0.9300
C7—C8	1.512 (4)	O1—H1A	0.8200
C8—C9	1.503 (3)	O2—H2B	0.8200
C6—C1—C2	119.2 (2)	C9—C8—H8B	108.3
C6—C1—C7	121.4 (2)	C7—C8—H8B	108.3
C2—C1—C7	119.4 (2)	H8A—C8—H8B	107.4
C3—C2—C1	120.5 (2)	C14—C9—C10	118.0 (2)
C3—C2—H2A	119.8	C14—C9—C8	121.6 (2)
C1—C2—H2A	119.8	C10—C9—C8	120.3 (2)
O1—C3—C2	119.6 (2)	C11—C10—C9	121.4 (2)
O1—C3—C4	120.3 (2)	C11—C10—H10A	119.3
C2—C3—C4	120.1 (2)	C9—C10—H10A	119.3
O2—C4—C5	124.3 (2)	C12—C11—C10	118.6 (3)
O2—C4—C3	116.2 (2)	C12—C11—H11A	120.7
C5—C4—C3	119.5 (2)	C10—C11—H11A	120.7
C4—C5—C6	120.9 (2)	C11—C12—F1	119.0 (3)
C4—C5—H5A	119.6	C11—C12—C13	122.6 (3)
C6—C5—H5A	119.6	F1—C12—C13	118.5 (3)
C5—C6—C1	119.9 (3)	C12—C13—C14	118.2 (3)
C5—C6—H6A	120.1	C12—C13—H13A	120.9
C1—C6—H6A	120.1	C14—C13—H13A	120.9
O3—C7—C1	121.2 (2)	C9—C14—C13	121.2 (3)
O3—C7—C8	120.5 (2)	C9—C14—H14A	119.4
C1—C7—C8	118.3 (2)	C13—C14—H14A	119.4
C9—C8—C7	115.7 (2)	C3—O1—H1A	109.5
C9—C8—H8A	108.3	C4—O2—H2B	109.5
C7—C8—H8A	108.3		

C6—C1—C2—C3	-1.2 (4)	C2—C1—C7—C8	-175.8 (2)
C7—C1—C2—C3	179.1 (2)	O3—C7—C8—C9	-8.8 (4)
C1—C2—C3—O1	178.1 (2)	C1—C7—C8—C9	169.9 (2)
C1—C2—C3—C4	0.9 (4)	C7—C8—C9—C14	112.3 (3)
O1—C3—C4—O2	2.9 (4)	C7—C8—C9—C10	-69.3 (3)
C2—C3—C4—O2	-179.9 (2)	C14—C9—C10—C11	0.3 (4)
O1—C3—C4—C5	-177.5 (3)	C8—C9—C10—C11	-178.2 (2)
C2—C3—C4—C5	-0.3 (4)	C9—C10—C11—C12	-0.1 (4)
O2—C4—C5—C6	179.6 (3)	C10—C11—C12—F1	-179.8 (3)
C3—C4—C5—C6	0.0 (4)	C10—C11—C12—C13	-0.3 (4)
C4—C5—C6—C1	-0.4 (4)	C11—C12—C13—C14	0.3 (5)
C2—C1—C6—C5	1.0 (4)	F1—C12—C13—C14	179.9 (3)
C7—C1—C6—C5	-179.4 (3)	C10—C9—C14—C13	-0.2 (4)
C6—C1—C7—O3	-176.9 (3)	C8—C9—C14—C13	178.3 (2)
C2—C1—C7—O3	2.8 (4)	C12—C13—C14—C9	-0.1 (4)
C6—C1—C7—C8	4.5 (4)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H1A...O2	0.82	2.28	2.690 (2)	111
O1—H1A...O2 ⁱ	0.82	2.16	2.876 (3)	146
O2—H2B...O3 ⁱⁱ	0.82	1.92	2.744 (2)	178

Symmetry codes: (i) $-x, -y, -z$; (ii) $x-1, y, z$.

Fig. 1

