

## 2,6-Dimethylanilinium chloride monohydrate

Wajda Smirani,\* Olfa Amri and Mohamed Rzaigui

Laboratoire de Chimie des Matériaux, Faculté des Sciences de Bizerte, 7021 Zarzouna Bizerte, Tunisia

Correspondence e-mail: wajda\_st@yahoo.fr


Received 20 November 2008; accepted 21 November 2008

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.031;  $wR$  factor = 0.080; data-to-parameter ratio = 14.4.

In the title hydrated molecular salt,  $\text{C}_8\text{H}_{12}\text{N}^+\cdot\text{Cl}^-\cdot\text{H}_2\text{O}$ , the component species interact by way of  $\text{N}-\text{H}\cdots\text{O}$ ,  $\text{N}-\text{H}\cdots\text{Cl}$  and  $\text{O}-\text{H}\cdots\text{Cl}$  hydrogen bonds, resulting in a three-dimensional network.

### Related literature

For related structures, see: Abid *et al.* (2007); Mrad *et al.* (2006). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



### Experimental

#### Crystal data



$M_r = 175.65$

Monoclinic,  $P\bar{2}_1/c$

$a = 8.676(3)\text{ \AA}$

$b = 14.144(3)\text{ \AA}$

$c = 7.913(6)\text{ \AA}$

$\beta = 101.88(5)^\circ$

$V = 950.2(8)\text{ \AA}^3$

$Z = 4$

Mo  $K\alpha$  radiation

$\mu = 0.35\text{ mm}^{-1}$

$T = 293(2)\text{ K}$

$0.20 \times 0.13 \times 0.10\text{ mm}$

### Data collection

Enraf–Nonius TurboCAD-4 diffractometer  
Absorption correction: none  
3722 measured reflections  
2244 independent reflections

1827 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.033$   
2 standard reflections  
frequency: 120 min  
intensity decay: 5%

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$   
 $wR(F^2) = 0.080$   
 $S = 1.04$   
2244 reflections

156 parameters  
H-atom parameters not refined  
 $\Delta\rho_{\text{max}} = 0.17\text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.24\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1···Cl1	0.90 (2)	2.41 (2)	3.305 (3)	173 (2)
O1—H2···Cl1 <sup>i</sup>	0.87 (3)	2.32 (3)	3.163 (3)	165 (2)
N1—H6···Cl1 <sup>ii</sup>	0.893 (18)	2.392 (18)	3.235 (3)	157.5 (15)
N1—H7···O1	0.896 (16)	1.835 (16)	2.731 (3)	177.3 (17)
N1—H8···Cl1 <sup>iii</sup>	0.883 (16)	2.414 (16)	3.265 (3)	162.8 (15)

Symmetry codes: (i)  $x, -y + \frac{1}{2}, z - \frac{1}{2}$ ; (ii)  $-x + 1, -y, -z + 1$ ; (iii)  $x, y, z - 1$ .

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2860).

### References

- Abid, S., Hemissi, H. & Rzaigui, M. (2007). *Acta Cryst. E63*, o3117.
- Bernstein, J., David, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Enraf–Nonius (1994). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- Mrad, M. L., Ben Nasr, C. & Rzaigui, M. (2006). *Anal. Sci.* **22**, x227–x228.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.

# supporting information

*Acta Cryst.* (2008). E64, o2463 [doi:10.1107/S1600536808039159]

## 2,6-Dimethylanilinium chloride monohydrate

**Wajda Smirani, Olfa Amri and Mohamed Rzaigui**

### S1. Comment

As part of our ongoing studies of organic-inorganic hybrid networks containing the 2,6-xylidinium cation (Mrad *et al.*, 2006; Abid *et al.*, 2007) we now report the synthesis and structure of the title compound, (I).

As shown in Fig. 1, the asymmetric unit of (I) contains a 2,6-xylidinium cation, a chloride anion and a water molecule. A perspective view of the structure along the *a* axis is given in Fig. 2. It shows that two 2,6-xylidinium cations are interconnected through two chloride anions into dimers *via* two N—H···Cl bonds, characterized by N···Cl separations of 3.264 (3) and 3.235 (3) Å and forming an 8-membered ring with graph-set  $R_2^4(8)$  (Bernstein *et al.*, 1995).

The title compound is a crystalline hydrate including one water of crystallization, which interconnect these dimers to each other to form layers parallel to the (*b*, *c*) plane, through N—H···O and O—H···Cl hydrogen bonds (Table 1).


Hydrogen bonds, electrostatic and van der Waals interactions participate to the cohesion of the three-dimensional network and add stability to this compound (Fig. 2). An examination of the organic group moiety geometrical features shows that the C—C and C—N bond lengths and the C—C—C and C—C—N angles are in the range usually found for this molecule (Abid *et al.*, 2007).

### S2. Experimental


2,6-xylidinie and HCl were mixed in water in a 1: 1 molar ratio. The obtained solution was slowly evaporated at room temperature to yield colourless blocks of (I).

### S3. Refinement

The H atoms were located in a difference map and their positions and  $U_{\text{iso}}$  values were freely refined.

**Figure 1**

View of (I) with displacement ellipsoids for non-H atoms drawn at the 30% probability level (arbitrary spheres for the H atoms).

**Figure 2**

A perspective view of the packing in (I).

### 2,6-Dimethylanilinium chloride monohydrate

#### Crystal data



$M_r = 175.65$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 8.676 (3) \text{ \AA}$

$b = 14.144 (3) \text{ \AA}$

$c = 7.913 (6) \text{ \AA}$

$\beta = 101.88 (5)^\circ$

$V = 950.2 (8) \text{ \AA}^3$

$Z = 4$

$F(000) = 376$

$D_x = 1.228 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 25 reflections

$\theta = 9.2\text{--}10.8^\circ$

$\mu = 0.35 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Block, colourless

$0.20 \times 0.13 \times 0.10 \text{ mm}$

*Data collection*

Enraf–Nonius TurboCAD-4  
diffractometer  
Radiation source: Enraf Nonius FR590  
Graphite monochromator  
Non-profiled  $\omega$  scans  
3722 measured reflections  
2244 independent reflections  
1827 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.033$   
 $\theta_{\max} = 28.0^\circ, \theta_{\min} = 2.8^\circ$   
 $h = -5 \rightarrow 11$   
 $k = -18 \rightarrow 0$   
 $l = -10 \rightarrow 10$   
2 standard reflections every 120 min  
intensity decay: 5%

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.031$   
 $wR(F^2) = 0.080$   
 $S = 1.04$   
2244 reflections  
156 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters not refined  
 $w = 1/[\sigma^2(F_o^2) + (0.0408P)^2 + 0.1063P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.17 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.24 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
H8	0.3326 (19)	0.0879 (11)	0.050 (2)	0.048 (4)*
H7	0.3721 (19)	0.1013 (11)	0.233 (2)	0.045 (4)*
H5	-0.159 (2)	0.2065 (13)	0.033 (2)	0.058 (4)*
H6	0.3494 (19)	0.0085 (13)	0.161 (2)	0.053 (4)*
H3	-0.121 (2)	-0.0291 (12)	0.318 (2)	0.055 (4)*
H13	0.171 (2)	0.2037 (14)	-0.109 (2)	0.066 (5)*
H11	0.265 (2)	-0.0493 (14)	0.427 (3)	0.070 (6)*
H17	0.056 (2)	0.2698 (15)	-0.066 (2)	0.073 (5)*
H2	0.488 (3)	0.2297 (19)	0.405 (3)	0.083 (7)*
H4	-0.278 (2)	0.0943 (13)	0.191 (2)	0.067 (5)*
H1	0.472 (3)	0.1562 (17)	0.518 (3)	0.090 (7)*
H10	0.215 (2)	-0.1099 (13)	0.268 (2)	0.062 (5)*
H9	0.124 (2)	-0.1045 (14)	0.406 (3)	0.071 (6)*
H12	0.215 (3)	0.2560 (15)	0.065 (3)	0.087 (7)*
N1	0.31170 (12)	0.06739 (8)	0.14878 (14)	0.0353 (2)
C7	0.1308 (2)	0.22609 (11)	-0.0155 (2)	0.0510 (3)

C8	0.1800 (2)	-0.07172 (12)	0.3454 (2)	0.0512 (3)
C1	0.14475 (13)	0.07516 (8)	0.15925 (14)	0.0322 (2)
C6	0.05793 (14)	0.15126 (9)	0.07822 (14)	0.0359 (3)
C5	-0.09996 (16)	0.15637 (10)	0.08975 (17)	0.0435 (3)
C2	0.08245 (14)	0.00745 (9)	0.25368 (15)	0.0367 (3)
C3	-0.07586 (16)	0.01607 (10)	0.26089 (17)	0.0445 (3)
C4	-0.16594 (15)	0.08936 (11)	0.18003 (18)	0.0466 (3)
O1	0.48995 (14)	0.16863 (9)	0.41224 (15)	0.0580 (3)
Cl1	0.45854 (4)	0.11499 (2)	0.81025 (4)	0.04716 (12)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N1	0.0319 (5)	0.0398 (5)	0.0364 (5)	0.0052 (4)	0.0125 (4)	0.0025 (4)
C7	0.0553 (9)	0.0467 (7)	0.0546 (8)	0.0130 (7)	0.0200 (7)	0.0119 (6)
C8	0.0580 (9)	0.0523 (8)	0.0493 (8)	0.0100 (7)	0.0253 (7)	0.0123 (7)
C1	0.0282 (5)	0.0398 (6)	0.0296 (5)	0.0024 (4)	0.0083 (4)	-0.0058 (4)
C6	0.0370 (6)	0.0389 (6)	0.0318 (5)	0.0050 (5)	0.0074 (4)	-0.0049 (4)
C5	0.0356 (6)	0.0504 (7)	0.0428 (6)	0.0102 (6)	0.0039 (5)	-0.0091 (6)
C2	0.0378 (6)	0.0416 (6)	0.0328 (5)	0.0007 (5)	0.0121 (4)	-0.0047 (5)
C3	0.0402 (7)	0.0524 (8)	0.0452 (6)	-0.0068 (6)	0.0186 (5)	-0.0077 (6)
C4	0.0298 (6)	0.0602 (8)	0.0508 (7)	-0.0003 (5)	0.0107 (5)	-0.0145 (6)
O1	0.0654 (7)	0.0543 (7)	0.0509 (6)	0.0072 (5)	0.0039 (5)	-0.0041 (5)
Cl1	0.04621 (19)	0.04719 (19)	0.0534 (2)	0.01013 (14)	0.02267 (14)	0.00643 (13)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

N1—C1	1.4718 (16)	C1—C2	1.3907 (17)
N1—H8	0.883 (18)	C1—C6	1.3921 (17)
N1—H7	0.896 (17)	C6—C5	1.3934 (18)
N1—H6	0.893 (18)	C5—C4	1.380 (2)
C7—C6	1.504 (2)	C5—H5	0.935 (18)
C7—H13	0.93 (2)	C2—C3	1.3917 (18)
C7—H17	0.92 (2)	C3—C4	1.374 (2)
C7—H12	0.96 (2)	C3—H3	0.918 (18)
C8—C2	1.498 (2)	C4—H4	1.00 (2)
C8—H11	0.93 (2)	O1—H2	0.87 (3)
C8—H10	0.92 (2)	O1—H1	0.90 (3)
C8—H9	0.88 (2)		
C1—N1—H8	114.1 (10)	C2—C1—N1	118.34 (11)
C1—N1—H7	110.5 (10)	C6—C1—N1	118.50 (11)
H8—N1—H7	106.5 (15)	C1—C6—C5	117.13 (12)
C1—N1—H6	114.0 (11)	C1—C6—C7	121.91 (11)
H8—N1—H6	105.3 (15)	C5—C6—C7	120.95 (12)
H7—N1—H6	105.9 (14)	C4—C5—C6	121.09 (13)
C6—C7—H13	114.5 (12)	C4—C5—H5	121.5 (11)
C6—C7—H17	110.9 (13)	C6—C5—H5	117.4 (11)

H13—C7—H17	103.3 (16)	C1—C2—C3	117.24 (12)
C6—C7—H12	108.9 (13)	C1—C2—C8	122.14 (12)
H13—C7—H12	108.2 (18)	C3—C2—C8	120.62 (12)
H17—C7—H12	111.0 (18)	C4—C3—C2	121.22 (13)
C2—C8—H11	111.7 (12)	C4—C3—H3	119.6 (11)
C2—C8—H10	110.5 (11)	C2—C3—H3	119.2 (11)
H11—C8—H10	109.9 (17)	C3—C4—C5	120.15 (12)
C2—C8—H9	109.7 (13)	C3—C4—H4	118.9 (11)
H11—C8—H9	104.3 (18)	C5—C4—H4	120.9 (11)
H10—C8—H9	110.5 (17)	H2—O1—H1	105 (2)
C2—C1—C6	123.14 (11)		
C2—C1—C6—C5	-2.12 (17)	N1—C1—C2—C3	-179.68 (10)
N1—C1—C6—C5	179.69 (10)	C6—C1—C2—C8	-177.60 (12)
C2—C1—C6—C7	176.85 (12)	N1—C1—C2—C8	0.60 (18)
N1—C1—C6—C7	-1.34 (17)	C1—C2—C3—C4	-0.96 (18)
C1—C6—C5—C4	0.94 (17)	C8—C2—C3—C4	178.78 (14)
C7—C6—C5—C4	-178.04 (13)	C2—C3—C4—C5	-0.1 (2)
C6—C1—C2—C3	2.13 (17)	C6—C5—C4—C3	0.1 (2)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1···Cl1	0.90 (2)	2.41 (2)	3.305 (3)	173 (2)
O1—H2···Cl1 <sup>i</sup>	0.87 (3)	2.32 (3)	3.163 (3)	165 (2)
N1—H6···Cl1 <sup>ii</sup>	0.893 (18)	2.392 (18)	3.235 (3)	157.5 (15)
N1—H7···O1	0.896 (16)	1.835 (16)	2.731 (3)	177.3 (17)
N1—H8···Cl1 <sup>iii</sup>	0.883 (16)	2.414 (16)	3.265 (3)	162.8 (15)

Symmetry codes: (i)  $x, -y+1/2, z-1/2$ ; (ii)  $-x+1, -y, -z+1$ ; (iii)  $x, y, z-1$ .