

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Phenyl naphthalene-2-sulfonate

Jasmine P. Vennila,^a Helen P. Kavitha,^b D. John Thiruvadigal^c and V. Manivannan^{d*}

^aDepartment of Physics, Panimalar Institute of Technology, Chennai 600 095, India,

^bDepartment of Chemistry, SRM University, Ramapuram, Chennai 600 089, India,

^cDepartment of Physics, SRM University, Kattankulathur Campus, Chennai 603 203, India, and ^dDepartment of Physics, Presidency College, Chennai 600 005, India

Correspondence e-mail: manivan_1999@yahoo.com

Received 24 October 2008; accepted 31 October 2008


Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å;

R factor = 0.044; wR factor = 0.106; data-to-parameter ratio = 28.1.

In the crystal structure of the title compound, $\text{C}_{16}\text{H}_{12}\text{O}_3\text{S}$, the dihedral angle between the naphthalene ring system and the phenyl ring is $65.21(3)^\circ$. The molecules are linked by intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, forming a chain along the a axis. The chains are connected through weak $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For general background, see: Spungin *et al.* (1984); Yachi *et al.* (1989). For related structures, see: Manivannan *et al.* (2005); Ramachandran *et al.* (2007).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{12}\text{O}_3\text{S}$

$M_r = 284.32$

Orthorhombic, $P2_12_12_1$

$a = 6.1525(2)$ Å

$b = 12.7466(7)$ Å

$c = 17.3414(10)$ Å

$V = 1359.97(12)$ Å³

$Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.24$ mm⁻¹

$T = 295(2)$ K
 $0.25 \times 0.18 \times 0.16$ mm

Data collection

Bruker Kappa APEXII diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.942$, $T_{\max} = 0.962$

11868 measured reflections
5093 independent reflections
3412 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.106$
 $S = 1.01$
5093 reflections
181 parameters
H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.26$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.31$ e Å⁻³
Absolute structure: Flack (1983),
1993 Friedel pairs
Flack parameter: $-0.03(7)$

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C2}-\text{H2}\cdots\text{O2}^{\text{i}}$	0.93	2.51	3.424 (2)	169
$\text{C5}-\text{H5}\cdots\text{Cg2}^{\text{ii}}$	0.93	2.96	3.486 (2)	117
$\text{C6}-\text{H6}\cdots\text{Cg3}^{\text{iii}}$	0.93	2.94	3.535 (2)	123
$\text{C12}-\text{H12}\cdots\text{Cg1}^{\text{iii}}$	0.93	2.94	3.788 (3)	152

Symmetry codes: (i) $x + 1, y, z$; (ii) $-x + \frac{3}{2}, -y + 2, z + \frac{1}{2}$; (iii) $-x, y + \frac{5}{2}, -z + \frac{1}{2}$. Cg1 is the centroid of the C1–C6 ring, Cg2 is the centroid of the C7–C9/C14–C16 ring and Cg3 is the centroid of the C9–C14 ring.

Data collection: APEX2 (Bruker, 2004); cell refinement: APEX2 and SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: PLATON (Spek, 2003); software used to prepare material for publication: SHELXL97.

The authors acknowledge the Sophisticated Analytical Instrument Facility, Indian Institute of Technology, Madras, for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2352).

References

- Bruker (2004). APEX2. Bruker AXS Inc., Madison, Wisconsin, USA.
Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
Manivannan, V., Vembu, N., Nallu, M., Sivakumar, K. & Fronczek, F. R. (2005). *Acta Cryst.* **E61**, o239–o241.
Ramachandran, G., Kanakam, C. C., Manivannan, V., Thiruvengatam, V. & Row, T. N. G. (2007). *Acta Cryst.* **E63**, o4638.
Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
Spungin, B., Levinshal, T., Rubenstein, S. & Breitbart, H. (1984). *Biochim. Biophys. Acta*, **769**, 531–542.
Yachi, K., Sugiyama, Y., Sawada, Y., Iga, T., Ikeda, Y., Toda, G. & Hananon, M. (1989). *Biochim. Biophys. Acta*, **978**, 1–7.

supplementary materials

Acta Cryst. (2008). E64, o2304 [doi:10.1107/S1600536808035824]

Phenyl naphthalene-2-sulfonate

J. P. Vennila, H. P. Kavitha, D. J. Thiruvadigal and V. Manivannan

Comment

Several compounds containing the *para*-toluene sulfonate moiety are used in the fields of biology and industry. The merging of lipids can be monitored using a derivative of *para*-toluene sulfonate (Yachi *et al.*, 1989). This method has been used in studying the membrane fusion during the acrosome reaction (Spungin *et al.*, 1984).

The geometric parameters in the title compound (I) agree well with the reported values of similar structures (Manivannan *et al.*, 2005; Ramachandran *et al.*, 2007). The phenyl ring makes a dihedral angle of 65.21 (3)° with the naphthalene ring system. The torsion angles of O2—S1—C7—C8 and O3—S1—C7—C16 [2.04 (15)° and 46.94 (15)°, respectively] indicate the *syn* conformation of the sulfonyl moiety. The crystal structure is stabilized through weak intermolecular C—H···O and C—H··· π interactions (Fig. 2 and Table 1). Cg1, Cg2 and Cg3 are the centroids of the C1—C6 ring, the C7—C9/C14—C16 ring and the C9—C14 ring, respectively.

Experimental

2-Naphthalene sulfonyl chloride (2.5 mmol) dissolved in acetone (4 ml) was added dropwise to phenol (2.5 mmol) in aqueous NaOH solution (4 ml, 5%) with constant shaking. The precipitated compound (2 mmol, yield 77%) recrystallized from ethanol yielded colourless crystals.

Refinement

H atoms were positioned geometrically and refined using riding model, with C—H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of (I), with atom labels and 50% probability displacement ellipsoids for non-H atoms.




Fig. 2. The packing of (I), viewed down the *b* axis. Hydrogen bonds are shown as dashed lines. H atoms not involved in hydrogen bonding have been omitted for the sake of clarity.

Phenyl naphthalene-2-sulfonate

Crystal data

$C_{16}H_{12}O_3S$	$F_{000} = 592$
$M_r = 284.32$	$D_x = 1.389 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
Hall symbol: P 2ac 2ab	$\lambda = 0.71073 \text{ \AA}$
$a = 6.1525 (2) \text{ \AA}$	Cell parameters from 4818 reflections
$b = 12.7466 (7) \text{ \AA}$	$\theta = 2.2\text{--}25.4^\circ$
$c = 17.3414 (10) \text{ \AA}$	$\mu = 0.24 \text{ mm}^{-1}$
$V = 1359.97 (12) \text{ \AA}^3$	$T = 295 (2) \text{ K}$
$Z = 4$	Tablet, colourless
	$0.25 \times 0.18 \times 0.16 \text{ mm}$

Data collection

Bruker Kappa APEXII diffractometer	5093 independent reflections
Radiation source: fine-focus sealed tube	3412 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.027$
$T = 295(2) \text{ K}$	$\theta_{\text{max}} = 34.9^\circ$
ω and φ scans	$\theta_{\text{min}} = 2.4^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -9 \rightarrow 5$
$T_{\text{min}} = 0.942$, $T_{\text{max}} = 0.962$	$k = -19 \rightarrow 18$
11868 measured reflections	$l = -25 \rightarrow 27$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.044$	$w = 1/[\sigma^2(F_o^2) + (0.0476P)^2 + 0.0596P]$
$wR(F^2) = 0.106$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.01$	$(\Delta/\sigma)_{\text{max}} < 0.001$
5093 reflections	$\Delta\rho_{\text{max}} = 0.26 \text{ e \AA}^{-3}$
181 parameters	$\Delta\rho_{\text{min}} = -0.31 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none
Secondary atom site location: difference Fourier map	Absolute structure: Flack (1983), 1993 Friedel pairs
	Flack parameter: $-0.03 (7)$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
-----	-----	-----	----------------------------------

S1	0.68361 (7)	0.89579 (3)	0.10630 (3)	0.03997 (11)
O1	0.8255 (2)	0.88973 (9)	0.18279 (6)	0.0403 (3)
O2	0.4945 (2)	0.95689 (12)	0.12058 (8)	0.0527 (4)
O3	0.6661 (3)	0.78918 (10)	0.08397 (9)	0.0646 (4)
C1	0.8523 (3)	0.98154 (12)	0.22766 (9)	0.0332 (3)
C2	1.0472 (3)	1.03331 (16)	0.22378 (11)	0.0438 (4)
H2	1.1557	1.0114	0.1901	0.053*
C3	1.0781 (3)	1.11934 (17)	0.27152 (13)	0.0537 (5)
H3	1.2089	1.1558	0.2701	0.064*
C4	0.9168 (3)	1.15065 (16)	0.32060 (12)	0.0513 (5)
H4	0.9389	1.2084	0.3524	0.062*
C5	0.7234 (3)	1.09789 (16)	0.32342 (10)	0.0497 (5)
H5	0.6144	1.1203	0.3567	0.060*
C6	0.6895 (3)	1.01146 (14)	0.27702 (10)	0.0422 (4)
H6	0.5594	0.9744	0.2792	0.051*
C7	0.8516 (3)	0.96307 (12)	0.04171 (9)	0.0334 (3)
C8	0.7855 (3)	1.05729 (12)	0.01241 (9)	0.0355 (4)
H8	0.6522	1.0855	0.0269	0.043*
C9	0.9203 (3)	1.11168 (13)	-0.03991 (10)	0.0380 (4)
C10	0.8568 (4)	1.20865 (14)	-0.07297 (12)	0.0522 (5)
H10	0.7231	1.2379	-0.0601	0.063*
C11	0.9892 (5)	1.25888 (17)	-0.12299 (12)	0.0652 (6)
H11	0.9458	1.3223	-0.1446	0.078*
C12	1.1911 (5)	1.2161 (2)	-0.14254 (12)	0.0709 (7)
H12	1.2808	1.2516	-0.1769	0.085*
C13	1.2580 (3)	1.12340 (18)	-0.11191 (12)	0.0569 (5)
H13	1.3933	1.0963	-0.1251	0.068*
C14	1.1230 (3)	1.06801 (14)	-0.06014 (10)	0.0405 (4)
C15	1.1837 (3)	0.96940 (15)	-0.02884 (10)	0.0439 (4)
H15	1.3164	0.9399	-0.0425	0.053*
C16	1.0520 (3)	0.91760 (13)	0.02050 (10)	0.0395 (4)
H16	1.0929	0.8527	0.0403	0.047*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0443 (2)	0.0406 (2)	0.0350 (2)	-0.00821 (18)	0.00007 (19)	-0.00279 (17)
O1	0.0525 (7)	0.0344 (5)	0.0339 (6)	0.0047 (6)	-0.0012 (5)	0.0021 (5)
O2	0.0361 (7)	0.0768 (9)	0.0451 (8)	-0.0027 (6)	0.0010 (6)	0.0017 (7)
O3	0.0897 (11)	0.0457 (7)	0.0584 (9)	-0.0245 (7)	0.0029 (9)	-0.0083 (6)
C1	0.0381 (10)	0.0353 (7)	0.0262 (7)	0.0012 (6)	-0.0017 (6)	0.0046 (6)
C2	0.0352 (10)	0.0579 (11)	0.0382 (9)	0.0016 (8)	0.0018 (7)	-0.0005 (8)
C3	0.0463 (12)	0.0649 (13)	0.0498 (12)	-0.0151 (10)	-0.0042 (9)	-0.0019 (10)
C4	0.0665 (14)	0.0508 (11)	0.0365 (11)	-0.0090 (9)	-0.0024 (9)	-0.0056 (8)
C5	0.0563 (13)	0.0562 (11)	0.0366 (9)	-0.0022 (9)	0.0132 (8)	-0.0060 (9)
C6	0.0413 (10)	0.0503 (9)	0.0348 (8)	-0.0094 (8)	0.0053 (8)	0.0016 (7)
C7	0.0373 (9)	0.0340 (7)	0.0290 (7)	-0.0004 (6)	-0.0011 (6)	-0.0049 (6)
C8	0.0374 (9)	0.0343 (7)	0.0348 (8)	0.0011 (7)	-0.0017 (7)	-0.0063 (6)

supplementary materials

C9	0.0478 (10)	0.0338 (8)	0.0325 (8)	-0.0053 (7)	-0.0064 (7)	-0.0034 (7)
C10	0.0675 (14)	0.0375 (9)	0.0516 (11)	-0.0044 (8)	-0.0138 (10)	0.0012 (8)
C11	0.0953 (18)	0.0482 (11)	0.0521 (14)	-0.0225 (12)	-0.0208 (13)	0.0102 (9)
C12	0.0927 (18)	0.0760 (15)	0.0440 (12)	-0.0440 (15)	-0.0050 (13)	0.0084 (11)
C13	0.0562 (12)	0.0748 (13)	0.0396 (10)	-0.0233 (10)	0.0020 (8)	-0.0062 (10)
C14	0.0419 (11)	0.0515 (10)	0.0280 (8)	-0.0102 (8)	-0.0035 (7)	-0.0074 (7)
C15	0.0378 (10)	0.0594 (10)	0.0347 (9)	0.0062 (9)	-0.0008 (8)	-0.0070 (8)
C16	0.0451 (10)	0.0425 (9)	0.0310 (9)	0.0101 (7)	-0.0023 (7)	-0.0024 (7)

Geometric parameters (Å, °)

S1—O3	1.4172 (13)	C7—C16	1.411 (2)
S1—O2	1.4217 (15)	C8—C9	1.411 (2)
S1—O1	1.5899 (13)	C8—H8	0.9300
S1—C7	1.7487 (17)	C9—C14	1.410 (3)
O1—C1	1.4149 (19)	C9—C10	1.417 (2)
C1—C2	1.370 (2)	C10—C11	1.351 (3)
C1—C6	1.372 (2)	C10—H10	0.9300
C2—C3	1.387 (3)	C11—C12	1.398 (4)
C2—H2	0.9300	C11—H11	0.9300
C3—C4	1.367 (3)	C12—C13	1.360 (3)
C3—H3	0.9300	C12—H12	0.9300
C4—C5	1.368 (3)	C13—C14	1.412 (3)
C4—H4	0.9300	C13—H13	0.9300
C5—C6	1.380 (3)	C14—C15	1.419 (3)
C5—H5	0.9300	C15—C16	1.351 (3)
C6—H6	0.9300	C15—H15	0.9300
C7—C8	1.366 (2)	C16—H16	0.9300
O3—S1—O2	120.72 (10)	C7—C8—C9	119.74 (16)
O3—S1—O1	102.89 (9)	C7—C8—H8	120.1
O2—S1—O1	109.31 (7)	C9—C8—H8	120.1
O3—S1—C7	109.88 (9)	C14—C9—C8	119.07 (16)
O2—S1—C7	109.05 (8)	C14—C9—C10	119.18 (18)
O1—S1—C7	103.52 (7)	C8—C9—C10	121.74 (17)
C1—O1—S1	118.86 (10)	C11—C10—C9	120.4 (2)
C2—C1—C6	122.41 (16)	C11—C10—H10	119.8
C2—C1—O1	118.25 (15)	C9—C10—H10	119.8
C6—C1—O1	119.20 (15)	C10—C11—C12	120.4 (2)
C1—C2—C3	118.10 (17)	C10—C11—H11	119.8
C1—C2—H2	121.0	C12—C11—H11	119.8
C3—C2—H2	121.0	C13—C12—C11	120.9 (2)
C4—C3—C2	120.22 (18)	C13—C12—H12	119.6
C4—C3—H3	119.9	C11—C12—H12	119.6
C2—C3—H3	119.9	C12—C13—C14	120.3 (2)
C3—C4—C5	120.68 (18)	C12—C13—H13	119.8
C3—C4—H4	119.7	C14—C13—H13	119.8
C5—C4—H4	119.7	C9—C14—C13	118.75 (19)
C4—C5—C6	120.22 (17)	C9—C14—C15	119.15 (17)
C4—C5—H5	119.9	C13—C14—C15	122.09 (19)

C6—C5—H5	119.9	C16—C15—C14	121.17 (18)
C1—C6—C5	118.37 (17)	C16—C15—H15	119.4
C1—C6—H6	120.8	C14—C15—H15	119.4
C5—C6—H6	120.8	C15—C16—C7	119.23 (16)
C8—C7—C16	121.62 (16)	C15—C16—H16	120.4
C8—C7—S1	119.58 (13)	C7—C16—H16	120.4
C16—C7—S1	118.80 (12)		
O3—S1—O1—C1	-171.58 (12)	S1—C7—C8—C9	-179.81 (12)
O2—S1—O1—C1	-42.13 (14)	C7—C8—C9—C14	-0.9 (2)
C7—S1—O1—C1	73.97 (13)	C7—C8—C9—C10	178.76 (16)
S1—O1—C1—C2	-104.08 (16)	C14—C9—C10—C11	-0.3 (3)
S1—O1—C1—C6	80.26 (17)	C8—C9—C10—C11	-179.92 (18)
C6—C1—C2—C3	-0.4 (3)	C9—C10—C11—C12	-0.3 (3)
O1—C1—C2—C3	-175.92 (16)	C10—C11—C12—C13	0.2 (3)
C1—C2—C3—C4	-0.1 (3)	C11—C12—C13—C14	0.6 (3)
C2—C3—C4—C5	0.0 (3)	C8—C9—C14—C13	-179.35 (16)
C3—C4—C5—C6	0.6 (3)	C10—C9—C14—C13	1.0 (2)
C2—C1—C6—C5	1.0 (3)	C8—C9—C14—C15	1.5 (2)
O1—C1—C6—C5	176.47 (15)	C10—C9—C14—C15	-178.11 (16)
C4—C5—C6—C1	-1.1 (3)	C12—C13—C14—C9	-1.2 (3)
O3—S1—C7—C8	132.35 (14)	C12—C13—C14—C15	177.93 (18)
O2—S1—C7—C8	-2.04 (15)	C9—C14—C15—C16	-0.8 (3)
O1—S1—C7—C8	-118.32 (13)	C13—C14—C15—C16	-179.89 (18)
O3—S1—C7—C16	-46.94 (15)	C14—C15—C16—C7	-0.6 (3)
O2—S1—C7—C16	178.67 (12)	C8—C7—C16—C15	1.3 (2)
O1—S1—C7—C16	62.39 (13)	S1—C7—C16—C15	-179.43 (14)
C16—C7—C8—C9	-0.5 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C2—H2 \cdots O2 ⁱ	0.93	2.51	3.424 (2)	169
C5—H5 \cdots Cg2 ⁱⁱ	0.93	2.96	3.486 (2)	117
C6—H6 \cdots Cg3 ⁱⁱ	0.93	2.94	3.535 (2)	123
C12—H12 \cdots Cg1 ⁱⁱⁱ	0.93	2.94	3.788 (3)	152

Symmetry codes: (i) $x+1, y, z$; (ii) $-x+3/2, -y+2, z+1/2$; (iii) $-x, y+5/2, -z+1/2$.

Fig. 1




Fig. 2

