

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

Tetra-*n*-butylammonium bis(1,1-dicyanoethylene-2,2-dithiolato)platinum(II)Nyasha Kanganga,<sup>a</sup> Kent R. Mann<sup>b</sup> and Daron E. Janzen<sup>a\*</sup>

<sup>a</sup>College of St Catherine, St Paul, Minnesota 55105, USA, and <sup>b</sup>Department of Chemistry, University of Minnesota, Minneapolis, Minnesota 55455, USA  
Correspondence e-mail: dejanzen@stkate.edu


Received 7 October 2008; accepted 7 November 2008

Key indicators: single-crystal X-ray study;  $T = 173$  K; mean  $\sigma(\text{C}-\text{C}) = 0.006$  Å;  
R factor = 0.027; wR factor = 0.077; data-to-parameter ratio = 17.3.

In the title compound,  $(\text{C}_{16}\text{H}_{36}\text{N})_2[\text{Pt}(\text{C}_4\text{N}_2\text{S}_2)_2]$ , the Pt<sup>II</sup> center adopts a distorted square-planar geometry due to the 4-membered chelate rings formed by coordination to the S atoms of the 1,1-dicyanoethylene-2,2-dithiolate (*i*-mnt) ligands [bite angle 74.35 (4)°]. The bond distances in the coordinated *i*-mnt ligands indicate some delocalization of the  $\pi$ -system.

## Related literature

For general background on the salts of metal complexes of  $[\text{Pt}(\textit{i}\text{-mnt})_2]^{2-}$  (*i*-mnt=1,1-dicyanoethylene-2,2-dithiolate), see: Cummings & Eisenberg (1996); Fackler & Coucouvanis (1966); Werden *et al.* (1966). For related structures, see: Gao *et al.* (2005, 2006); Hummel (1987); Li *et al.* (2004); Sun *et al.* (2006).



## Experimental

## Crystal data

 $(\text{C}_{16}\text{H}_{36}\text{N})_2[\text{Pt}(\text{C}_4\text{N}_2\text{S}_2)_2]$  $M_r = 960.40$ 

Monoclinic,  $P2_1/n$   
 $a = 9.8687$  (6) Å  
 $b = 16.9556$  (11) Å  
 $c = 13.8274$  (9) Å  
 $\beta = 92.840$  (1)°  
 $V = 2310.9$  (3) Å<sup>3</sup>

$Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 3.25$  mm<sup>-1</sup>  
 $T = 173$  (2) K  
 $0.40 \times 0.20 \times 0.15$  mm

## Data collection

Bruker SMART CCD area-detector diffractometer  
Absorption correction: multi-scan (SADABS; Bruker 2003)  
 $T_{\min} = 0.468$ ,  $T_{\max} = 0.612$

22256 measured reflections  
4087 independent reflections  
3226 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.033$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.027$   
 $wR(F^2) = 0.077$   
 $S = 1.09$   
4087 reflections

236 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 1.02$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.33$  e Å<sup>-3</sup>

Data collection: *SMART* (Bruker,2003); cell refinement: *SAINTE* (Bruker, 2006); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

This work was supported by funding from the NSF through a Research Site for Educators in Chemistry grant. The authors acknowledge Victor G. Young, Jr and the X-ray Crystallographic Laboratory in the Department of Chemistry at the University of Minnesota.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2127).

## References

- Bruker (2003). *SMART* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Bruker (2006). *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Cummings, S. D. & Eisenberg, R. (1996). *Inorg. Chim. Acta*, **242**, 225–231.  
Fackler, J. P. & Coucouvanis, D. (1966). *J. Am. Chem. Soc.* **88**, 3913–3920.  
Gao, X.-K., Dou, J.-M., Li, D.-C., Dong, F.-Y. & Wang, D.-Q. (2005). *J. Incl. Phen. Macro. Chem.* **53**, 111–119.  
Gao, X.-K., Dou, J.-M., Li, D.-C., Dong, F.-Y. & Wang, D.-Q. (2006). *J. Mol. Struct.* **733**, 181–186.  
Hummel, H.-U. (1987). *Transition Met. Chem.* **12**, 172–174.  
Li, B., Li, D.-C., Dong, F.-Y., Dou, J.-M. & Wang, D.-Q. (2004). *Z. Kristallogr.* **219**, 413–414.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Sun, Y.-M., Dong, F.-Y., Dou, J.-M., Li, D.-C., Gao, X.-K. & Wang, D.-Q. (2006). *J. Inorg. Organomet. Poly. Mater.* **16**, 61–67.  
Werden, B. G., Billig, E. & Gray, H. B. (1966). *Inorg. Chem.* **5**, 78–81.

**supplementary materials**

*Acta Cryst.* (2008). E64, m1552 [ doi:10.1107/S1600536808036726 ]

## Tetra-*n*-butylammonium bis(1,1-dicyanoethylene-2,2-dithiolato)platinum(II)

N. Kanganga, K. R. Mann and D. E. Janzen

### Comment

Salts of metal complexes of  $[\text{Pt}(i\text{-mnt})_2]^{2-}$  ( $i\text{-mnt}$ =1,1-dicyanoethylene-2,2-dithiolate) have been studied for their interesting electronic properties including their photoluminescence (Cummings & Eisenberg, 1996) and their redox behavior especially in relation to the analogous isomeric ligand 1,2-dicyanoethylene-1,2-dithiolate ( $\text{mnt}^{2-}$ ) complexes (Fackler & Coucouvanis, 1966; Werden *et al.*, 1966). In sharp contrast to  $\text{mnt}$  complexes of the form  $[\text{M}(\text{mnt})_2]^{2-}$  ( $\text{M} = \text{Ni}^{\text{II}}, \text{Pd}^{\text{II}}, \text{Pt}^{\text{II}}$ ) which do exhibit reversible oxidation behavior, analogous  $i\text{-mnt}$  complexes of the form  $[\text{M}(i\text{-mnt})_2]^{2-}$  do not. This effect is attributed to better  $\pi$ -delocalization of the five-membered rings formed by complexation of  $\text{mnt}$  compared with four-membered chelate rings of  $i\text{-mnt}$  complexes. Salts of  $[\text{Pt}(i\text{-mnt})_2]^{2-}$  have also been studied as supramolecular linker groups in organic-inorganic hybrid coordination polymers (Gao *et al.* 2005, 2006; Li *et al.* 2004; Sun *et al.* 2006). While several  $x$ -ray structures of  $[\text{Pt}(i\text{-mnt})_2]^{2-}$  with alkali metal-complexed crown ether salts have been reported, only one other simple non-coordinating cation salt (tetraethylammonium, Hummel 1987) has been structurally characterized.

The structure of the anion in the title compound  $(\text{C}_{16}\text{H}_{36}\text{N})_2[\text{Pt}(\text{S}_2\text{C}_4\text{N}_2)_2]$  shows significant distortions from a square planar environment as forced by the four-membered chelate rings of the  $i\text{-mnt}$  ligands, with the  $i\text{-mnt}$  bite angle  $\text{S}(2)\text{—Pt}(1)\text{—S}(1) = 74.35(4)^\circ$ . As the Pt sits on the special position  $(1/2, 1/2, 1/2)$  in the space group  $P2_1/n$ ,  $Z = 0.5$ , the anion is quite planar, with a calculated r.m.s. deviation from a least-squares plane formed by all atoms of the complex anion of  $0.042(3) \text{ \AA}$ . The bond lengths within coordinated  $i\text{-mnt}$  ligand, in particular the bonds  $\text{C}(1)\text{—C}(2) 1.361(6) \text{ \AA}$ ,  $\text{C}(2)\text{—C}(3) 1.429(6) \text{ \AA}$ , and  $\text{C}(2)\text{—C}(4) 1.430(6) \text{ \AA}$  are very similar to those observed in the tetraethylammonium salt, showing significant  $\pi$ -delocalization. No columnar stacking is observed amongst the complex anions. As expected, upon comparison of the structure of the title compound and the tetraethylammonium salt, little effect was observed on the intramolecular features of the complex anion.

### Experimental

The title compound  $(\text{C}_{16}\text{H}_{36}\text{N})_2[\text{Pt}(\text{S}_2\text{C}_4\text{N}_2)_2]$  was prepared using a procedure similar to that described by Fackler and Coucouvanis (1966) substituting the use of tetra-*n*-propylammonium iodide with tetra-*n*-butylammonium bromide. The title compound has been previously characterized by Werden *et al.* (1966). Spectroscopic analysis of the present sample obtained by this procedure was consistent with the data previously reported. Crystals were obtained by diffusion of diethyl ether into a concentrated solution of the title compound dissolved in dichloromethane.

### Refinement

The H atoms were geometrically placed ( $\text{C—H} = 0.98\text{--}0.99 \text{ \AA}$ ) and refined as riding with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  or  $1.5U_{\text{eq}}(\text{methyl C})$ .

## Figures




Fig. 1. The molecular structure of  $(C_{16}H_{36}N)_2[Pt(S_2C_4N_2)_2]$  showing 50% displacement ellipsoids for the non-hydrogen atoms. Only the crystallographically independent atoms are labelled.

## Tetra-*n*-butylammonium bis(1,1-dicyanoethylene-2,2-dithiolato)platinum(II)

### Crystal data

$(C_{16}H_{36}N)_2[Pt(C_4N_2S_2)_2]$

$M_r = 960.40$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 9.8687$  (6) Å

$b = 16.9556$  (11) Å

$c = 13.8274$  (9) Å

$\beta = 92.840$  (1)°

$V = 2310.9$  (3) Å<sup>3</sup>

$Z = 2$

$F(000) = 992$

$D_x = 1.380$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71069$  Å

Cell parameters from 3207 reflections

$\theta = 5.5$ – $50.0$ °

$\mu = 3.25$  mm<sup>-1</sup>

$T = 173$  K

Plate, yellow

$0.4 \times 0.2 \times 0.15$  mm

### Data collection

Bruker SMART CCD area-detector diffractometer

Radiation source: normal-focus sealed tube graphite

$\varphi$  scans

Absorption correction: multi-scan (SADABS; Bruker 2003)

$T_{\min} = 0.468$ ,  $T_{\max} = 0.612$

22256 measured reflections

4087 independent reflections

3226 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.033$

$\theta_{\max} = 25.1$ °,  $\theta_{\min} = 1.9$ °

$h = -11 \rightarrow 11$

$k = -20 \rightarrow 20$

$l = -16 \rightarrow 16$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.027$

$wR(F^2) = 0.077$

$S = 1.09$

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0339P)^2 + 3.6638P]$

where  $P = (F_o^2 + 2F_c^2)/3$

4087 reflections	$(\Delta/\sigma)_{\max} = 0.001$
236 parameters	$\Delta\rho_{\max} = 1.02 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.33 \text{ e } \text{\AA}^{-3}$
0 constraints	

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Pt1	0.5000	0.5000	0.5000	0.03643 (9)
S1	0.56122 (11)	0.43791 (7)	0.64610 (8)	0.0434 (3)
S2	0.30447 (11)	0.49674 (7)	0.58620 (8)	0.0439 (2)
C1	0.3957 (4)	0.4489 (2)	0.6793 (3)	0.0365 (9)
C2	0.3440 (4)	0.4241 (2)	0.7636 (3)	0.0395 (9)
C3	0.2028 (5)	0.4321 (2)	0.7797 (3)	0.0432 (10)
C4	0.4294 (5)	0.3889 (2)	0.8383 (3)	0.0419 (10)
N1	0.0899 (4)	0.4381 (3)	0.7927 (3)	0.0585 (11)
N2	0.4951 (5)	0.3608 (2)	0.8982 (3)	0.0558 (10)
N3	0.6156 (3)	0.15041 (17)	0.7976 (2)	0.0311 (7)
C5	0.6718 (4)	0.0849 (2)	0.7366 (3)	0.0325 (8)
H5A	0.5998	0.0449	0.7250	0.039*
H5B	0.7471	0.0592	0.7747	0.039*
C6	0.7239 (5)	0.1092 (3)	0.6392 (3)	0.0513 (12)
H6A	0.6473	0.1284	0.5965	0.062*
H6B	0.7899	0.1528	0.6485	0.062*
C7	0.7915 (5)	0.0399 (3)	0.5914 (3)	0.0530 (12)
H7A	0.8370	0.0594	0.5338	0.064*
H7B	0.8626	0.0185	0.6372	0.064*
C8	0.6986 (7)	-0.0250 (4)	0.5610 (5)	0.088 (2)
H8A	0.7497	-0.0664	0.5293	0.131*
H8B	0.6276	-0.0047	0.5156	0.131*
H8C	0.6568	-0.0470	0.6180	0.131*
C9	0.7213 (4)	0.2133 (2)	0.8215 (3)	0.0364 (9)
H9A	0.6795	0.2539	0.8620	0.044*
H9B	0.7446	0.2391	0.7603	0.044*
C10	0.8523 (4)	0.1859 (2)	0.8736 (3)	0.0407 (10)
H10A	0.8317	0.1624	0.9368	0.049*

## supplementary materials

---

H10B	0.8954	0.1448	0.8346	0.049*
C11	0.9494 (5)	0.2548 (3)	0.8897 (3)	0.0454 (10)
H11A	0.9651	0.2796	0.8264	0.054*
H11B	0.9063	0.2947	0.9305	0.054*
C12	1.0841 (5)	0.2320 (3)	0.9372 (4)	0.0548 (12)
H12A	1.1439	0.2781	0.9408	0.082*
H12B	1.1256	0.1905	0.8991	0.082*
H12C	1.0706	0.2124	1.0027	0.082*
C13	0.5705 (4)	0.1106 (2)	0.8892 (3)	0.0321 (8)
H13A	0.5059	0.0680	0.8703	0.039*
H13B	0.6507	0.0855	0.9223	0.039*
C14	0.5040 (5)	0.1638 (2)	0.9614 (3)	0.0416 (10)
H14A	0.5685	0.2056	0.9834	0.050*
H14B	0.4234	0.1896	0.9297	0.050*
C15	0.4618 (4)	0.1166 (2)	1.0474 (3)	0.0389 (9)
H15A	0.5434	0.0922	1.0794	0.047*
H15B	0.4007	0.0736	1.0242	0.047*
C16	0.3900 (5)	0.1654 (3)	1.1214 (3)	0.0499 (11)
H16A	0.3599	0.1309	1.1729	0.075*
H16B	0.3112	0.1916	1.0897	0.075*
H16C	0.4526	0.2052	1.1492	0.075*
C17	0.4978 (4)	0.1918 (2)	0.7443 (3)	0.0354 (9)
H17A	0.4689	0.2363	0.7848	0.042*
H17B	0.5309	0.2145	0.6839	0.042*
C18	0.3751 (4)	0.1418 (2)	0.7184 (3)	0.0395 (9)
H18A	0.3417	0.1178	0.7780	0.047*
H18B	0.4014	0.0985	0.6750	0.047*
C19	0.2614 (5)	0.1896 (3)	0.6683 (3)	0.0517 (12)
H19A	0.2422	0.2362	0.7084	0.062*
H19B	0.2914	0.2086	0.6052	0.062*
C20	0.1329 (5)	0.1419 (3)	0.6520 (4)	0.0549 (12)
H20A	0.0649	0.1735	0.6153	0.082*
H20B	0.0979	0.1273	0.7146	0.082*
H20C	0.1526	0.0941	0.6154	0.082*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Pt1	0.03067 (12)	0.04558 (14)	0.03345 (13)	-0.00168 (10)	0.00587 (8)	-0.00881 (11)
S1	0.0354 (5)	0.0561 (6)	0.0392 (6)	0.0020 (5)	0.0066 (4)	-0.0062 (5)
S2	0.0336 (5)	0.0607 (7)	0.0379 (5)	0.0017 (5)	0.0067 (4)	-0.0022 (5)
C1	0.037 (2)	0.033 (2)	0.040 (2)	0.0008 (17)	0.0046 (17)	-0.0123 (17)
C2	0.040 (2)	0.034 (2)	0.045 (2)	0.0013 (18)	0.0059 (19)	-0.0068 (18)
C3	0.049 (3)	0.039 (2)	0.043 (2)	-0.001 (2)	0.009 (2)	-0.0047 (19)
C4	0.053 (3)	0.032 (2)	0.043 (3)	-0.0012 (19)	0.014 (2)	-0.0088 (19)
N1	0.045 (2)	0.063 (3)	0.069 (3)	0.001 (2)	0.022 (2)	0.004 (2)
N2	0.071 (3)	0.047 (2)	0.050 (2)	0.006 (2)	0.008 (2)	0.0006 (19)
N3	0.0361 (17)	0.0295 (16)	0.0282 (17)	0.0022 (13)	0.0071 (13)	0.0044 (13)

C5	0.038 (2)	0.0298 (19)	0.031 (2)	0.0043 (16)	0.0081 (16)	-0.0007 (15)
C6	0.067 (3)	0.047 (3)	0.041 (3)	0.003 (2)	0.024 (2)	0.007 (2)
C7	0.060 (3)	0.056 (3)	0.045 (3)	0.002 (2)	0.026 (2)	0.000 (2)
C8	0.080 (4)	0.102 (5)	0.083 (4)	-0.016 (4)	0.034 (4)	-0.046 (4)
C9	0.043 (2)	0.030 (2)	0.037 (2)	-0.0031 (17)	0.0095 (18)	0.0057 (17)
C10	0.041 (2)	0.038 (2)	0.043 (2)	-0.0033 (18)	0.0023 (19)	0.0032 (18)
C11	0.053 (3)	0.041 (2)	0.043 (2)	-0.007 (2)	0.000 (2)	0.0058 (19)
C12	0.054 (3)	0.057 (3)	0.052 (3)	-0.011 (2)	-0.008 (2)	-0.004 (2)
C13	0.039 (2)	0.030 (2)	0.028 (2)	0.0020 (16)	0.0077 (16)	0.0059 (15)
C14	0.053 (3)	0.033 (2)	0.039 (2)	0.0041 (18)	0.0141 (19)	0.0023 (18)
C15	0.047 (2)	0.042 (2)	0.028 (2)	0.0022 (19)	0.0080 (18)	-0.0009 (17)
C16	0.055 (3)	0.055 (3)	0.040 (3)	0.003 (2)	0.016 (2)	0.000 (2)
C17	0.041 (2)	0.034 (2)	0.032 (2)	0.0081 (17)	0.0054 (17)	0.0061 (16)
C18	0.042 (2)	0.039 (2)	0.037 (2)	0.0074 (18)	0.0019 (18)	0.0018 (18)
C19	0.049 (3)	0.055 (3)	0.051 (3)	0.012 (2)	-0.001 (2)	0.008 (2)
C20	0.044 (3)	0.068 (3)	0.052 (3)	0.012 (2)	-0.003 (2)	0.001 (2)

*Geometric parameters (Å, °)*

Pt1—S2	2.3184 (10)	C10—H10B	0.9900
Pt1—S2 <sup>i</sup>	2.3185 (10)	C11—C12	1.504 (6)
Pt1—S1	2.3310 (11)	C11—H11A	0.9900
Pt1—S1 <sup>i</sup>	2.3310 (11)	C11—H11B	0.9900
S1—C1	1.729 (4)	C12—H12A	0.9800
S2—C1	1.735 (4)	C12—H12B	0.9800
C1—C2	1.361 (6)	C12—H12C	0.9800
C2—C3	1.429 (6)	C13—C14	1.519 (5)
C2—C4	1.430 (6)	C13—H13A	0.9900
C3—N1	1.142 (5)	C13—H13B	0.9900
C4—N2	1.132 (6)	C14—C15	1.509 (6)
N3—C9	1.516 (5)	C14—H14A	0.9900
N3—C5	1.516 (4)	C14—H14B	0.9900
N3—C17	1.518 (5)	C15—C16	1.517 (6)
N3—C13	1.522 (4)	C15—H15A	0.9900
C5—C6	1.522 (5)	C15—H15B	0.9900
C5—H5A	0.9900	C16—H16A	0.9800
C5—H5B	0.9900	C16—H16B	0.9800
C6—C7	1.518 (6)	C16—H16C	0.9800
C6—H6A	0.9900	C17—C18	1.508 (6)
C6—H6B	0.9900	C17—H17A	0.9900
C7—C8	1.480 (8)	C17—H17B	0.9900
C7—H7A	0.9900	C18—C19	1.522 (6)
C7—H7B	0.9900	C18—H18A	0.9900
C8—H8A	0.9800	C18—H18B	0.9900
C8—H8B	0.9800	C19—C20	1.511 (7)
C8—H8C	0.9800	C19—H19A	0.9900
C9—C10	1.521 (6)	C19—H19B	0.9900
C9—H9A	0.9900	C20—H20A	0.9800
C9—H9B	0.9900	C20—H20B	0.9800

## supplementary materials

---

C10—C11	1.521 (6)	C20—H20C	0.9800
C10—H10A	0.9900		
S2—Pt1—S2 <sup>i</sup>	180.0	C12—C11—H11A	108.8
S2—Pt1—S1	74.35 (4)	C10—C11—H11A	108.8
S2 <sup>i</sup> —Pt1—S1	105.65 (4)	C12—C11—H11B	108.8
S2—Pt1—S1 <sup>i</sup>	105.65 (4)	C10—C11—H11B	108.8
S2 <sup>i</sup> —Pt1—S1 <sup>i</sup>	74.35 (4)	H11A—C11—H11B	107.7
S1—Pt1—S1 <sup>i</sup>	180.0	C11—C12—H12A	109.5
C1—S1—Pt1	88.49 (15)	C11—C12—H12B	109.5
C1—S2—Pt1	88.74 (14)	H12A—C12—H12B	109.5
C2—C1—S1	126.4 (3)	C11—C12—H12C	109.5
C2—C1—S2	125.2 (3)	H12A—C12—H12C	109.5
S1—C1—S2	108.4 (2)	H12B—C12—H12C	109.5
C1—C2—C3	120.9 (4)	C14—C13—N3	115.9 (3)
C1—C2—C4	121.0 (4)	C14—C13—H13A	108.3
C3—C2—C4	118.1 (4)	N3—C13—H13A	108.3
N1—C3—C2	179.6 (5)	C14—C13—H13B	108.3
N2—C4—C2	178.8 (5)	N3—C13—H13B	108.3
C9—N3—C5	111.7 (3)	H13A—C13—H13B	107.4
C9—N3—C17	106.4 (3)	C15—C14—C13	110.3 (3)
C5—N3—C17	111.2 (3)	C15—C14—H14A	109.6
C9—N3—C13	110.9 (3)	C13—C14—H14A	109.6
C5—N3—C13	105.6 (3)	C15—C14—H14B	109.6
C17—N3—C13	111.2 (3)	C13—C14—H14B	109.6
N3—C5—C6	116.3 (3)	H14A—C14—H14B	108.1
N3—C5—H5A	108.2	C14—C15—C16	113.5 (4)
C6—C5—H5A	108.2	C14—C15—H15A	108.9
N3—C5—H5B	108.2	C16—C15—H15A	108.9
C6—C5—H5B	108.2	C14—C15—H15B	108.9
H5A—C5—H5B	107.4	C16—C15—H15B	108.9
C7—C6—C5	110.5 (4)	H15A—C15—H15B	107.7
C7—C6—H6A	109.5	C15—C16—H16A	109.5
C5—C6—H6A	109.5	C15—C16—H16B	109.5
C7—C6—H6B	109.5	H16A—C16—H16B	109.5
C5—C6—H6B	109.5	C15—C16—H16C	109.5
H6A—C6—H6B	108.1	H16A—C16—H16C	109.5
C8—C7—C6	114.8 (5)	H16B—C16—H16C	109.5
C8—C7—H7A	108.6	C18—C17—N3	116.2 (3)
C6—C7—H7A	108.6	C18—C17—H17A	108.2
C8—C7—H7B	108.6	N3—C17—H17A	108.2
C6—C7—H7B	108.6	C18—C17—H17B	108.2
H7A—C7—H7B	107.5	N3—C17—H17B	108.2
C7—C8—H8A	109.5	H17A—C17—H17B	107.4
C7—C8—H8B	109.5	C17—C18—C19	112.0 (3)
H8A—C8—H8B	109.5	C17—C18—H18A	109.2
C7—C8—H8C	109.5	C19—C18—H18A	109.2
H8A—C8—H8C	109.5	C17—C18—H18B	109.2
H8B—C8—H8C	109.5	C19—C18—H18B	109.2

N3—C9—C10	116.6 (3)	H18A—C18—H18B	107.9
N3—C9—H9A	108.1	C20—C19—C18	112.2 (4)
C10—C9—H9A	108.1	C20—C19—H19A	109.2
N3—C9—H9B	108.1	C18—C19—H19A	109.2
C10—C9—H9B	108.1	C20—C19—H19B	109.2
H9A—C9—H9B	107.3	C18—C19—H19B	109.2
C11—C10—C9	110.5 (3)	H19A—C19—H19B	107.9
C11—C10—H10A	109.6	C19—C20—H20A	109.5
C9—C10—H10A	109.6	C19—C20—H20B	109.5
C11—C10—H10B	109.6	H20A—C20—H20B	109.5
C9—C10—H10B	109.6	C19—C20—H20C	109.5
H10A—C10—H10B	108.1	H20A—C20—H20C	109.5
C12—C11—C10	113.8 (4)	H20B—C20—H20C	109.5
S2—Pt1—S1—C1	0.38 (13)	C5—C6—C7—C8	67.3 (6)
S2 <sup>i</sup> —Pt1—S1—C1	-179.62 (13)	C5—N3—C9—C10	-57.1 (4)
S1—Pt1—S2—C1	-0.38 (13)	C17—N3—C9—C10	-178.6 (3)
S1 <sup>i</sup> —Pt1—S2—C1	179.62 (13)	C13—N3—C9—C10	60.4 (4)
Pt1—S1—C1—C2	179.1 (4)	N3—C9—C10—C11	178.2 (3)
Pt1—S1—C1—S2	-0.52 (17)	C9—C10—C11—C12	-177.5 (4)
Pt1—S2—C1—C2	-179.1 (3)	C9—N3—C13—C14	61.4 (4)
Pt1—S2—C1—S1	0.52 (17)	C5—N3—C13—C14	-177.5 (3)
S1—C1—C2—C3	-175.8 (3)	C17—N3—C13—C14	-56.8 (4)
S2—C1—C2—C3	3.8 (6)	N3—C13—C14—C15	178.8 (3)
S1—C1—C2—C4	3.9 (6)	C13—C14—C15—C16	-178.0 (4)
S2—C1—C2—C4	-176.6 (3)	C9—N3—C17—C18	-176.2 (3)
C9—N3—C5—C6	-59.8 (4)	C5—N3—C17—C18	62.0 (4)
C17—N3—C5—C6	58.9 (4)	C13—N3—C17—C18	-55.3 (4)
C13—N3—C5—C6	179.6 (4)	N3—C17—C18—C19	178.0 (3)
N3—C5—C6—C7	173.5 (4)	C17—C18—C19—C20	-173.8 (4)

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ .

Fig. 1

