

Bis(isopropoxido- κ O)bis(2-methylquinolin-8-olato- κ^2 N,O)titanium(IV)

 Yousef Fazaeli,^a Mostafa M. Amini^a and Seik Weng Ng^{b*}
^aDepartment of Chemistry, Shahid Beheshti University, Tehran, Iran, and

^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia


Correspondence e-mail: seikweng@um.edu.my

Received 29 October 2008; accepted 29 October 2008

 Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.032; wR factor = 0.088; data-to-parameter ratio = 18.0.

 The two 2-methylquinolin-8-olate anions in the title complex, $[\text{Ti}(\text{C}_{10}\text{H}_8\text{NO})_2(\text{C}_3\text{H}_7\text{O})_2]$, chelate the Ti^{IV} atom, which shows an all-*cis* distorted octahedral N_2O_4 coordination geometry.

Related literature

 For the synthesis, see: Bickley & Nick (1979); Harrod & Taylor (1975). For the crystal structure of bis(isopropoxy)bis(quinolin-8-olato)titanium, see: Zeng *et al.* (2002).


Experimental

Crystal data

 $[\text{Ti}(\text{C}_{10}\text{H}_8\text{NO})_2(\text{C}_3\text{H}_7\text{O})_2]$
 $M_r = 482.42$

 Monoclinic, $P2_1/c$
 $a = 9.5851$ (2) Å

 $b = 13.5768$ (2) Å

 $c = 18.7779$ (3) Å

 $\beta = 102.559$ (1)°

 $V = 2385.19$ (7) Å³
 $Z = 4$

 Mo $K\alpha$ radiation

 $\mu = 0.39$ mm⁻¹
 $T = 100$ (2) K

 $0.35 \times 0.25 \times 0.15$ mm

Data collection

Bruker SMART APEX

diffractometer

Absorption correction: multi-scan

(SADABS; Sheldrick, 1996)

 $T_{\text{min}} = 0.875$, $T_{\text{max}} = 0.943$

16391 measured reflections

5467 independent reflections

 4651 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.088$
 $S = 1.03$

5467 reflections

304 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.39$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.39$ e Å⁻³

Data collection: APEX2 (Bruker, 2007); cell refinement: APEX2; data reduction: SAINT (Bruker, 2007); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2008).

The authors thank the Vice-President's Office for Research Affairs of Shahid Beheshti University and the University of Malaya for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2320).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bickley, D. G. & Nick, S. (1979). *Inorg. Chem.* **18**, 2200–2204.
 Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Harrod, J. F. & Taylor, K. R. (1975). *Inorg. Chem.* **14**, 1541–1545.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2008). publCIF. In preparation.
 Zeng, W. F., Chen, Y. S., Chiang, M. Y., Chern, S. S. & Cheng, C. P. (2002). *Polyhedron*, **21**, 1081–1087.

supplementary materials

Acta Cryst. (2008). E64, m1509 [doi:10.1107/S1600536808035460]

Bis(isopropoxido- κO)bis(2-methylquinolin-8-olato- $\kappa^2 N, O$)titanium(IV)

Y. Fazaeli, M. M. Amini and S. W. Ng

Experimental

8-Hydroxy-2-methylquinoline (1.59 g, 10 mmol) was added to the titanium isopropoxide (2.84 g, 10 mmol) in toluene (20 ml) at room temperature. The mixture was stirred for a day and then solvent was removed under reduced pressure to furnish an orange solid. The solid was crystallized from dichloromethane and *n*-hexane (1:1) to give yellow crystals, m.p. 445 K. IR (KBr, cm^{-1}): 1575 (C=C, C=N), 1236 (C—O). ^1H NMR (CDCl_3 , p.p.m.): 0.94 (CH_3 , doublet), 1.14 (CH_3 , doublet), 2.83 (CH_3 , singlet), 4.61 (CH, quartet), 6.9–8.58 (aromatic H atoms).

Refinement

Hydrogen atoms were placed in their calculated positions (C—H 0.95–0.98 Å) and were treated as riding on their parent atoms, with $U(\text{H})$ set to 1.2–1.5 times $U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of $\text{Ti}(\text{C}_{10}\text{H}_8\text{NO})_2(\text{C}_3\text{H}_7\text{O})_2$ at the 70% probability level showing atom labelling. Hydrogen atoms are drawn as spheres of arbitrary radius.

Bis(isopropoxido- κO)bis(2-methylquinolin-8-olato- $\kappa^2 N, O$)titanium(IV)

Crystal data

$[\text{Ti}(\text{C}_{10}\text{H}_8\text{NO})_2(\text{C}_3\text{H}_7\text{O})_2]$

$M_r = 482.42$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 9.5851$ (2) Å

$b = 13.5768$ (2) Å

$c = 18.7779$ (3) Å

$\beta = 102.559$ (1)°

$V = 2385.19$ (7) Å³

$F_{000} = 1016$

$D_x = 1.343$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 6457 reflections

$\theta = 2.2$ – 28.3 °

$\mu = 0.39$ mm⁻¹

$T = 100$ (2) K

Irregular chip, yellow

$0.35 \times 0.25 \times 0.15$ mm

supplementary materials

Z = 4

Data collection

Bruker SMART APEX diffractometer	5467 independent reflections
Radiation source: fine-focus sealed tube	4651 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.025$
$T = 100(2)$ K	$\theta_{\text{max}} = 27.5^\circ$
ω scans	$\theta_{\text{min}} = 2.2^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -12 \rightarrow 11$
$T_{\text{min}} = 0.875$, $T_{\text{max}} = 0.943$	$k = -17 \rightarrow 17$
16391 measured reflections	$l = -24 \rightarrow 24$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.032$	H-atom parameters constrained
$wR(F^2) = 0.088$	$w = 1/[\sigma^2(F_o^2) + (0.0394P)^2 + 1.3426P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
5467 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
304 parameters	$\Delta\rho_{\text{max}} = 0.39 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.39 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Ti1	0.36712 (3)	0.433571 (18)	0.241664 (14)	0.01288 (8)
O1	0.26537 (11)	0.55637 (7)	0.23824 (6)	0.0166 (2)
O2	0.46090 (11)	0.31617 (7)	0.28355 (5)	0.0153 (2)
O3	0.49414 (11)	0.48288 (8)	0.19514 (6)	0.0187 (2)
O4	0.24248 (11)	0.37144 (8)	0.16802 (5)	0.0171 (2)
N1	0.19366 (13)	0.41035 (9)	0.31601 (6)	0.0142 (2)
N2	0.51257 (13)	0.48222 (9)	0.35581 (6)	0.0145 (2)
C1	0.16171 (16)	0.57482 (11)	0.27306 (8)	0.0169 (3)
C2	0.09311 (17)	0.66473 (12)	0.27024 (9)	0.0219 (3)

H2	0.1206	0.7172	0.2427	0.026*
C3	-0.01715 (18)	0.67913 (12)	0.30789 (9)	0.0261 (4)
H3	-0.0628	0.7415	0.3055	0.031*
C4	-0.05984 (17)	0.60535 (13)	0.34771 (9)	0.0259 (4)
H4	-0.1351	0.6164	0.3724	0.031*
C5	0.00847 (16)	0.51221 (12)	0.35212 (8)	0.0204 (3)
C6	0.12004 (15)	0.49752 (11)	0.31484 (8)	0.0155 (3)
C7	-0.02862 (17)	0.43106 (13)	0.39098 (9)	0.0244 (3)
H7	-0.1022	0.4370	0.4174	0.029*
C8	0.04179 (17)	0.34421 (13)	0.39039 (9)	0.0228 (3)
H8	0.0155	0.2890	0.4157	0.027*
C9	0.15426 (16)	0.33490 (11)	0.35243 (8)	0.0170 (3)
C10	0.22910 (17)	0.23783 (11)	0.35446 (9)	0.0210 (3)
H10A	0.2668	0.2303	0.3102	0.032*
H10B	0.1615	0.1845	0.3570	0.032*
H10C	0.3081	0.2352	0.3975	0.032*
C11	0.54608 (15)	0.31020 (11)	0.34980 (8)	0.0151 (3)
C12	0.60985 (17)	0.22382 (12)	0.37911 (8)	0.0197 (3)
H12	0.5961	0.1647	0.3514	0.024*
C13	0.69565 (17)	0.22374 (12)	0.45060 (9)	0.0226 (3)
H13	0.7381	0.1638	0.4706	0.027*
C14	0.71930 (17)	0.30763 (12)	0.49184 (9)	0.0215 (3)
H14	0.7760	0.3054	0.5401	0.026*
C15	0.65868 (16)	0.39776 (12)	0.46212 (8)	0.0170 (3)
C16	0.57156 (15)	0.39851 (11)	0.39100 (8)	0.0145 (3)
C17	0.68312 (16)	0.49006 (12)	0.49736 (8)	0.0199 (3)
H17	0.7367	0.4940	0.5462	0.024*
C18	0.62969 (17)	0.57309 (11)	0.46108 (8)	0.0197 (3)
H18	0.6493	0.6353	0.4842	0.024*
C19	0.54487 (16)	0.56820 (11)	0.38889 (8)	0.0171 (3)
C20	0.49429 (18)	0.66222 (11)	0.35024 (9)	0.0235 (3)
H20A	0.4595	0.6491	0.2981	0.035*
H20B	0.5736	0.7093	0.3570	0.035*
H20C	0.4166	0.6899	0.3704	0.035*
C21	0.63093 (17)	0.48966 (12)	0.17852 (9)	0.0199 (3)
H21	0.6754	0.5535	0.1980	0.024*
C22	0.6130 (2)	0.48986 (15)	0.09632 (9)	0.0321 (4)
H22A	0.5505	0.5445	0.0754	0.048*
H22B	0.5703	0.4274	0.0764	0.048*
H22C	0.7066	0.4978	0.0840	0.048*
C23	0.72535 (18)	0.40664 (13)	0.21459 (10)	0.0274 (4)
H23A	0.7328	0.4091	0.2674	0.041*
H23B	0.8207	0.4135	0.2042	0.041*
H23C	0.6838	0.3434	0.1956	0.041*
C24	0.17566 (17)	0.41249 (11)	0.09929 (8)	0.0181 (3)
H24	0.2249	0.4755	0.0922	0.022*
C25	0.1915 (2)	0.34129 (13)	0.03960 (9)	0.0308 (4)
H25A	0.2932	0.3306	0.0411	0.046*
H25B	0.1451	0.3687	-0.0079	0.046*

supplementary materials

H25C	0.1464	0.2784	0.0470	0.046*
C26	0.02027 (18)	0.43437 (14)	0.09889 (10)	0.0279 (4)
H26A	0.0150	0.4809	0.1381	0.042*
H26B	-0.0286	0.3731	0.1065	0.042*
H26C	-0.0259	0.4632	0.0518	0.042*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ti1	0.01437 (14)	0.01316 (13)	0.01128 (13)	0.00042 (10)	0.00319 (9)	0.00035 (9)
O1	0.0175 (5)	0.0147 (5)	0.0178 (5)	0.0012 (4)	0.0042 (4)	0.0013 (4)
O2	0.0170 (5)	0.0156 (5)	0.0125 (5)	0.0015 (4)	0.0016 (4)	-0.0008 (4)
O3	0.0177 (5)	0.0219 (5)	0.0178 (5)	0.0005 (4)	0.0068 (4)	0.0032 (4)
O4	0.0201 (5)	0.0170 (5)	0.0130 (5)	-0.0005 (4)	0.0013 (4)	0.0005 (4)
N1	0.0133 (6)	0.0169 (6)	0.0119 (6)	-0.0007 (5)	0.0017 (5)	-0.0018 (4)
N2	0.0132 (6)	0.0164 (6)	0.0144 (6)	-0.0013 (5)	0.0041 (5)	0.0001 (5)
C1	0.0155 (7)	0.0176 (7)	0.0158 (7)	-0.0004 (6)	-0.0006 (5)	-0.0037 (5)
C2	0.0216 (8)	0.0172 (7)	0.0240 (8)	0.0025 (6)	-0.0016 (6)	-0.0030 (6)
C3	0.0224 (8)	0.0231 (8)	0.0296 (9)	0.0087 (7)	-0.0013 (7)	-0.0095 (7)
C4	0.0170 (8)	0.0343 (9)	0.0262 (8)	0.0060 (7)	0.0039 (6)	-0.0099 (7)
C5	0.0136 (7)	0.0289 (8)	0.0177 (7)	0.0009 (6)	0.0011 (6)	-0.0054 (6)
C6	0.0131 (7)	0.0185 (7)	0.0133 (7)	0.0000 (6)	-0.0005 (5)	-0.0034 (5)
C7	0.0162 (8)	0.0388 (10)	0.0197 (8)	-0.0011 (7)	0.0071 (6)	-0.0023 (7)
C8	0.0192 (8)	0.0305 (9)	0.0189 (8)	-0.0048 (7)	0.0049 (6)	0.0035 (6)
C9	0.0154 (7)	0.0214 (7)	0.0129 (7)	-0.0034 (6)	0.0003 (5)	0.0001 (5)
C10	0.0216 (8)	0.0205 (8)	0.0214 (8)	-0.0019 (6)	0.0056 (6)	0.0045 (6)
C11	0.0128 (7)	0.0189 (7)	0.0139 (7)	0.0002 (5)	0.0034 (5)	0.0007 (5)
C12	0.0196 (8)	0.0188 (7)	0.0205 (8)	0.0022 (6)	0.0042 (6)	0.0016 (6)
C13	0.0208 (8)	0.0237 (8)	0.0223 (8)	0.0055 (6)	0.0025 (6)	0.0077 (6)
C14	0.0173 (8)	0.0296 (8)	0.0164 (7)	0.0009 (6)	0.0010 (6)	0.0047 (6)
C15	0.0136 (7)	0.0240 (8)	0.0140 (7)	-0.0016 (6)	0.0042 (6)	0.0013 (6)
C16	0.0122 (7)	0.0182 (7)	0.0141 (7)	-0.0002 (5)	0.0049 (5)	0.0014 (5)
C17	0.0152 (7)	0.0301 (8)	0.0142 (7)	-0.0047 (6)	0.0030 (6)	-0.0032 (6)
C18	0.0183 (8)	0.0221 (8)	0.0191 (7)	-0.0062 (6)	0.0052 (6)	-0.0061 (6)
C19	0.0151 (7)	0.0188 (7)	0.0182 (7)	-0.0037 (6)	0.0057 (6)	-0.0018 (6)
C20	0.0278 (9)	0.0167 (7)	0.0238 (8)	-0.0025 (6)	0.0009 (7)	-0.0008 (6)
C21	0.0189 (8)	0.0216 (8)	0.0210 (8)	-0.0041 (6)	0.0081 (6)	-0.0003 (6)
C22	0.0325 (10)	0.0447 (11)	0.0233 (9)	0.0086 (8)	0.0156 (7)	0.0071 (8)
C23	0.0180 (8)	0.0340 (9)	0.0305 (9)	0.0017 (7)	0.0059 (7)	0.0053 (7)
C24	0.0218 (8)	0.0189 (7)	0.0130 (7)	0.0004 (6)	0.0024 (6)	0.0019 (5)
C25	0.0447 (11)	0.0302 (9)	0.0160 (8)	0.0091 (8)	0.0031 (7)	-0.0011 (7)
C26	0.0225 (9)	0.0362 (10)	0.0245 (8)	0.0057 (7)	0.0037 (7)	0.0071 (7)

Geometric parameters (\AA , $^\circ$)

Ti1—O3	1.7766 (11)	C12—C13	1.414 (2)
Ti1—O4	1.8255 (10)	C12—H12	0.9500
Ti1—O2	1.9130 (10)	C13—C14	1.368 (2)
Ti1—O1	1.9255 (10)	C13—H13	0.9500

Ti1—N2	2.3822 (12)	C14—C15	1.416 (2)
Ti1—N1	2.4130 (12)	C14—H14	0.9500
O1—C1	1.3266 (18)	C15—C16	1.413 (2)
O2—C11	1.3338 (17)	C15—C17	1.413 (2)
O3—C21	1.4153 (18)	C17—C18	1.359 (2)
O4—C24	1.4239 (17)	C17—H17	0.9500
N1—C9	1.3311 (19)	C18—C19	1.422 (2)
N1—C6	1.3756 (19)	C18—H18	0.9500
N2—C19	1.3270 (19)	C19—C20	1.496 (2)
N2—C16	1.3731 (19)	C20—H20A	0.9800
C1—C2	1.382 (2)	C20—H20B	0.9800
C1—C6	1.419 (2)	C20—H20C	0.9800
C2—C3	1.407 (2)	C21—C23	1.511 (2)
C2—H2	0.9500	C21—C22	1.515 (2)
C3—C4	1.365 (3)	C21—H21	1.0000
C3—H3	0.9500	C22—H22A	0.9800
C4—C5	1.418 (2)	C22—H22B	0.9800
C4—H4	0.9500	C22—H22C	0.9800
C5—C7	1.409 (2)	C23—H23A	0.9800
C5—C6	1.414 (2)	C23—H23B	0.9800
C7—C8	1.360 (2)	C23—H23C	0.9800
C7—H7	0.9500	C24—C25	1.512 (2)
C8—C9	1.421 (2)	C24—C26	1.517 (2)
C8—H8	0.9500	C24—H24	1.0000
C9—C10	1.497 (2)	C25—H25A	0.9800
C10—H10A	0.9800	C25—H25B	0.9800
C10—H10B	0.9800	C25—H25C	0.9800
C10—H10C	0.9800	C26—H26A	0.9800
C11—C12	1.380 (2)	C26—H26B	0.9800
C11—C16	1.419 (2)	C26—H26C	0.9800
O3—Ti1—O4	101.96 (5)	C14—C13—C12	121.83 (15)
O3—Ti1—O2	101.84 (5)	C14—C13—H13	119.1
O4—Ti1—O2	95.62 (4)	C12—C13—H13	119.1
O3—Ti1—O1	93.07 (5)	C13—C14—C15	119.54 (14)
O4—Ti1—O1	97.56 (5)	C13—C14—H14	120.2
O2—Ti1—O1	157.58 (4)	C15—C14—H14	120.2
O3—Ti1—N2	90.48 (5)	C16—C15—C17	116.11 (14)
O4—Ti1—N2	165.52 (4)	C16—C15—C14	119.09 (14)
O2—Ti1—N2	74.36 (4)	C17—C15—C14	124.74 (14)
O1—Ti1—N2	89.03 (4)	N2—C16—C15	123.90 (14)
O3—Ti1—N1	164.98 (5)	N2—C16—C11	115.65 (13)
O4—Ti1—N1	87.60 (4)	C15—C16—C11	120.37 (14)
O2—Ti1—N1	88.54 (4)	C18—C17—C15	119.67 (14)
O1—Ti1—N1	74.02 (4)	C18—C17—H17	120.2
N2—Ti1—N1	81.82 (4)	C15—C17—H17	120.2
C1—O1—Ti1	125.12 (9)	C17—C18—C19	120.95 (14)
C11—O2—Ti1	124.87 (9)	C17—C18—H18	119.5
C21—O3—Ti1	154.13 (10)	C19—C18—H18	119.5
C24—O4—Ti1	126.66 (9)	N2—C19—C18	120.93 (14)

supplementary materials

C9—N1—C6	117.94 (13)	N2—C19—C20	120.36 (14)
C9—N1—Ti1	134.90 (10)	C18—C19—C20	118.70 (13)
C6—N1—Ti1	107.09 (9)	C19—C20—H20A	109.5
C19—N2—C16	118.24 (13)	C19—C20—H20B	109.5
C19—N2—Ti1	134.14 (10)	H20A—C20—H20B	109.5
C16—N2—Ti1	107.62 (9)	C19—C20—H20C	109.5
O1—C1—C2	123.29 (14)	H20A—C20—H20C	109.5
O1—C1—C6	117.65 (13)	H20B—C20—H20C	109.5
C2—C1—C6	119.06 (14)	O3—C21—C23	110.24 (12)
C1—C2—C3	120.43 (15)	O3—C21—C22	108.63 (13)
C1—C2—H2	119.8	C23—C21—C22	112.47 (14)
C3—C2—H2	119.8	O3—C21—H21	108.5
C4—C3—C2	121.38 (15)	C23—C21—H21	108.5
C4—C3—H3	119.3	C22—C21—H21	108.5
C2—C3—H3	119.3	C21—C22—H22A	109.5
C3—C4—C5	119.79 (15)	C21—C22—H22B	109.5
C3—C4—H4	120.1	H22A—C22—H22B	109.5
C5—C4—H4	120.1	C21—C22—H22C	109.5
C7—C5—C6	116.69 (14)	H22A—C22—H22C	109.5
C7—C5—C4	124.21 (15)	H22B—C22—H22C	109.5
C6—C5—C4	119.09 (15)	C21—C23—H23A	109.5
N1—C6—C5	123.64 (14)	C21—C23—H23B	109.5
N1—C6—C1	116.11 (13)	H23A—C23—H23B	109.5
C5—C6—C1	120.24 (14)	C21—C23—H23C	109.5
C8—C7—C5	119.49 (15)	H23A—C23—H23C	109.5
C8—C7—H7	120.3	H23B—C23—H23C	109.5
C5—C7—H7	120.3	O4—C24—C25	108.89 (12)
C7—C8—C9	120.86 (15)	O4—C24—C26	109.19 (12)
C7—C8—H8	119.6	C25—C24—C26	112.20 (14)
C9—C8—H8	119.6	O4—C24—H24	108.8
N1—C9—C8	121.34 (14)	C25—C24—H24	108.8
N1—C9—C10	120.18 (13)	C26—C24—H24	108.8
C8—C9—C10	118.48 (13)	C24—C25—H25A	109.5
C9—C10—H10A	109.5	C24—C25—H25B	109.5
C9—C10—H10B	109.5	H25A—C25—H25B	109.5
H10A—C10—H10B	109.5	C24—C25—H25C	109.5
C9—C10—H10C	109.5	H25A—C25—H25C	109.5
H10A—C10—H10C	109.5	H25B—C25—H25C	109.5
H10B—C10—H10C	109.5	C24—C26—H26A	109.5
O2—C11—C12	123.48 (13)	C24—C26—H26B	109.5
O2—C11—C16	117.07 (13)	H26A—C26—H26B	109.5
C12—C11—C16	119.45 (13)	C24—C26—H26C	109.5
C11—C12—C13	119.68 (15)	H26A—C26—H26C	109.5
C11—C12—H12	120.2	H26B—C26—H26C	109.5
C13—C12—H12	120.2		
O3—Ti1—O1—C1	-171.18 (11)	Ti1—N1—C6—C1	-0.15 (14)
O4—Ti1—O1—C1	86.32 (11)	C7—C5—C6—N1	-1.2 (2)
O2—Ti1—O1—C1	-39.19 (19)	C4—C5—C6—N1	179.55 (14)
N2—Ti1—O1—C1	-80.75 (11)	C7—C5—C6—C1	178.57 (14)

N1—Ti1—O1—C1	1.02 (11)	C4—C5—C6—C1	-0.6 (2)
O3—Ti1—O2—C11	91.30 (11)	O1—C1—C6—N1	0.93 (19)
O4—Ti1—O2—C11	-165.20 (11)	C2—C1—C6—N1	-179.29 (13)
O1—Ti1—O2—C11	-39.37 (18)	O1—C1—C6—C5	-178.89 (13)
N2—Ti1—O2—C11	4.16 (10)	C2—C1—C6—C5	0.9 (2)
N1—Ti1—O2—C11	-77.75 (11)	C6—C5—C7—C8	-0.6 (2)
O4—Ti1—O3—C21	-112.2 (2)	C4—C5—C7—C8	178.58 (15)
O2—Ti1—O3—C21	-13.8 (2)	C5—C7—C8—C9	1.4 (2)
O1—Ti1—O3—C21	149.4 (2)	C6—N1—C9—C8	-1.4 (2)
N2—Ti1—O3—C21	60.3 (2)	Ti1—N1—C9—C8	-177.92 (10)
N1—Ti1—O3—C21	119.1 (2)	C6—N1—C9—C10	178.99 (13)
O3—Ti1—O4—C24	-50.83 (12)	Ti1—N1—C9—C10	2.4 (2)
O2—Ti1—O4—C24	-154.22 (11)	C7—C8—C9—N1	-0.4 (2)
O1—Ti1—O4—C24	43.96 (12)	C7—C8—C9—C10	179.26 (15)
N2—Ti1—O4—C24	160.41 (16)	Ti1—O2—C11—C12	178.57 (11)
N1—Ti1—O4—C24	117.49 (11)	Ti1—O2—C11—C16	-1.88 (18)
O3—Ti1—N1—C9	-152.04 (17)	O2—C11—C12—C13	-178.34 (14)
O4—Ti1—N1—C9	77.85 (14)	C16—C11—C12—C13	2.1 (2)
O2—Ti1—N1—C9	-17.84 (14)	C11—C12—C13—C14	-0.8 (2)
O1—Ti1—N1—C9	176.41 (14)	C12—C13—C14—C15	-1.1 (2)
N2—Ti1—N1—C9	-92.25 (14)	C13—C14—C15—C16	1.7 (2)
O3—Ti1—N1—C6	31.1 (2)	C13—C14—C15—C17	-175.58 (15)
O4—Ti1—N1—C6	-98.96 (9)	C19—N2—C16—C15	3.5 (2)
O2—Ti1—N1—C6	165.34 (9)	Ti1—N2—C16—C15	-176.65 (11)
O1—Ti1—N1—C6	-0.40 (9)	C19—N2—C16—C11	-173.27 (13)
N2—Ti1—N1—C6	90.94 (9)	Ti1—N2—C16—C11	6.53 (14)
O3—Ti1—N2—C19	71.95 (14)	C17—C15—C16—N2	0.4 (2)
O4—Ti1—N2—C19	-138.53 (18)	C14—C15—C16—N2	-177.12 (13)
O2—Ti1—N2—C19	174.12 (14)	C17—C15—C16—C11	177.10 (13)
O1—Ti1—N2—C19	-21.11 (14)	C14—C15—C16—C11	-0.4 (2)
N1—Ti1—N2—C19	-95.11 (14)	O2—C11—C16—N2	-4.13 (19)
O3—Ti1—N2—C16	-107.81 (9)	C12—C11—C16—N2	175.45 (13)
O4—Ti1—N2—C16	41.7 (2)	O2—C11—C16—C15	178.93 (12)
O2—Ti1—N2—C16	-5.64 (9)	C12—C11—C16—C15	-1.5 (2)
O1—Ti1—N2—C16	159.13 (9)	C16—C15—C17—C18	-3.3 (2)
N1—Ti1—N2—C16	85.13 (9)	C14—C15—C17—C18	174.09 (15)
Ti1—O1—C1—C2	178.74 (11)	C15—C17—C18—C19	2.3 (2)
Ti1—O1—C1—C6	-1.49 (18)	C16—N2—C19—C18	-4.6 (2)
O1—C1—C2—C3	179.34 (14)	Ti1—N2—C19—C18	175.66 (10)
C6—C1—C2—C3	-0.4 (2)	C16—N2—C19—C20	174.44 (13)
C1—C2—C3—C4	-0.3 (2)	Ti1—N2—C19—C20	-5.3 (2)
C2—C3—C4—C5	0.6 (2)	C17—C18—C19—N2	1.8 (2)
C3—C4—C5—C7	-179.23 (16)	C17—C18—C19—C20	-177.27 (14)
C3—C4—C5—C6	-0.1 (2)	Ti1—O3—C21—C23	8.2 (3)
C9—N1—C6—C5	2.2 (2)	Ti1—O3—C21—C22	131.9 (2)
Ti1—N1—C6—C5	179.66 (12)	Ti1—O4—C24—C25	132.10 (12)
C9—N1—C6—C1	-177.60 (13)	Ti1—O4—C24—C26	-105.08 (14)

Fig. 1

