

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# 5-*n*-Butyl-4-[2-(2-ethyl-1-benzothiophen-3-yl)-3,3,4,4,5,5-hexafluorocyclopent-1-en-1-yl]thiophene-2-carbaldehyde

 Zhen Yang,<sup>a</sup> Congbin Fan,<sup>a\*</sup> Min Li,<sup>a</sup> Weijun Liu,<sup>a</sup> Gang Liu<sup>a</sup> and Seik Weng Ng<sup>b</sup>

<sup>a</sup>Jiangxi Key Laboratory of Organic Chemistry, Jiangxi Science & Technology Normal University, Nanchang 330013, People's Republic of China, and <sup>b</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: congbinfan@yahoo.com.cn


Received 23 August 2008; accepted 6 November 2008

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å; disorder in main residue;  $R$  factor = 0.046;  $wR$  factor = 0.151; data-to-parameter ratio = 16.3.

The title compound,  $\text{C}_{24}\text{H}_{20}\text{F}_6\text{OS}_2$ , exhibiting photochromic behaviour, has thienyl and benzothieryl substituents attached to the double-bond C atoms of the envelope-shaped cyclopentene ring. The mean planes of aromatic systems form dihedral angles of 43.0 (1) (thienyl) and 73.8 (1)° (benzothieryl) with the mean plane of the C=C—C portion of the cyclopentene ring. This conformation avoids steric hindrance between the *n*-butyl and ethyl substituents. The formyl substituent of the thienyl group, as well as the ethyl substituent of the benzothieryl group, are disordered [occupancies of 0.788 (17):0.212 (17) and 0.64 (5):0.36 (5), respectively].

## Related literature

For the synthesis of the precursors, see: Pu *et al.* (2008); Ramamurthy & Venkatesan (1987); Kobatake & Irie (2004); Zheng *et al.* (2007). For the crystal structures of other photochromic dithienyl-substituted hexafluorocyclopentenes, see: Congbin *et al.* (2007); Li *et al.* (2008); Liu *et al.* (2008); Pu & Zhou (2007); Tu *et al.* (2008); Zhu *et al.* (2007).



## Experimental

## Crystal data

$\text{C}_{24}\text{H}_{20}\text{F}_6\text{OS}_2$	$\gamma = 103.542$ (1)°
$M_r = 502.52$	$V = 1157.5$ (2) Å <sup>3</sup>
Triclinic, $P\bar{1}$	$Z = 2$
$a = 10.051$ (1) Å	Mo $K\alpha$ radiation
$b = 11.031$ (1) Å	$\mu = 0.29$ mm <sup>-1</sup>
$c = 12.019$ (1) Å	$T = 296$ (2) K
$\alpha = 113.126$ (1)°	$0.43 \times 0.43 \times 0.43$ mm
$\beta = 96.882$ (1)°	

## Data collection

Bruker SMART area-detector diffractometer	5195 independent reflections
Absorption correction: none	3669 reflections with $I > 2\sigma(I)$
10086 measured reflections	$R_{\text{int}} = 0.019$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$	30 restraints
$wR(F^2) = 0.151$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\text{max}} = 0.34$ e Å <sup>-3</sup>
5195 reflections	$\Delta\rho_{\text{min}} = -0.25$ e Å <sup>-3</sup>
318 parameters	

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINTE* (Bruker, 1997); data reduction: *SAINTE* program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *pubCIF* (Westrip, 2008).

This work was supported by the Natural Science Foundation of Jiangxi (0620012) and the Science Fund of the Education Office of Jiangxi ([2007]279).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: YA2080).

## References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
 Bruker (1997). *SMART* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Congbin, F., Tianshe, Y., Qidong, T. & Gang, L. (2007). *Acta Cryst.* **E63**, o4721.  
 Kobatake, S. & Irie, M. (2004). *Bull. Chem. Soc. Jpn.* **77**, 195–210.  
 Li, M., Pu, S.-Z., Fan, C.-B. & Le, Z.-G. (2008). *Acta Cryst.* **E64**, o517.  
 Liu, G., Tu, Q., Zhang, Q., Fan, C. & Yang, T. (2008). *Acta Cryst.* **E64**, o938.  
 Pu, S.-Z., Li, M., Fan, C.-B., Liu, G. & Shen, L. (2008). *J. Mol. Struct.* doi:10.1016/j.molstruc.2008.08.023.  
 Pu, S.-Z. & Zhou, Q.-F. (2007). *Acta Cryst.* **E63**, o927–o928.  
 Ramamurthy, V. & Venkatesan, K. (1987). *Chem. Rev.* **87**, 433–481.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Tu, Q., Fan, C., Liu, G., Li, M. & Ng, S. W. (2008). *Acta Cryst.* **E64**, o1007.  
 Westrip, S. P. (2008). *pubCIF*. In preparation.  
 Zheng, C.-H., Pu, S.-Z., Xu, J.-K., Luo, M.-B., Huang, D.-C. & Shen, L. (2007). *Tetrahedron*, **63**, 5437–5449.  
 Zhu, S., Rao, Y., Liu, G., Xie, Y. & Pu, S. (2007). *Acta Cryst.* **E63**, o4175.

**supplementary materials**

*Acta Cryst.* (2008). E64, o2311 [ doi:10.1107/S1600536808036453 ]

## 5-*n*-Butyl-4-[2-(2-ethyl-1-benzothiophen-3-yl)-3,3,4,4,5,5-hexafluorocyclopent-1-en-1-yl]thiophene-2-carbaldehyde

Z. Yang, C. Fan, M. Li, W. Liu, G. Liu and S. W. Ng

### Comment

The title compound has thienyl and benzothienyl substituents on the double-bond C atoms of the envelope-shaped cyclopentene ring. The planes of aromatic systems form dihedral angles 43.0 (1)° (thienyl) and 73.8 (1)° (benzothienyl) with the mean plane of the C4–C11=C15–C16 portion of the cyclopentene ring. Such conformation allows to avoid steric hindrance between the *n*-butyl and ethyl substituents (Fig. 1). The terminal atoms of formyl substituent on the thienyl group as well as the ethyl substituent on the benzothienyl group are disordered over two positions. The intramolecular distance between C3 and C20 is 3.989 (8) Å. This distance indicates that the compound may undergo photochromism in crystalline phase to form the closed ring isomer, as photochromic reactivity was shown to occur when the distance between the potentially reactive C atoms is less than 4.2 Å (Ramamurthy & Venkatesan, 1987; Kobatake & Irie, 2004).

Indeed, crystals of the title compound show photochromism: upon irradiation with 365 nm light, the colourless crystals rapidly turn blue, and the blue crystals turn colourless again upon irradiation with visible light (>510 nm). When dissolved in hexane, the blue compound displays an absorption maximum at 581 nm; the solution of colourless compound has absorption maximum at 257 nm.

### Experimental

2.0 mL (5 mmol) of *n*-butyllithium was added under stirring in nitrogen atmosphere to 30 mL of THF solution containing 1.46 g (5.0 mmol) of 4-bromo-5-*n*-butyl-2-(1,3-dioxolane)-thiophene (Zheng *et al.*, 2007) at 195 K. 40 min later, 10 mL of THF solution containing 1.77 g (5.0 mmol) of 1-(2-ethyl-1-benzothien-3-yl)heptafluorocyclopentene (Pu *et al.*, 2008) was added to the reaction mixture and stirring under nitrogen atmosphere at 195 K was continued for two more hours. The reaction mixture was extracted with diethyl ether and evaporated in vacuum. Then the obtained compound was hydrolyzed by *p*-toluenesulfonic acid (0.4 g) in mixture of water (30 ml) and acetone (90 ml). Pyridine (2 ml) was added, and the solution was refluxed for 24 hours and then washed with aqueous sodium bicarbonate. The mixed compound was extracted with diethyl ether and evaporated in vacuum. The crude product was purified by column chromatography on silica, with ethyl acetate and petroleum ether (v/v 1/6) as the eluent, to give 1.58 g (3.15 mmol, 63% yield) of the title compound. Elemental analysis: calc. for C<sub>24</sub>H<sub>20</sub>F<sub>6</sub>OS<sub>2</sub>: C 57.36, H, 4.01%. Found C 57.22, H 3.90%.

### Refinement

All H atoms were positioned geometrically and treated as riding with C–H = 0.97 Å (methylene), 0.96 Å (methyl) or 0.93 Å (aromatic and formyl) with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  ( $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$  for methyl H atoms).

The methyl group of the ethyl chain is disordered over two sites C1 and C1'; the the C3–C2 distance was restrained to 1.50±0.01 Å, C2–C1 and C2–C1' distances were restrained to be equal within 0.01 Å, and C3...C1 and C3...C1' restrained to 2.51±0.01 Å. The occupancies of the disorder components refined to a 0.79 (1):0.21 ratio.

## supplementary materials

The formyl oxygen atom is also disordered over two positions; the C19–O1 and C19–O1' distances were restrained to be equal within 0.01 Å. The occupancies of the O1 and O1' atoms refined to a 0.64 (1):0.36 ratio.

The anisotropic displacement parameters of the disordered atoms were restrained to be nearly isotropic.

### Figures




Fig. 1. Molecular structure of the title compound and the atom-labelling scheme; thermal displacement ellipsoids are drawn at the 30% probability level. Minor components of the disorder are shown with the dashed bonds; H atoms are omitted.

### 5-*n*-Butyl-4-[2-(2-ethyl-1-benzothiophen-3-yl)-3,3,4,4,5,5-hexafluorocyclopent-1-en-1-yl]thiophene-2-carbaldehyde

#### Crystal data

C<sub>24</sub>H<sub>20</sub>F<sub>6</sub>OS<sub>2</sub>

*M<sub>r</sub>* = 502.52

Triclinic, *PT*

Hall symbol: -P 1

*a* = 10.051 (1) Å

*b* = 11.031 (1) Å

*c* = 12.019 (1) Å

α = 113.126 (1)°

β = 96.882 (1)°

γ = 103.542 (1)°

*V* = 1157.5 (2) Å<sup>3</sup>

*Z* = 2

*F*<sub>000</sub> = 516

*D<sub>x</sub>* = 1.442 Mg m<sup>-3</sup>

Mo *K*α radiation

λ = 0.71073 Å

Cell parameters from 4259 reflections

θ = 2.5–28.2°

μ = 0.29 mm<sup>-1</sup>

*T* = 296 (2) K

Block, colourless

0.43 × 0.43 × 0.43 mm

#### Data collection

Bruker SMART area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

*T* = 296(2) K

φ and ω scans

Absorption correction: None

10086 measured reflections

5195 independent reflections

3669 reflections with *I* > 2σ(*I*)

*R*<sub>int</sub> = 0.019

θ<sub>max</sub> = 27.5°

θ<sub>min</sub> = 2.5°

*h* = -12→13

*k* = -14→14

*l* = -15→15

Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.046$	H-atom parameters constrained
$wR(F^2) = 0.151$	$w = 1/[\sigma^2(F_o^2) + (0.0768P)^2 + 0.3784P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
5195 reflections	$(\Delta/\sigma)_{\max} = 0.001$
318 parameters	$\Delta\rho_{\max} = 0.34 \text{ e } \text{\AA}^{-3}$
30 restraints	$\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
S1	0.43383 (7)	0.52869 (8)	0.19023 (8)	0.0736 (2)	
S2	0.86262 (7)	0.60297 (6)	0.65571 (5)	0.05353 (19)	
F1	0.89175 (19)	0.80884 (18)	0.17554 (17)	0.0805 (5)	
F2	0.8985 (2)	0.6035 (2)	0.06303 (14)	0.0868 (6)	
F3	1.16124 (19)	0.8540 (2)	0.22574 (18)	0.0943 (6)	
F4	1.1390 (2)	0.6431 (2)	0.19936 (18)	0.0921 (6)	
F5	1.10459 (17)	0.92568 (14)	0.44135 (15)	0.0707 (4)	
F6	1.19297 (14)	0.76341 (19)	0.43741 (16)	0.0735 (5)	
O1'	0.947 (3)	0.8292 (9)	0.9172 (7)	0.091 (4)	0.64 (5)
O1	1.010 (3)	0.8304 (18)	0.9249 (10)	0.080 (4)	0.36 (5)
C1	0.5164 (7)	0.8436 (7)	0.3633 (7)	0.104 (3)	0.788 (17)
H1A	0.5476	0.9370	0.4276	0.156*	0.788 (17)
H1B	0.4403	0.7889	0.3812	0.156*	0.788 (17)
H1C	0.4848	0.8435	0.2846	0.156*	0.788 (17)
C1'	0.577 (3)	0.8783 (13)	0.3189 (17)	0.077 (6)	0.212 (17)
H1'A	0.6029	0.9682	0.3880	0.116*	0.212 (17)
H1'B	0.4767	0.8414	0.2910	0.116*	0.212 (17)
H1'C	0.6169	0.8861	0.2522	0.116*	0.212 (17)
C2	0.6323 (4)	0.7854 (3)	0.3580 (3)	0.0885 (10)	
H2A	0.6646	0.7900	0.4393	0.106*	
H2B	0.7092	0.8446	0.3433	0.106*	
C3	0.6050 (3)	0.6395 (3)	0.2628 (2)	0.0565 (6)	
C4	0.7027 (2)	0.5769 (2)	0.2221 (2)	0.0454 (5)	
C5	0.6422 (3)	0.4353 (2)	0.1294 (2)	0.0519 (6)	
C6	0.7084 (3)	0.3380 (3)	0.0696 (2)	0.0677 (7)	
H6	0.8061	0.3627	0.0841	0.081*	
C7	0.6267 (5)	0.2048 (3)	-0.0111 (3)	0.0914 (11)	
H7	0.6703	0.1400	-0.0517	0.110*	
C8	0.4127 (4)	0.2581 (4)	0.0223 (3)	0.0846 (10)	

## supplementary materials

H8	0.3149	0.2315	0.0058	0.102*	
C9	0.4813 (5)	0.1652 (4)	-0.0332 (3)	0.0991 (13)	
H9	0.4292	0.0739	-0.0867	0.119*	
C10	0.4937 (3)	0.3945 (3)	0.1048 (2)	0.0615 (7)	
C11	0.8549 (2)	0.6521 (2)	0.2655 (2)	0.0424 (5)	
C12	0.9296 (3)	0.6987 (3)	0.1805 (2)	0.0560 (6)	
C13	1.0846 (3)	0.7499 (3)	0.2438 (2)	0.0576 (6)	
C14	1.0847 (2)	0.7868 (2)	0.3798 (2)	0.0480 (5)	
C15	0.9421 (2)	0.7024 (2)	0.37843 (19)	0.0387 (4)	
C16	0.9193 (2)	0.6924 (2)	0.49357 (19)	0.0385 (4)	
C17	0.9764 (2)	0.8081 (2)	0.6132 (2)	0.0467 (5)	
H17	1.0247	0.8966	0.6245	0.056*	
C18	0.9532 (3)	0.7762 (2)	0.7089 (2)	0.0533 (6)	
C19	0.9934 (4)	0.8673 (3)	0.8416 (3)	0.0801 (9)	
H19	1.0552	0.9562	0.8702	0.096*	0.64 (5)
H19'	1.0075	0.9613	0.8654	0.096*	0.36 (5)
C20	0.8536 (2)	0.5729 (2)	0.50312 (19)	0.0401 (4)	
C21	0.7831 (2)	0.4277 (2)	0.4049 (2)	0.0425 (5)	
H21A	0.8296	0.3668	0.4216	0.051*	
H21B	0.7949	0.4238	0.3247	0.051*	
C22	0.6266 (2)	0.3746 (2)	0.3974 (2)	0.0510 (5)	
H22A	0.6139	0.3879	0.4796	0.061*	
H22B	0.5782	0.4289	0.3713	0.061*	
C23	0.5601 (3)	0.2226 (3)	0.3076 (3)	0.0634 (7)	
H23A	0.6027	0.1675	0.3378	0.076*	
H23B	0.5800	0.2077	0.2271	0.076*	
C24	0.4014 (4)	0.1731 (4)	0.2918 (3)	0.0940 (11)	
H24A	0.3648	0.0766	0.2346	0.141*	
H24B	0.3583	0.2255	0.2598	0.141*	
H24C	0.3810	0.1860	0.3709	0.141*	

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0440 (4)	0.0835 (5)	0.0917 (6)	0.0164 (3)	-0.0006 (3)	0.0430 (4)
S2	0.0674 (4)	0.0499 (3)	0.0462 (3)	0.0127 (3)	0.0157 (3)	0.0262 (3)
F1	0.0839 (11)	0.0906 (12)	0.0950 (12)	0.0251 (9)	0.0131 (9)	0.0717 (11)
F2	0.0934 (13)	0.0972 (13)	0.0451 (9)	-0.0039 (10)	0.0150 (8)	0.0255 (9)
F3	0.0792 (12)	0.1137 (15)	0.0913 (13)	-0.0117 (10)	0.0169 (10)	0.0701 (12)
F4	0.0853 (12)	0.1121 (15)	0.0909 (13)	0.0522 (11)	0.0407 (10)	0.0382 (12)
F5	0.0776 (10)	0.0444 (8)	0.0744 (10)	-0.0017 (7)	0.0125 (8)	0.0233 (7)
F6	0.0408 (7)	0.1085 (13)	0.0919 (11)	0.0184 (8)	0.0082 (7)	0.0688 (11)
O1'	0.127 (8)	0.085 (3)	0.047 (2)	0.009 (4)	0.025 (3)	0.0263 (19)
O1	0.107 (9)	0.084 (5)	0.053 (4)	0.032 (5)	0.015 (4)	0.033 (3)
C1	0.098 (4)	0.102 (4)	0.114 (4)	0.061 (3)	0.027 (3)	0.031 (3)
C1'	0.082 (10)	0.072 (8)	0.084 (9)	0.033 (7)	0.013 (6)	0.036 (6)
C2	0.080 (2)	0.0667 (19)	0.102 (2)	0.0363 (16)	0.0055 (18)	0.0168 (18)
C3	0.0474 (13)	0.0579 (14)	0.0639 (15)	0.0171 (11)	0.0035 (11)	0.0285 (12)

C4	0.0462 (12)	0.0475 (12)	0.0416 (11)	0.0112 (9)	0.0022 (9)	0.0226 (10)
C5	0.0569 (14)	0.0512 (13)	0.0419 (12)	0.0071 (10)	0.0020 (10)	0.0228 (10)
C6	0.0835 (19)	0.0575 (15)	0.0530 (15)	0.0147 (14)	0.0191 (14)	0.0182 (12)
C7	0.128 (3)	0.0590 (18)	0.0636 (19)	0.0153 (19)	0.028 (2)	0.0098 (15)
C8	0.077 (2)	0.075 (2)	0.0695 (19)	-0.0144 (17)	-0.0182 (16)	0.0302 (17)
C9	0.126 (3)	0.062 (2)	0.063 (2)	-0.012 (2)	-0.001 (2)	0.0117 (16)
C10	0.0552 (14)	0.0627 (15)	0.0549 (14)	0.0001 (12)	-0.0070 (11)	0.0298 (12)
C11	0.0446 (11)	0.0412 (11)	0.0440 (11)	0.0122 (9)	0.0084 (9)	0.0225 (9)
C12	0.0625 (15)	0.0605 (14)	0.0468 (13)	0.0110 (12)	0.0099 (11)	0.0308 (12)
C13	0.0538 (14)	0.0631 (15)	0.0651 (16)	0.0132 (12)	0.0200 (12)	0.0381 (13)
C14	0.0416 (11)	0.0476 (12)	0.0573 (14)	0.0091 (9)	0.0057 (10)	0.0296 (11)
C15	0.0392 (10)	0.0366 (10)	0.0445 (11)	0.0130 (8)	0.0096 (8)	0.0213 (9)
C16	0.0366 (10)	0.0395 (10)	0.0430 (11)	0.0134 (8)	0.0057 (8)	0.0215 (9)
C17	0.0537 (12)	0.0384 (11)	0.0466 (12)	0.0122 (9)	0.0082 (10)	0.0192 (9)
C18	0.0645 (15)	0.0470 (12)	0.0453 (12)	0.0153 (11)	0.0112 (11)	0.0186 (10)
C19	0.126 (3)	0.0564 (16)	0.0457 (15)	0.0148 (17)	0.0188 (16)	0.0175 (13)
C20	0.0404 (10)	0.0419 (11)	0.0431 (11)	0.0151 (8)	0.0092 (9)	0.0224 (9)
C21	0.0455 (11)	0.0382 (10)	0.0476 (12)	0.0153 (9)	0.0087 (9)	0.0218 (9)
C22	0.0471 (12)	0.0458 (12)	0.0566 (14)	0.0088 (10)	0.0090 (10)	0.0229 (11)
C23	0.0630 (16)	0.0483 (13)	0.0684 (17)	0.0043 (12)	0.0049 (13)	0.0253 (12)
C24	0.073 (2)	0.079 (2)	0.091 (2)	-0.0168 (17)	0.0107 (17)	0.0238 (18)

*Geometric parameters (Å, °)*

S1—C10	1.726 (3)	C7—H7	0.9300
S1—C3	1.742 (3)	C8—C9	1.369 (5)
S2—C20	1.717 (2)	C8—C10	1.403 (4)
S2—C18	1.725 (2)	C8—H8	0.9300
F1—C12	1.376 (3)	C9—H9	0.9300
F2—C12	1.331 (3)	C11—C15	1.346 (3)
F3—C13	1.335 (3)	C11—C12	1.512 (3)
F4—C13	1.360 (3)	C12—C13	1.513 (4)
F5—C14	1.366 (3)	C13—C14	1.521 (3)
F6—C14	1.350 (3)	C14—C15	1.510 (3)
O1'—C19	1.236 (5)	C15—C16	1.471 (3)
O1—C19	1.228 (8)	C16—C20	1.385 (3)
C1—C2	1.453 (5)	C16—C17	1.430 (3)
C1—H1A	0.9600	C17—C18	1.358 (3)
C1—H1B	0.9600	C17—H17	0.9300
C1—H1C	0.9600	C18—C19	1.453 (4)
C1'—C2	1.468 (8)	C19—H19	0.9300
C1'—H1'A	0.9600	C19—H19'	0.9300
C1'—H1'B	0.9600	C20—C21	1.497 (3)
C1'—H1'C	0.9600	C21—C22	1.523 (3)
C2—C3	1.499 (4)	C21—H21A	0.9700
C2—H2A	0.9700	C21—H21B	0.9700
C2—H2B	0.9700	C22—C23	1.518 (3)
C3—C4	1.359 (3)	C22—H22A	0.9700
C4—C5	1.441 (3)	C22—H22B	0.9700

## supplementary materials

---

C4—C11	1.475 (3)	C23—C24	1.522 (4)
C5—C6	1.399 (4)	C23—H23A	0.9700
C5—C10	1.412 (4)	C23—H23B	0.9700
C6—C7	1.378 (4)	C24—H24A	0.9600
C6—H6	0.9300	C24—H24B	0.9600
C7—C9	1.384 (6)	C24—H24C	0.9600
C10—S1—C3	91.85 (12)	F4—C13—C12	108.6 (2)
C20—S2—C18	92.17 (11)	F3—C13—C14	113.6 (2)
C2—C1—H1A	109.5	F4—C13—C14	108.9 (2)
C2—C1—H1B	109.5	C12—C13—C14	103.50 (19)
H1A—C1—H1B	109.5	F6—C14—F5	105.70 (19)
C2—C1—H1C	109.5	F6—C14—C15	113.35 (17)
H1A—C1—H1C	109.5	F5—C14—C15	111.64 (19)
H1B—C1—H1C	109.5	F6—C14—C13	111.9 (2)
C2—C1'—H1'A	109.5	F5—C14—C13	108.90 (18)
C2—C1'—H1'B	109.5	C15—C14—C13	105.39 (18)
H1'A—C1'—H1'B	109.5	C11—C15—C16	131.72 (19)
C2—C1'—H1'C	109.5	C11—C15—C14	109.62 (18)
H1'A—C1'—H1'C	109.5	C16—C15—C14	118.62 (18)
H1'B—C1'—H1'C	109.5	C20—C16—C17	111.60 (18)
C1—C2—C3	117.8 (3)	C20—C16—C15	126.15 (19)
C1'—C2—C3	117.9 (6)	C17—C16—C15	122.02 (18)
C1—C2—H2A	107.9	C18—C17—C16	113.4 (2)
C1'—C2—H2A	132.3	C18—C17—H17	123.3
C3—C2—H2A	107.9	C16—C17—H17	123.3
C1—C2—H2B	107.9	C17—C18—C19	128.5 (2)
C3—C2—H2B	107.9	C17—C18—S2	111.32 (18)
H2A—C2—H2B	107.2	C19—C18—S2	120.2 (2)
C4—C3—C2	127.0 (2)	O1—C19—C18	125.4 (8)
C4—C3—S1	111.98 (18)	O1'—C19—C18	121.5 (5)
C2—C3—S1	121.0 (2)	O1—C19—H19	107.4
C3—C4—C5	113.4 (2)	O1'—C19—H19	119.3
C3—C4—C11	121.9 (2)	C18—C19—H19	119.3
C5—C4—C11	124.6 (2)	O1—C19—H19'	117.3
C6—C5—C10	119.1 (2)	O1'—C19—H19'	112.4
C6—C5—C4	129.8 (2)	C18—C19—H19'	117.3
C10—C5—C4	111.1 (2)	C16—C20—C21	130.78 (19)
C7—C6—C5	119.0 (3)	C16—C20—S2	111.48 (16)
C7—C6—H6	120.5	C21—C20—S2	117.72 (15)
C5—C6—H6	120.5	C20—C21—C22	113.64 (18)
C6—C7—C9	121.5 (3)	C20—C21—H21A	108.8
C6—C7—H7	119.2	C22—C21—H21A	108.8
C9—C7—H7	119.2	C20—C21—H21B	108.8
C9—C8—C10	118.4 (3)	C22—C21—H21B	108.8
C9—C8—H8	120.8	H21A—C21—H21B	107.7
C10—C8—H8	120.8	C23—C22—C21	113.0 (2)
C8—C9—C7	121.1 (3)	C23—C22—H22A	109.0
C8—C9—H9	119.4	C21—C22—H22A	109.0
C7—C9—H9	119.4	C23—C22—H22B	109.0

C8—C10—C5	120.9 (3)	C21—C22—H22B	109.0
C8—C10—S1	127.5 (3)	H22A—C22—H22B	107.8
C5—C10—S1	111.61 (19)	C22—C23—C24	112.8 (2)
C15—C11—C4	130.17 (19)	C22—C23—H23A	109.0
C15—C11—C12	110.50 (19)	C24—C23—H23A	109.0
C4—C11—C12	118.92 (19)	C22—C23—H23B	109.0
F2—C12—F1	105.7 (2)	C24—C23—H23B	109.0
F2—C12—C11	114.8 (2)	H23A—C23—H23B	107.8
F1—C12—C11	109.4 (2)	C23—C24—H24A	109.5
F2—C12—C13	113.8 (2)	C23—C24—H24B	109.5
F1—C12—C13	108.2 (2)	H24A—C24—H24B	109.5
C11—C12—C13	104.97 (18)	C23—C24—H24C	109.5
F3—C13—F4	107.5 (2)	H24A—C24—H24C	109.5
F3—C13—C12	114.5 (2)	H24B—C24—H24C	109.5
C1—C2—C3—C4	163.3 (6)	C11—C12—C13—C14	-23.4 (2)
C1'—C2—C3—C4	119.3 (14)	F3—C13—C14—F6	-88.3 (3)
C1—C2—C3—S1	-16.7 (7)	F4—C13—C14—F6	31.4 (3)
C1'—C2—C3—S1	-60.7 (14)	C12—C13—C14—F6	146.9 (2)
C10—S1—C3—C4	0.8 (2)	F3—C13—C14—F5	28.1 (3)
C10—S1—C3—C2	-179.3 (3)	F4—C13—C14—F5	147.89 (19)
C2—C3—C4—C5	-179.9 (3)	C12—C13—C14—F5	-96.7 (2)
S1—C3—C4—C5	0.1 (3)	F3—C13—C14—C15	148.0 (2)
C2—C3—C4—C11	-3.0 (4)	F4—C13—C14—C15	-92.2 (2)
S1—C3—C4—C11	176.99 (17)	C12—C13—C14—C15	23.2 (2)
C3—C4—C5—C6	-177.8 (3)	C4—C11—C15—C16	4.6 (4)
C11—C4—C5—C6	5.4 (4)	C12—C11—C15—C16	177.0 (2)
C3—C4—C5—C10	-1.1 (3)	C4—C11—C15—C14	-172.9 (2)
C11—C4—C5—C10	-177.9 (2)	C12—C11—C15—C14	-0.5 (2)
C10—C5—C6—C7	-0.4 (4)	F6—C14—C15—C11	-137.4 (2)
C4—C5—C6—C7	176.1 (3)	F5—C14—C15—C11	103.4 (2)
C5—C6—C7—C9	-0.6 (5)	C13—C14—C15—C11	-14.7 (2)
C10—C8—C9—C7	-1.4 (5)	F6—C14—C15—C16	44.7 (3)
C6—C7—C9—C8	1.6 (6)	F5—C14—C15—C16	-74.5 (2)
C9—C8—C10—C5	0.3 (4)	C13—C14—C15—C16	167.42 (19)
C9—C8—C10—S1	-177.5 (3)	C11—C15—C16—C20	47.7 (3)
C6—C5—C10—C8	0.5 (4)	C14—C15—C16—C20	-135.0 (2)
C4—C5—C10—C8	-176.5 (2)	C11—C15—C16—C17	-138.3 (2)
C6—C5—C10—S1	178.73 (19)	C14—C15—C16—C17	39.1 (3)
C4—C5—C10—S1	1.7 (3)	C20—C16—C17—C18	-0.5 (3)
C3—S1—C10—C8	176.7 (3)	C15—C16—C17—C18	-175.3 (2)
C3—S1—C10—C5	-1.4 (2)	C16—C17—C18—C19	-178.8 (3)
C3—C4—C11—C15	68.7 (3)	C16—C17—C18—S2	0.8 (3)
C5—C4—C11—C15	-114.7 (3)	C20—S2—C18—C17	-0.7 (2)
C3—C4—C11—C12	-103.1 (3)	C20—S2—C18—C19	178.9 (3)
C5—C4—C11—C12	73.5 (3)	C17—C18—C19—O1	-156.0 (19)
C15—C11—C12—F2	141.2 (2)	S2—C18—C19—O1	25 (2)
C4—C11—C12—F2	-45.4 (3)	C17—C18—C19—O1'	168.9 (15)
C15—C11—C12—F1	-100.3 (2)	S2—C18—C19—O1'	-10.6 (16)
C4—C11—C12—F1	73.1 (3)	C17—C16—C20—C21	-178.2 (2)

## supplementary materials

---

C15—C11—C12—C13	15.6 (3)	C15—C16—C20—C21	-3.6 (3)
C4—C11—C12—C13	-171.1 (2)	C17—C16—C20—S2	-0.1 (2)
F2—C12—C13—F3	86.1 (3)	C15—C16—C20—S2	174.49 (16)
F1—C12—C13—F3	-30.9 (3)	C18—S2—C20—C16	0.43 (17)
C11—C12—C13—F3	-147.6 (2)	C18—S2—C20—C21	178.80 (17)
F2—C12—C13—F4	-34.0 (3)	C16—C20—C21—C22	-117.1 (2)
F1—C12—C13—F4	-151.1 (2)	S2—C20—C21—C22	64.9 (2)
C11—C12—C13—F4	92.2 (2)	C20—C21—C22—C23	-173.65 (19)
F2—C12—C13—C14	-149.7 (2)	C21—C22—C23—C24	-175.0 (2)
F1—C12—C13—C14	93.3 (2)		

Fig. 1

