

1-Dichloroacetyl-*r*-2,*c*-6-bis(4-methoxyphenyl)-*c*-3,*t*-3-dimethylpiperidin-4-one

M. Thenmozhi,^a S. Ponnuswamy,^b V. Mohanraj,^b
R. Vijayalakshmi^c and M. N. Ponnuswamy^{a*}

^aCentre of Advanced Study in Crystallography and Biophysics, University of Madras, Guindy Campus, Chennai 600 025, India, ^bDepartment of Chemistry, Government Arts College (Autonomous), Coimbatore 641 018, India, and ^cDepartment of Chemistry, Queen Mary's College (Autonomous), Chennai 600 004, India
Correspondence e-mail: mnpsy2004@yahoo.com


Received 15 November 2008; accepted 27 November 2008

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.056; wR factor = 0.184; data-to-parameter ratio = 25.1.

In the title compound, $\text{C}_{23}\text{H}_{25}\text{Cl}_2\text{NO}_4$, the piperidine ring adopts a distorted boat conformation. Inversion-related molecules are linked into centrosymmetric $R_2^2(16)$ dimers by paired $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, and the dimers are connected via $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds into a chain running along [101].

Related literature

For general background, see: Eller *et al.* (2002); Ribeiro da Silva *et al.* (2007). For hybridization, see: Beddoes *et al.* (1986). For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For ring conformational analysis, see: Cremer & Pople (1975); Nardelli (1983).



Experimental

Crystal data

$\text{C}_{23}\text{H}_{25}\text{Cl}_2\text{NO}_4$	$V = 4508.5$ (3) Å ³
$M_r = 450.34$	$Z = 8$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 23.6295$ (9) Å	$\mu = 0.32$ mm ⁻¹
$b = 10.3999$ (4) Å	$T = 293$ (2) K
$c = 19.2617$ (9) Å	$0.30 \times 0.25 \times 0.20$ mm
$\beta = 107.734$ (1)°	

Data collection

Bruker Kappa APEXII area-detector diffractometer	29354 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 2001)	6905 independent reflections
$T_{\min} = 0.911$, $T_{\max} = 0.939$	4138 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$	275 parameters
$wR(F^2) = 0.184$	H-atom parameters constrained
$S = 1.00$	$\Delta\rho_{\text{max}} = 0.60$ e Å ⁻³
6905 reflections	$\Delta\rho_{\text{min}} = -0.63$ e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C6}-\text{H6}\cdots\text{O1}^i$	0.98	2.30	3.216 (2)	155
$\text{C13}-\text{H13}\cdots\text{O3}^{ii}$	0.93	2.58	3.453 (3)	155

Symmetry codes: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x + 1, -y + 1, -z + 1$.

Data collection: APEX2 (Bruker, 2004); cell refinement: APEX2; data reduction: SAINT (Bruker, 2004); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXS97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997); software used to prepare material for publication: SHELXL97 and PLATON (Spek, 2003).

SP thanks the UGC, India, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2726).

References

- Beddoes, R. L., Dalton, L., Joule, T. A., Mills, O. S., Street, J. D. & Watt, C. I. F. (1986). *J. Chem. Soc. Perkin Trans. 2*, pp. 787–797.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Eller, K., Henkes, E., Rossbacher, R. & Hoke, H. (2002). *Amines, Aliphatic*, in *Ullmann's Encyclopedia of Industrial Chemistry*, Vol. 1, p. 379. Weinheim: Wiley-VCH Verlag.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Nardelli, M. (1983). *Acta Cryst.* **C39**, 1141–1142.
- Ribeiro da Silva, M. A. V. & Cabral, J. I. T. A. (2007). *J. Therm. Anal. Calorim.* **90**, 865–871.
- Sheldrick, G. M. (2001). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2009). E65, o11 [doi:10.1107/S1600536808040063]

1-Dichloroacetyl-*r*-2,*c*-6-bis(4-methoxyphenyl)-*c*-3,*t*-3-dimethylpiperidin-4-one

M. Thenmozhi, S. Ponnuswamy, V. Mohanraj, R. Vijayalakshmi and M. N. Ponnuswamy

Comment

The piperidine compounds are used in chemical industry for the synthesis of pharmacological drugs, either as reactants, solvents or being units of molecular chemical structure of the final compounds. These compounds have significant biological importance with more environmental impact (Ribeiro da Silva *et al.*, 2007). A significant industrial application of piperidine is for the production of dipiperidinyl dithiuram tetrasulfide, which is used as a rubber vulcanization accelerator (Eller *et al.*, 2002).

The piperidine ring adopts a distorted boat conformation, with puckering parameters (Cremer & Pople, 1975) $q_2 = 0.606$ (2) Å, $q_3 = 0.127$ (2) Å, and $\varphi = 77.5$ (2)°, and asymmetry parameter $\Delta_s(C2) = 18.26$ (17)° (Nardelli, 1983). The torsion angles C13—C12—O2—C15 [2.6 (3)°] and C20—C21—O4—C24 [-11.7 (3)°] indicate that the methoxy groups are almost coplanar with the attached rings (Fig. 1). The sum of the bond angles around atom N1 (359.4°) of the piperidine ring is in accordance with sp^2 hybridization (Beddoes *et al.*, 1986). The best plane through the piperidine ring (N1/C3/C4/C6) and methoxyphenyl ring (C9—C14) are orthogonal to one another, with a dihedral angle of 87.18 (7)°, whereas, the other methoxyphenyl ring (C18—C23) is oriented at an angle of 54.47 (7)°.

The molecules at positions (x, y, z) and $(1 - x, 1 - y, 1 - z)$ are linked through a pair of C13—H13 \cdots O3 hydrogen bonds forming a cyclic centrosymmetric $R_2^2(16)$ dimer (Bernstein *et al.*, 1995). The dimers are linked by intermolecular C6—H6 \cdots O1 hydrogen bonds (Table 1) into a chain running along the [101] (Fig. 2).

Experimental

To the solution of *r*-2,*c*-6-bis(4-methoxyphenyl)-*c*-3,*t*-3-dimethylpiperidin-4-one (2 g) in benzene (25 ml), triethylamine (2.0 ml) and dichloroacetyl chloride (1.40 ml) were added and allowed to reflux on a water bath for 5 h. The course of the reaction was monitored by TLC. The solution was concentrated and the resulting mass was crystallized from ethanol.

Refinement

H atoms were positioned geometrically (C—H = 0.93–0.98 Å) and allowed to ride on their parent atoms, with $U_{iso}(H) = 1.2$ – 1.5 (methyl) $U_{eq}(C)$.

Figures




Fig. 1. The molecular structure of the title compound, showing 50% probability displacement ellipsoids.




Fig. 2. The crystal packing of the molecules viewed down *b* axis. Hydrogen bonds are shown as dashed lines.

1-Dichloroacetyl-r-2,c-6-bis(4-methoxyphenyl)-c-3,t-3-dimethylpiperidin-4-one

Crystal data

$C_{23}H_{25}Cl_2NO_4$

$M_r = 450.34$

Monoclinic, $C2/c$

Hall symbol: $-C\ 2yc$

$a = 23.6295\ (9)\ \text{\AA}$

$b = 10.3999\ (4)\ \text{\AA}$

$c = 19.2617\ (9)\ \text{\AA}$

$\beta = 107.734\ (1)^\circ$

$V = 4508.5\ (3)\ \text{\AA}^3$

$Z = 8$

$F_{000} = 1888$

$D_x = 1.327\ \text{Mg m}^{-3}$

Melting point = $377\text{--}379\ \text{K}$

Mo $K\alpha$ radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 6905 reflections

$\theta = 2.2\text{--}30.6^\circ$

$\mu = 0.32\ \text{mm}^{-1}$

$T = 293\ (2)\ \text{K}$

Block, colourless

$0.30 \times 0.25 \times 0.20\ \text{mm}$

Data collection

Bruker Kappa APEXII area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293\ (2)\ \text{K}$

ω and φ scans

Absorption correction: multi-scan (SADABS; Sheldrick, 2001)

$T_{\min} = 0.911$, $T_{\max} = 0.939$

29354 measured reflections

6905 independent reflections

4138 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.031$

$\theta_{\text{max}} = 30.6^\circ$

$\theta_{\text{min}} = 2.2^\circ$

$h = -32 \rightarrow 33$

$k = -14 \rightarrow 14$

$l = -27 \rightarrow 27$

Refinement

Refinement on F^2

Secondary atom site location: difference Fourier map

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.056$$

$$wR(F^2) = 0.184$$

$$S = 1.01$$

6905 reflections

275 parameters

Primary atom site location: structure-invariant direct methods

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0869P)^2 + 2.9239P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.60 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.63 \text{ e } \text{\AA}^{-3}$$

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.18029 (3)	0.52254 (7)	0.14172 (4)	0.0790 (2)
C12	0.26274 (4)	0.62339 (11)	0.07261 (4)	0.1127 (4)
O1	0.26407 (7)	0.74756 (16)	0.21261 (9)	0.0635 (4)
O2	0.56012 (6)	0.89254 (16)	0.36906 (9)	0.0604 (4)
O3	0.37036 (9)	0.4204 (2)	0.48165 (10)	0.0798 (6)
O4	0.48349 (7)	0.30777 (15)	0.10880 (9)	0.0592 (4)
N1	0.31916 (6)	0.58476 (15)	0.27717 (8)	0.0402 (3)
C2	0.33601 (8)	0.66580 (19)	0.34377 (10)	0.0407 (4)
H2	0.3092	0.7400	0.3325	0.049*
C3	0.32172 (8)	0.5947 (2)	0.40675 (11)	0.0481 (5)
C4	0.35641 (9)	0.4707 (2)	0.42265 (12)	0.0530 (5)
C5	0.37393 (9)	0.4121 (2)	0.36076 (12)	0.0521 (5)
H5A	0.3721	0.3194	0.3651	0.063*
H5B	0.4151	0.4343	0.3676	0.063*
C6	0.33818 (8)	0.44832 (19)	0.28249 (11)	0.0435 (4)
H6	0.3023	0.3950	0.2685	0.052*
C7	0.28061 (8)	0.6362 (2)	0.21725 (11)	0.0460 (4)
C8	0.25568 (9)	0.5486 (2)	0.15090 (12)	0.0550 (5)
H8	0.2771	0.4665	0.1587	0.066*
C9	0.39837 (8)	0.72041 (18)	0.35651 (10)	0.0402 (4)
C10	0.40476 (8)	0.81147 (19)	0.30699 (10)	0.0431 (4)
H10	0.3716	0.8357	0.2689	0.052*

supplementary materials

C11	0.45865 (9)	0.8669 (2)	0.31249 (11)	0.0476 (5)
H11	0.4616	0.9282	0.2786	0.057*
C12	0.50864 (8)	0.83169 (19)	0.36834 (11)	0.0453 (4)
C13	0.50374 (8)	0.7419 (2)	0.41826 (11)	0.0475 (5)
H13	0.5371	0.7176	0.4560	0.057*
C14	0.44863 (8)	0.6872 (2)	0.41224 (11)	0.0474 (5)
H14	0.4456	0.6268	0.4466	0.057*
C15	0.61272 (11)	0.8573 (4)	0.4232 (2)	0.1002 (11)
H15A	0.6096	0.8793	0.4703	0.150*
H15B	0.6457	0.9022	0.4154	0.150*
H15C	0.6188	0.7663	0.4210	0.150*
C16	0.25568 (9)	0.5548 (3)	0.38209 (14)	0.0630 (6)
H16A	0.2487	0.4941	0.3428	0.095*
H16B	0.2313	0.6293	0.3659	0.095*
H16C	0.2459	0.5160	0.4222	0.095*
C17	0.33296 (11)	0.6809 (3)	0.47360 (12)	0.0646 (6)
H17A	0.3258	0.6332	0.5128	0.097*
H17B	0.3067	0.7536	0.4621	0.097*
H17C	0.3734	0.7101	0.4881	0.097*
C18	0.37492 (8)	0.41315 (18)	0.23272 (11)	0.0416 (4)
C19	0.37935 (9)	0.2855 (2)	0.21607 (12)	0.0498 (5)
H19	0.3576	0.2247	0.2326	0.060*
C20	0.41538 (9)	0.2454 (2)	0.17531 (12)	0.0507 (5)
H20	0.4180	0.1587	0.1650	0.061*
C21	0.44721 (8)	0.3353 (2)	0.15026 (11)	0.0458 (4)
C22	0.44278 (9)	0.4630 (2)	0.16603 (13)	0.0533 (5)
H22	0.4640	0.5240	0.1488	0.064*
C23	0.40738 (9)	0.5018 (2)	0.20705 (13)	0.0508 (5)
H23	0.4052	0.5885	0.2177	0.061*
C24	0.49748 (14)	0.1779 (2)	0.10196 (17)	0.0750 (7)
H24A	0.5128	0.1402	0.1495	0.113*
H24B	0.5269	0.1721	0.0770	0.113*
H24C	0.4623	0.1326	0.0747	0.113*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0512 (3)	0.0880 (5)	0.0926 (5)	-0.0197 (3)	0.0140 (3)	0.0036 (4)
C12	0.1188 (7)	0.1699 (10)	0.0577 (4)	-0.0632 (6)	0.0394 (4)	-0.0210 (5)
O1	0.0641 (9)	0.0589 (10)	0.0558 (9)	0.0225 (8)	0.0008 (7)	0.0010 (7)
O2	0.0427 (7)	0.0669 (10)	0.0689 (10)	-0.0084 (7)	0.0130 (7)	0.0073 (8)
O3	0.0847 (13)	0.0963 (14)	0.0611 (11)	0.0211 (11)	0.0265 (9)	0.0364 (10)
O4	0.0632 (9)	0.0547 (9)	0.0686 (10)	0.0045 (7)	0.0332 (8)	-0.0007 (8)
N1	0.0348 (7)	0.0415 (8)	0.0421 (8)	0.0034 (6)	0.0087 (6)	0.0038 (7)
C2	0.0355 (8)	0.0449 (10)	0.0413 (9)	0.0042 (7)	0.0110 (7)	0.0027 (8)
C3	0.0389 (9)	0.0617 (13)	0.0448 (10)	0.0023 (8)	0.0146 (8)	0.0073 (9)
C4	0.0407 (9)	0.0641 (14)	0.0543 (12)	0.0010 (9)	0.0148 (9)	0.0185 (10)
C5	0.0486 (10)	0.0480 (12)	0.0609 (13)	0.0096 (9)	0.0183 (9)	0.0164 (10)

C6	0.0369 (8)	0.0398 (10)	0.0541 (11)	-0.0008 (7)	0.0143 (8)	0.0053 (8)
C7	0.0381 (9)	0.0522 (12)	0.0449 (10)	0.0084 (8)	0.0084 (8)	0.0022 (9)
C8	0.0396 (9)	0.0657 (14)	0.0514 (12)	0.0050 (9)	0.0015 (8)	-0.0046 (10)
C9	0.0384 (8)	0.0412 (10)	0.0408 (9)	0.0021 (7)	0.0117 (7)	0.0000 (8)
C10	0.0394 (9)	0.0470 (11)	0.0412 (9)	0.0023 (8)	0.0097 (7)	0.0009 (8)
C11	0.0477 (10)	0.0495 (11)	0.0462 (11)	-0.0009 (8)	0.0151 (8)	0.0052 (9)
C12	0.0398 (9)	0.0460 (11)	0.0505 (11)	-0.0016 (8)	0.0145 (8)	-0.0059 (9)
C13	0.0386 (9)	0.0509 (12)	0.0479 (11)	0.0033 (8)	0.0057 (8)	0.0023 (9)
C14	0.0426 (9)	0.0511 (12)	0.0457 (10)	0.0006 (8)	0.0093 (8)	0.0080 (9)
C15	0.0412 (12)	0.117 (3)	0.126 (3)	-0.0143 (14)	0.0005 (14)	0.044 (2)
C16	0.0400 (10)	0.0862 (18)	0.0664 (14)	-0.0007 (10)	0.0214 (10)	0.0106 (13)
C17	0.0607 (13)	0.0883 (18)	0.0478 (12)	0.0039 (12)	0.0212 (10)	0.0006 (12)
C18	0.0352 (8)	0.0387 (10)	0.0502 (10)	0.0000 (7)	0.0119 (7)	0.0025 (8)
C19	0.0501 (10)	0.0419 (11)	0.0592 (12)	-0.0088 (8)	0.0193 (9)	0.0019 (9)
C20	0.0554 (11)	0.0383 (11)	0.0584 (12)	-0.0018 (9)	0.0172 (10)	-0.0034 (9)
C21	0.0419 (9)	0.0483 (11)	0.0463 (10)	0.0025 (8)	0.0122 (8)	0.0024 (9)
C22	0.0521 (11)	0.0441 (11)	0.0693 (14)	-0.0048 (9)	0.0268 (10)	0.0041 (10)
C23	0.0504 (11)	0.0363 (10)	0.0708 (14)	-0.0005 (8)	0.0258 (10)	0.0009 (9)
C24	0.0880 (18)	0.0611 (16)	0.0897 (19)	0.0135 (13)	0.0475 (16)	-0.0023 (14)

Geometric parameters (Å, °)

C11—C8	1.757 (2)	C11—C12	1.383 (3)
C12—C8	1.749 (2)	C11—H11	0.93
O1—C7	1.217 (2)	C12—C13	1.370 (3)
O2—C12	1.368 (2)	C13—C14	1.393 (3)
O2—C15	1.406 (3)	C13—H13	0.93
O3—C4	1.202 (3)	C14—H14	0.93
O4—C21	1.368 (2)	C15—H15A	0.96
O4—C24	1.406 (3)	C15—H15B	0.96
N1—C7	1.344 (2)	C15—H15C	0.96
N1—C6	1.483 (2)	C16—H16A	0.96
N1—C2	1.484 (2)	C16—H16B	0.96
C2—C9	1.528 (2)	C16—H16C	0.96
C2—C3	1.544 (3)	C17—H17A	0.96
C2—H2	0.98	C17—H17B	0.96
C3—C4	1.508 (3)	C17—H17C	0.96
C3—C17	1.524 (3)	C18—C19	1.377 (3)
C3—C16	1.543 (3)	C18—C23	1.383 (3)
C4—C5	1.505 (3)	C19—C20	1.387 (3)
C5—C6	1.534 (3)	C19—H19	0.93
C5—H5A	0.97	C20—C21	1.376 (3)
C5—H5B	0.97	C20—H20	0.93
C6—C18	1.521 (3)	C21—C22	1.373 (3)
C6—H6	0.98	C22—C23	1.374 (3)
C7—C8	1.533 (3)	C22—H22	0.93
C8—H8	0.98	C23—H23	0.93
C9—C14	1.380 (3)	C24—H24A	0.96
C9—C10	1.385 (3)	C24—H24B	0.96

supplementary materials

C10—C11	1.372 (3)	C24—H24C	0.96
C10—H10	0.93		
C12—O2—C15	117.99 (19)	O2—C12—C11	115.57 (18)
C21—O4—C24	117.58 (18)	C13—C12—C11	119.45 (18)
C7—N1—C6	123.51 (16)	C12—C13—C14	119.78 (18)
C7—N1—C2	116.70 (15)	C12—C13—H13	120.1
C6—N1—C2	119.19 (15)	C14—C13—H13	120.1
N1—C2—C9	109.97 (14)	C9—C14—C13	121.55 (19)
N1—C2—C3	109.99 (16)	C9—C14—H14	119.2
C9—C2—C3	118.89 (16)	C13—C14—H14	119.2
N1—C2—H2	105.7	O2—C15—H15A	109.5
C9—C2—H2	105.7	O2—C15—H15B	109.5
C3—C2—H2	105.7	H15A—C15—H15B	109.5
C4—C3—C17	112.76 (18)	O2—C15—H15C	109.5
C4—C3—C16	105.59 (19)	H15A—C15—H15C	109.5
C17—C3—C16	108.54 (17)	H15B—C15—H15C	109.5
C4—C3—C2	109.49 (15)	C3—C16—H16A	109.5
C17—C3—C2	111.01 (18)	C3—C16—H16B	109.5
C16—C3—C2	109.25 (17)	H16A—C16—H16B	109.5
O3—C4—C5	120.7 (2)	C3—C16—H16C	109.5
O3—C4—C3	122.7 (2)	H16A—C16—H16C	109.5
C5—C4—C3	116.68 (17)	H16B—C16—H16C	109.5
C4—C5—C6	118.50 (17)	C3—C17—H17A	109.5
C4—C5—H5A	107.7	C3—C17—H17B	109.5
C6—C5—H5A	107.7	H17A—C17—H17B	109.5
C4—C5—H5B	107.7	C3—C17—H17C	109.5
C6—C5—H5B	107.7	H17A—C17—H17C	109.5
H5A—C5—H5B	107.1	H17B—C17—H17C	109.5
N1—C6—C18	113.94 (15)	C19—C18—C23	117.99 (18)
N1—C6—C5	111.57 (16)	C19—C18—C6	118.50 (17)
C18—C6—C5	108.14 (15)	C23—C18—C6	123.37 (18)
N1—C6—H6	107.6	C18—C19—C20	121.69 (18)
C18—C6—H6	107.6	C18—C19—H19	119.2
C5—C6—H6	107.6	C20—C19—H19	119.2
O1—C7—N1	124.06 (19)	C21—C20—C19	119.30 (19)
O1—C7—C8	118.08 (18)	C21—C20—H20	120.3
N1—C7—C8	117.85 (18)	C19—C20—H20	120.3
C7—C8—C12	109.96 (16)	O4—C21—C22	115.80 (18)
C7—C8—C11	107.21 (15)	O4—C21—C20	124.66 (19)
C12—C8—C11	110.00 (12)	C22—C21—C20	119.54 (18)
C7—C8—H8	109.9	C21—C22—C23	120.75 (19)
C12—C8—H8	109.9	C21—C22—H22	119.6
C11—C8—H8	109.9	C23—C22—H22	119.6
C14—C9—C10	117.29 (17)	C22—C23—C18	120.72 (19)
C14—C9—C2	126.23 (17)	C22—C23—H23	119.6
C10—C9—C2	116.48 (16)	C18—C23—H23	119.6
C11—C10—C9	121.84 (18)	O4—C24—H24A	109.5
C11—C10—H10	119.1	O4—C24—H24B	109.5
C9—C10—H10	119.1	H24A—C24—H24B	109.5

C10—C11—C12	120.10 (19)	O4—C24—H24C	109.5
C10—C11—H11	120.0	H24A—C24—H24C	109.5
C12—C11—H11	120.0	H24B—C24—H24C	109.5
O2—C12—C13	124.98 (18)		
C7—N1—C2—C9	-104.12 (18)	N1—C2—C9—C14	-110.8 (2)
C6—N1—C2—C9	84.47 (19)	C3—C2—C9—C14	17.3 (3)
C7—N1—C2—C3	123.10 (17)	N1—C2—C9—C10	68.9 (2)
C6—N1—C2—C3	-48.31 (19)	C3—C2—C9—C10	-163.05 (18)
N1—C2—C3—C4	60.4 (2)	C14—C9—C10—C11	0.1 (3)
C9—C2—C3—C4	-67.6 (2)	C2—C9—C10—C11	-179.62 (18)
N1—C2—C3—C17	-174.47 (15)	C9—C10—C11—C12	0.4 (3)
C9—C2—C3—C17	57.5 (2)	C15—O2—C12—C13	2.6 (3)
N1—C2—C3—C16	-54.8 (2)	C15—O2—C12—C11	-177.9 (3)
C9—C2—C3—C16	177.18 (18)	C10—C11—C12—O2	-179.95 (18)
C17—C3—C4—O3	29.1 (3)	C10—C11—C12—C13	-0.5 (3)
C16—C3—C4—O3	-89.2 (3)	O2—C12—C13—C14	179.43 (19)
C2—C3—C4—O3	153.3 (2)	C11—C12—C13—C14	0.0 (3)
C17—C3—C4—C5	-150.11 (19)	C10—C9—C14—C13	-0.6 (3)
C16—C3—C4—C5	91.5 (2)	C2—C9—C14—C13	179.11 (18)
C2—C3—C4—C5	-26.0 (2)	C12—C13—C14—C9	0.5 (3)
O3—C4—C5—C6	158.4 (2)	N1—C6—C18—C19	-160.22 (17)
C3—C4—C5—C6	-22.3 (3)	C5—C6—C18—C19	75.1 (2)
C7—N1—C6—C18	66.7 (2)	N1—C6—C18—C23	24.2 (3)
C2—N1—C6—C18	-122.54 (17)	C5—C6—C18—C23	-100.5 (2)
C7—N1—C6—C5	-170.53 (17)	C23—C18—C19—C20	0.4 (3)
C2—N1—C6—C5	0.3 (2)	C6—C18—C19—C20	-175.39 (19)
C4—C5—C6—N1	36.3 (2)	C18—C19—C20—C21	-0.5 (3)
C4—C5—C6—C18	162.32 (18)	C24—O4—C21—C22	169.2 (2)
C6—N1—C7—O1	178.95 (18)	C24—O4—C21—C20	-11.7 (3)
C2—N1—C7—O1	7.9 (3)	C19—C20—C21—O4	-179.0 (2)
C6—N1—C7—C8	-0.6 (3)	C19—C20—C21—C22	0.0 (3)
C2—N1—C7—C8	-171.63 (16)	O4—C21—C22—C23	179.7 (2)
O1—C7—C8—C12	49.9 (2)	C20—C21—C22—C23	0.6 (3)
N1—C7—C8—C12	-130.51 (17)	C21—C22—C23—C18	-0.7 (4)
O1—C7—C8—C11	-69.7 (2)	C19—C18—C23—C22	0.2 (3)
N1—C7—C8—C11	109.92 (18)	C6—C18—C23—C22	175.8 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C6—H6...O1 ⁱ	0.98	2.30	3.216 (2)	155
C13—H13...O3 ⁱⁱ	0.93	2.58	3.453 (3)	155

Symmetry codes: (i) $-x+1/2, y-1/2, -z+1/2$; (ii) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2

