

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 2,4-Difluorophenylboronic acid

Patricia Rodríguez-Cuamatzi,<sup>a</sup> Hugo Tlahuext<sup>b</sup> and Herbert Höpfl<sup>b\*</sup>

<sup>a</sup>Universidad Politécnica de Tlaxcala, Carretera Federal Tlaxcala-Puebla Km 9.5, Tepeyanco, Tlaxcala, Mexico, and <sup>b</sup>Centro de Investigaciones Químicas, Universidad Autónoma del Estado de Morelos. Av. Universidad 1001 Col., Chamilpa, CP 62209, Cuernavaca Mor., Mexico

Correspondence e-mail: hhopfl@uaem.mx


Received 24 November 2008; accepted 3 December 2008

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.056;  $wR$  factor = 0.127; data-to-parameter ratio = 11.2.

The molecular structure of the title compound,  $\text{C}_6\text{H}_5\text{BF}_2\text{O}_2$ , is essentially planar (mean deviation = 0.019 Å), indicating electronic delocalization between the dihydroxyboryl group and the aromatic ring. In the crystal structure, inversion dimers linked by two  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds arise. An intramolecular  $\text{O}-\text{H}\cdots\text{F}$  hydrogen bond reinforces the conformation and the same H atom is also involved in an intermolecular  $\text{O}-\text{H}\cdots\text{F}$  link, leading to molecular sheets in the crystal.

## Related literature

For general background to boronic acids, see: Hall (2005); Höpfl (2002); Fujita *et al.* (2008); Soloway *et al.* (1998). For hydrogen-bond motifs, see: Bernstein *et al.* (1995); Desiraju (2002). For related structures, see: Wu *et al.* (2006); Bradley *et al.* (1996); Horton *et al.* (2004). For crystal engineering, see: Fournier *et al.* (2003); Rodríguez-Cuamatzi *et al.* (2004, 2005).



## Experimental

## Crystal data

$\text{C}_6\text{H}_5\text{BF}_2\text{O}_2$   
 $M_r = 157.91$   
 Monoclinic,  $P2_1/n$   
 $a = 3.7617$  (11) Å  
 $b = 12.347$  (4) Å  
 $c = 14.620$  (4) Å  
 $\beta = 95.450$  (5)°

$V = 676.0$  (3) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.15$  mm<sup>-1</sup>  
 $T = 293$  (2) K  
 $0.37 \times 0.35 \times 0.22$  mm

## Data collection

Bruker SMART APEX CCD area-detector diffractometer  
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.947$ ,  $T_{\max} = 0.968$

3196 measured reflections  
 1190 independent reflections  
 1012 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.028$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$   
 $wR(F^2) = 0.127$   
 $S = 1.15$   
 1190 reflections  
 106 parameters  
 2 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.14$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.18$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1}\cdots\text{F1}$	0.841 (15)	2.16 (3)	2.799 (3)	133 (2)
$\text{O1}-\text{H1}\cdots\text{F2}^i$	0.841 (15)	2.39 (2)	3.086 (3)	140 (3)
$\text{O2}-\text{H2}\cdots\text{O1}^{ii}$	0.841 (19)	1.97 (2)	2.809 (3)	174 (3)

Symmetry codes: (i)  $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$ ; (ii)  $-x + 1, -y + 1, -z + 1$ .

Data collection: SMART (Bruker, 2000); cell refinement: SAINT-Plus-NT (Bruker, 2001); data reduction: SAINT-Plus-NT; program(s) used to solve structure: SHELXTL-NT (Sheldrick, 2008); program(s) used to refine structure: SHELXTL-NT; molecular graphics: CAMERON (Watkin *et al.*, 1996); software used to prepare material for publication: PLATON (Spek, 2003) and publCIF (Westrip, 2009).

This work was supported by Consejo Nacional de Ciencia y Tecnología (CIAM-59213 for HH).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2865).

## References

- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bradley, D. C., Harding, I. S., Keefe, A. D., Motevalli, M. & Zheng, D. H. (1996). *J. Chem. Soc. Dalton Trans.* pp. 3931–3936.
- Bruker (2000). SMART. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2001). SAINT-Plus NT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Desiraju, G. R. (2002). *Acc. Chem. Res.* **35**, 565–573.
- Fournier, J.-H., Maris, T., Wuest, J. D., Guo, W. & Galoppini, E. (2003). *J. Am. Chem. Soc.* **125**, 1002–1006.
- Fujita, N., Shinkai, S. & James, T. D. (2008). *Chem. Asian J.* **3**, 1076–1091.
- Hall, D. G. (2005). *Boronic Acids: Preparation, Applications in Organic Synthesis and Medicine*. Weinheim: Wiley-VCH.
- Höpfl, H. (2002). *Structure and Bonding*, edited by H. W. Roesky & D. A. Atwood, Vol. 103, pp. 1–56. Berlin: Springer Verlag.
- Horton, P. N., Hursthouse, M. B., Beckett, M. A. & Rugen-Hankey, M. P. (2004). *Acta Cryst. E* **60**, o2204–o2206.
- Rodríguez-Cuamatzi, P., Arillo-Flores, O. I., Bernal-Uruchurtu, M. I. & Höpfl, H. (2005). *Cryst. Growth Des.* **5**, 167–175.
- Rodríguez-Cuamatzi, P., Vargas-Díaz, G. & Höpfl, H. (2004). *Angew. Chem. Int. Ed.* **43**, 3041–3044.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

Soloway, A. H., Tjarks, W., Barnum, B. A., Rong, R.-A., Barth, R. F., Codogni, I. M. & Wilson, J. G. (1998). *Chem. Rev.* **98**, 1515–1562.  
Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.  
Watkin, D. J., Prout, C. K. & Pearce, L. J. (1996). *CAMERON*. Chemical Crystallography Laboratory Oxford, Oxford, England.

Westrip, S. P. (2009). *pubCIF*. In preparation.  
Wu, Y.-M., Dong, C.-C., Liu, S., Zhu, H.-J. & Wu, Y.-Z. (2006). *Acta Cryst.* **E62**, o4236–o4237.

**supplementary materials**

*Acta Cryst.* (2009). E65, o44-o45 [ doi:10.1107/S1600536808040646 ]

## 2,4-Difluorophenylboronic acid

P. Rodríguez-Cuamatzi, H. Tlahuext and H. Höpfl

### Comment

Boronic acids,  $\text{RB}(\text{OH})_2$  with  $R$  = alkyl and aryl, have applications in organic synthesis (Hall, 2005), host–guest chemistry (Höpfl, 2002), the molecular recognition of biochemically active molecules (Fujita *et al.*, 2008) and in medicine as antibiotics, inhibitors and for the treatment of tumors (Soloway *et al.*, 1998). Similar to carboxylic acids they are capable to form hydrogen-bonded dimeric units and, therefore, boronic acids have been used recently as new building blocks in crystal engineering (Fournier *et al.*, 2003; Rodríguez-Cuamatzi *et al.*, 2004; Rodríguez-Cuamatzi *et al.*, 2005). Previously, the structures of 3-fluorophenylboronic acid (Wu *et al.*, 2006), 2,6-difluoroboronic acid (Bradley *et al.*, 1996) and pentafluoroboronic acid (Horton *et al.*, 2004) had been reported. We now present the crystal structure of (**I**).

The molecular structure is essentially planar,  $\text{O1—B1—C1—C2} = 4.4$  (4) $^\circ$ , indicating that there is a  $\pi\cdots\pi$  interaction between the dihydroxyboryl group and the aromatic ring, to which it is attached (Fig. 1). This interaction is also evidenced by the B—C bond length of 1.566 (3) Å, which is significantly shorter than that observed in boronates containing tetra-coordinate boron atoms (Höpfl, 2002). The crystal structure is stabilized by strong  $\text{O2—H2}\cdots\text{O1}$  hydrogen-bonding interactions, forming  $R_2^2(8)$  motifs (Bernstein *et al.*, 1995), as well as,  $\text{O1—H1}\cdots\text{F1}$  and  $\text{O1—H1}\cdots\text{F2}$  bifurcated hydrogen bonds (Fig. 2; Table 1) (Desiraju, 2002). Due to these interactions each boronic acid homodimer is linked to two neighboring homodimeric units, thus creating a two-dimensional hydrogen-bonded network, in which fluorine is therefore an essential structural component.


### Experimental


2,4-Difluorophenylboronic acid was purchased from Aldrich and crystallized from water to yield colourless blocks of (**I**).

### Refinement

The aromatic H atoms were positioned geometrically ( $\text{C—H} = 0.93\text{Å}$ ) and refined as riding with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . The O—H hydrogen atoms were localized in a difference map and their coordinates were refined with  $\text{O—H} = 0.84\pm 0.01\text{Å}$  and  $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$ .

### Figures

 Fig. 1. The molecular structure of (**I**) with displacement ellipsoids drawn at the 30% probability level and H atoms shown as small spheres of arbitrary radius.

 Fig. 2. View of the packing arrangement of the two-dimensional network of (**I**)(**I**).

## 2,4-Difluorophenylboronic acid

### Crystal data

$C_6H_5BF_2O_2$	$F_{000} = 320$
$M_r = 157.91$	$D_x = 1.552 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Melting point = 521–522 K
Hall symbol: -P 2yn	Mo $K\alpha$ radiation
$a = 3.7617 (11) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 12.347 (4) \text{ \AA}$	Cell parameters from 1052 reflections
$c = 14.620 (4) \text{ \AA}$	$\theta = 2.3\text{--}26.2^\circ$
$\beta = 95.450 (5)^\circ$	$\mu = 0.15 \text{ mm}^{-1}$
$V = 676.0 (3) \text{ \AA}^3$	$T = 293 (2) \text{ K}$
$Z = 4$	Block, colorless
	$0.37 \times 0.35 \times 0.22 \text{ mm}$

### Data collection

Bruker SMART APEX CCD area-detector diffractometer	1190 independent reflections
Radiation source: fine-focus sealed tube	1012 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.028$
Detector resolution: 8.3 pixels $\text{mm}^{-1}$	$\theta_{\text{max}} = 25.0^\circ$
$T = 293(2) \text{ K}$	$\theta_{\text{min}} = 2.2^\circ$
$\varphi$ and $\omega$ scans	$h = -3 \rightarrow 4$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$k = -14 \rightarrow 12$
$T_{\text{min}} = 0.947$ , $T_{\text{max}} = 0.968$	$l = -17 \rightarrow 17$
3196 measured reflections	

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.056$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.127$	$w = 1/[\sigma^2(F_o^2) + (0.0442P)^2 + 0.2673P]$
$S = 1.15$	where $P = (F_o^2 + 2F_c^2)/3$
1190 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
106 parameters	$\Delta\rho_{\text{max}} = 0.14 \text{ e \AA}^{-3}$
2 restraints	$\Delta\rho_{\text{min}} = -0.18 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
B1	0.7704 (8)	0.4548 (2)	0.62567 (18)	0.0452 (7)
O1	0.6825 (6)	0.38623 (15)	0.55419 (13)	0.0695 (6)
H1	0.748 (9)	0.3215 (9)	0.562 (2)	0.104*
O2	0.6880 (6)	0.55977 (14)	0.61557 (12)	0.0630 (6)
H2	0.593 (8)	0.577 (3)	0.5632 (10)	0.094*
F1	1.0385 (5)	0.23728 (11)	0.67473 (11)	0.0768 (6)
F2	1.4329 (5)	0.33016 (14)	0.97634 (10)	0.0822 (6)
C1	0.9591 (6)	0.41789 (18)	0.72069 (15)	0.0424 (6)
C2	1.0796 (7)	0.31430 (18)	0.74175 (16)	0.0470 (6)
C3	1.2380 (7)	0.2819 (2)	0.82553 (17)	0.0539 (7)
H3	1.3138	0.2109	0.8363	0.065*
C4	1.2785 (7)	0.3593 (2)	0.89223 (17)	0.0547 (7)
C5	1.1696 (8)	0.4640 (2)	0.87828 (17)	0.0586 (7)
H5	1.2013	0.5150	0.9251	0.070*
C6	1.0119 (7)	0.49169 (19)	0.79282 (16)	0.0498 (6)
H6	0.9371	0.5628	0.7827	0.060*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
B1	0.0451 (16)	0.0425 (15)	0.0471 (16)	-0.0009 (12)	-0.0005 (12)	0.0033 (12)
O1	0.1000 (17)	0.0484 (11)	0.0540 (11)	0.0158 (10)	-0.0241 (10)	-0.0009 (9)
O2	0.0850 (15)	0.0441 (10)	0.0552 (11)	0.0087 (9)	-0.0172 (10)	0.0048 (8)
F1	0.1192 (15)	0.0456 (9)	0.0602 (10)	0.0151 (9)	-0.0187 (9)	-0.0046 (7)
F2	0.1070 (14)	0.0804 (12)	0.0527 (10)	-0.0120 (10)	-0.0262 (9)	0.0181 (8)
C1	0.0383 (13)	0.0412 (13)	0.0473 (13)	-0.0039 (10)	0.0019 (10)	0.0048 (10)
C2	0.0523 (16)	0.0410 (13)	0.0467 (13)	-0.0017 (11)	0.0001 (11)	0.0008 (10)
C3	0.0584 (17)	0.0449 (14)	0.0564 (15)	0.0002 (12)	-0.0053 (13)	0.0124 (12)
C4	0.0579 (17)	0.0613 (17)	0.0426 (13)	-0.0099 (13)	-0.0075 (12)	0.0135 (12)
C5	0.0696 (19)	0.0552 (16)	0.0489 (14)	-0.0109 (13)	-0.0058 (13)	-0.0020 (12)
C6	0.0559 (16)	0.0399 (13)	0.0522 (14)	-0.0011 (11)	-0.0011 (12)	0.0029 (11)

## supplementary materials

---

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

B1—O2	1.338 (3)	C1—C6	1.394 (3)
B1—O1	1.361 (3)	C2—C3	1.370 (3)
B1—C1	1.566 (3)	C3—C4	1.363 (4)
O1—H1	0.841 (15)	C3—H3	0.93
O2—H2	0.841 (15)	C4—C5	1.366 (4)
F1—C2	1.364 (3)	C5—C6	1.374 (3)
F2—C4	1.358 (3)	C5—H5	0.93
C1—C2	1.382 (3)	C6—H6	0.93
O2—B1—O1	118.7 (2)	C4—C3—H3	121.8
O2—B1—C1	117.4 (2)	C2—C3—H3	121.8
O1—B1—C1	123.8 (2)	F2—C4—C3	118.1 (2)
B1—O1—H1	116 (2)	F2—C4—C5	118.8 (2)
B1—O2—H2	115 (2)	C3—C4—C5	123.0 (2)
C2—C1—C6	114.6 (2)	C4—C5—C6	117.9 (2)
C2—C1—B1	125.3 (2)	C4—C5—H5	121.0
C6—C1—B1	120.1 (2)	C6—C5—H5	121.0
F1—C2—C3	116.7 (2)	C5—C6—C1	122.9 (2)
F1—C2—C1	118.2 (2)	C5—C6—H6	118.5
C3—C2—C1	125.1 (2)	C1—C6—H6	118.5
C4—C3—C2	116.4 (2)		
O2—B1—C1—C2	-176.5 (2)	C1—C2—C3—C4	-0.3 (4)
O1—B1—C1—C2	4.5 (4)	C2—C3—C4—F2	179.7 (2)
O2—B1—C1—C6	4.6 (4)	C2—C3—C4—C5	0.0 (4)
O1—B1—C1—C6	-174.5 (2)	F2—C4—C5—C6	-179.6 (2)
C6—C1—C2—F1	-179.9 (2)	C3—C4—C5—C6	0.1 (4)
B1—C1—C2—F1	1.1 (4)	C4—C5—C6—C1	0.0 (4)
C6—C1—C2—C3	0.4 (4)	C2—C1—C6—C5	-0.3 (4)
B1—C1—C2—C3	-178.6 (2)	B1—C1—C6—C5	178.8 (2)
F1—C2—C3—C4	180.0 (2)		

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1 $\cdots$ F1	0.841 (15)	2.16 (3)	2.799 (3)	133 (2)
O1—H1 $\cdots$ F2 <sup>i</sup>	0.841 (15)	2.39 (2)	3.086 (3)	140 (3)
O2—H2 $\cdots$ O1 <sup>ii</sup>	0.841 (19)	1.97 (2)	2.809 (3)	174 (3)

Symmetry codes: (i)  $x-1/2, -y+1/2, z-1/2$ ; (ii)  $-x+1, -y+1, -z+1$ .

Fig. 1




Fig. 2

