

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

***trans*-Dimethanolbis(1,1,1-trifluoro-5,5-dimethylhexane-2,4-dionato)zinc(II)**

Gerald O. Hunter, Matthias Zeller and Brian D. Leskiw*

Youngstown State University, Department of Chemistry, 1 University Plaza,
Youngstown, OH 44555, USA
Correspondence e-mail: bdleskiw@ysu.edu


Received 13 November 2008; accepted 14 November 2008

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å;
R factor = 0.052; wR factor = 0.114; data-to-parameter ratio = 18.4.

The title compound, $[\text{Zn}(\text{C}_8\text{H}_{10}\text{F}_3\text{O}_2)_2(\text{CH}_4\text{O})_2]$, is a dimethanol coordinated zinc complex with the acetyl acetate derivative 1,1,1-trifluoro-5,5-dimethylhexane-2,4-dionate. The *bis*- β -diketonate complex, which is isostructural with its Co analogue, is located on a crystallographic inversion center. The complex is octahedral with basically no distortion, and the methanol molecules are in *trans* positions with respect to one another. The planes of the β -diketonate and the ZnO_4 unit are tilted by 18.64 (10)° against each other. $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds between the methanol hydroxyl groups and neighboring diketonate O atoms create chains running along $[100]$.

Related literature

For information regarding the synthesis of various metal β -diketonates refer to Watson & Lin (1966). For mass spectrometry related articles see Lerach & Leskiw (2008) and Schilderout (1976). For a variety of applications and properties of metal β -diketonate complexes refer to Burtoloso (2005), Katok *et al.* (2006) and Condorelli *et al.* (2007). Lerach *et al.* (2007) report the structure of the Co analogue of the title compound.



Experimental

Crystal data

$[\text{Zn}(\text{C}_8\text{H}_{10}\text{F}_3\text{O}_2)_2(\text{CH}_4\text{O})_2]$
 $M_r = 519.79$
Triclinic, $P\bar{1}$
 $a = 5.470$ (2) Å
 $b = 8.755$ (3) Å
 $c = 12.031$ (4) Å
 $\alpha = 78.785$ (5)°
 $\beta = 80.542$ (5)°

$\gamma = 88.083$ (5)°
 $V = 557.5$ (3) Å³
 $Z = 1$
Mo $K\alpha$ radiation
 $\mu = 1.18$ mm⁻¹
 $T = 100$ (2) K
 $0.55 \times 0.26 \times 0.05$ mm

Data collection

Bruker SMART APEX CCD
diffractometer
Absorption correction: multi-scan
(*APEX2*; Bruker, 2008)
 $T_{\text{min}} = 0.603$, $T_{\text{max}} = 0.943$

5584 measured reflections
2736 independent reflections
2103 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.044$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$
 $wR(F^2) = 0.114$
 $S = 1.04$
2736 reflections
149 parameters
1 restraint

H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\text{max}} = 0.85$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.97$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O}3-\text{H}3A\cdots\text{O}1^i$	0.82 (2)	2.06 (2)	2.869 (3)	168 (4)

Symmetry code: (i) $x - 1, y, z$.

Data collection: *APEX2* (Bruker, 2008); cell refinement: *APEX2*;
data reduction: *APEX2*; program(s) used to solve structure:
SHELXTL (Sheldrick, 2008); program(s) used to refine structure:
SHELXTL; molecular graphics: *SHELXTL*; software used to
prepare material for publication: *SHELXTL*.

GOH would like to thank Mr Jordan Lerach for his
fundamental contributions in the initial stages of this new
research project and his continued help and assistance. The
diffractometer was funded by NSF grant 0087210, by Ohio
Board of Regents grant CAP-491, and by YSU.

Supplementary data and figures for this paper are available from the
IUCr electronic archives (Reference: LX2078).

References

- Bruker (2008). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
Burtoloso, A. (2005). *Synlett*, **18**, 2859.
Condorelli, G. G., Motta, A., Bedoya, C., Di Mauro, G. P. & Smecca, E. (2007).
Inorg. Chim. Acta, **360**, 170–178.
Katok, K. V., Tertykh, V. A., Brichka, S. Y. & Prikhod, G. P. (2006). *J. Therm.
Anal. Calorim.* **86**, 109–114.
Lerach, O. J. & Leskiw, B. D. (2008). *Rapid Commun. Mass Spectrom.* 4139–
4146.
Lerach, J. O., Zeller, M. & Leskiw, B. D. (2007). *Acta Cryst.* **E63**, m2639.
Schilderout, S. M. (1976). *J. Phys. Chem.* **80**, 2834–2838.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Watson, W. H. & Lin, C. (1966). *Inorg. Chem.* **5**, 1074–1077.

supplementary materials

Acta Cryst. (2009). E65, m24 [doi:10.1107/S1600536808037963]

***trans*-Dimethanolbis(1,1,1-trifluoro-5,5-dimethylhexane-2,4-dionato)zinc(II)**

G. O. Hunter, M. Zeller and B. D. Leskiw

Comment

β -Diketonates and especially metal β -diketonate complexes have been widely studied for both their intrinsic properties as well as a variety of scientific and technological applications. Especially interesting applications include, but are not limited to, catalysis (Burtoloso, 2005), carbon-nanotube structures (Katok *et al.*, 2006), or the deposition of metallic or ceramic thin films (Condorelli *et al.*, 2007). In our own laboratory we are investigating gas phase reactions of a series of metal acetylacetonate (acac) complexes. Through mass spectrometric analysis, several acetylacetonate and substituted acetylacetonate species were observed to undergo various reactions including ligand exchange and association (Schildercrout, 1976; Lerach & Leskiw, 2008). In this context fluorinated metal- β -diketonates are especially interesting because of their increased volatility, thermal stability, and also their ease of preparation.

The title compound, $[\text{Zn}(\text{C}_8\text{H}_{10}\text{F}_3\text{O}_2)_2(\text{CH}_3\text{OH})_2]$, which is isostructural with its Co analogue (Lerach *et al.*, 2007) is a dimethanol coordinate of a zinc complex with the ligand 1,1,1-trifluoro-5,5-dimethylhexane-2,4-dionate, an acetylacetonate derivative with each a *tert*-butyl and a trifluoromethyl substituent. A thermal ellipsoid plot of the molecule is shown in Fig. 1. The bis- β -diketonate complex is located on a crystallographic inversion center with the two methanol molecules in *trans* position to each other. The coordination environment of the central zinc cation is octahedral with only a very slight distortion: angles around the Zn atom deviate from 90° by $0.36(8)^\circ$ or less, and Zn—O distances are 2.054(2) and 2.040(2) Å for the zinc β -diketonate bonds and 2.161(2) Å towards the methanol molecules. The mean planes of the diketonate ligands, defined by the atoms O1, O2 and C1 to C5, and that of the ZnO_4 unit are tilted against each other by an angle of $18.64(10)^\circ$, which is virtually identical to the value of $17.41(7)^\circ$ observed in the structure of the Co analogue of the title compound.

Packing of the molecules within the structure is assisted by hydrogen bonds between the methanol hydroxyl groups and diketonate oxygen atoms of neighboring molecules (Table 1). The O—H \cdots O interactions create hydrogen bonded chains that stretch along the *a*-axis of the structure.

Experimental

The synthesis of the title compound was adapted from Watson & Lin (1966). 0.80 ml (5.0 mmol) of the ligand were added to a stirring solution of 0.22 g of ZnCl_2 (1.6 mmol) and 50 ml of de-ionized water. Diluted 1:1 (v/v) NH_4OH was added dropwise to the mixture until no more visible precipitate formed. The solution was stirred for another hour at room temperature, and the precipitate was isolated by vacuum filtration. The desired product was re-crystallized by overnight evaporation of a concentrated methanolic solution.


Refinement

The hydroxyl H atom was located in a difference density Fourier map and the O—H distance was restrained to $0.84(2)$ Å. All other H atoms were placed in calculated positions with C—H distances of 0.98 (methyl) and 0.95 Å (CH). The methyl and hydroxyl H's were refined with an isotropic displacement parameter U_{iso} of 1.5 times U_{eq} of the adjacent carbon or

supplementary materials

oxygen atom, and the C—H hydrogen atom with $U_{\text{iso}} = 1.2 U_{\text{eq}}(\text{C})$. Methyl hydrogen atoms were allowed to rotate to best fit the experimental electron density.

Figures

 Fig. 1. ORTEP representation of the asymmetric unit of the title compound (50% probability displacement ellipsoids). H atoms are shown as circles of arbitrary radii.

trans-Dimethanolbis(1,1,1-trifluoro-5,5-dimethylhexane-2,4-dionato)zinc(II)

Crystal data

$[\text{Zn}(\text{C}_8\text{H}_{10}\text{F}_3\text{O}_2)_2(\text{CH}_4\text{O})_2]$	$Z = 1$
$M_r = 519.79$	$F_{000} = 268$
Triclinic, $P\bar{1}$	$D_x = 1.548 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation
$a = 5.470$ (2) Å	$\lambda = 0.71073$ Å
$b = 8.755$ (3) Å	Cell parameters from 1117 reflections
$c = 12.031$ (4) Å	$\theta = 2.4\text{--}29.6^\circ$
$\alpha = 78.785$ (5)°	$\mu = 1.18 \text{ mm}^{-1}$
$\beta = 80.542$ (5)°	$T = 100$ (2) K
$\gamma = 88.083$ (5)°	Plate, colourless
$V = 557.5$ (3) Å ³	$0.55 \times 0.26 \times 0.05 \text{ mm}$

Data collection

Bruker SMART APEX CCD diffractometer	2736 independent reflections
Radiation source: fine-focus sealed tube	2103 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.044$
Detector resolution: 10.0 pixels mm^{-1}	$\theta_{\text{max}} = 28.3^\circ$
$T = 100$ (2) K	$\theta_{\text{min}} = 1.8^\circ$
ω scans	$h = -7 \rightarrow 7$
Absorption correction: multi-scan (APEX2; Bruker, 2008)	$k = -11 \rightarrow 11$
$T_{\text{min}} = 0.603$, $T_{\text{max}} = 0.943$	$l = -15 \rightarrow 16$
5584 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.052$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.114$	$w = 1/[\sigma^2(F_o^2) + (0.0494P)^2 + 0.1141P]$
	where $P = (F_o^2 + 2F_c^2)/3$

$S = 1.04$ $(\Delta/\sigma)_{\max} < 0.001$
 2736 reflections $\Delta\rho_{\max} = 0.85 \text{ e } \text{\AA}^{-3}$
 149 parameters $\Delta\rho_{\min} = -0.97 \text{ e } \text{\AA}^{-3}$
 1 restraint Extinction correction: none
 Primary atom site location: structure-invariant direct methods

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	1.1698 (6)	1.2211 (4)	0.7642 (3)	0.0245 (7)
C2	1.0749 (5)	1.0950 (3)	0.7105 (3)	0.0196 (6)
C3	0.9063 (5)	0.9904 (3)	0.7791 (3)	0.0211 (6)
H3	0.8553	1.0023	0.8563	0.025*
C4	0.8021 (5)	0.8648 (3)	0.7425 (3)	0.0199 (6)
C5	0.6476 (5)	0.7405 (3)	0.8316 (3)	0.0200 (6)
C6	0.4765 (6)	0.8131 (4)	0.9221 (3)	0.0266 (7)
H6A	0.3670	0.8895	0.8840	0.040*
H6B	0.5768	0.8647	0.9643	0.040*
H6C	0.3764	0.7312	0.9758	0.040*
C7	0.8326 (6)	0.6306 (4)	0.8926 (3)	0.0265 (7)
H7A	0.7410	0.5512	0.9521	0.040*
H7B	0.9354	0.6910	0.9278	0.040*
H7C	0.9389	0.5801	0.8364	0.040*
C8	0.4976 (6)	0.6471 (4)	0.7721 (3)	0.0298 (8)
H8A	0.3999	0.5681	0.8297	0.045*
H8B	0.6103	0.5961	0.7184	0.045*
H8C	0.3861	0.7171	0.7300	0.045*
C9	0.7190 (6)	1.3221 (4)	0.5071 (3)	0.0290 (7)
H9A	0.7125	1.3480	0.5832	0.044*
H9B	0.5854	1.3768	0.4702	0.044*
H9C	0.8793	1.3542	0.4599	0.044*
F1	1.4150 (3)	1.2087 (2)	0.76434 (17)	0.0350 (5)
F2	1.0662 (4)	1.2202 (2)	0.87310 (16)	0.0385 (5)
F3	1.1312 (4)	1.3633 (2)	0.70461 (18)	0.0384 (5)
O1	1.1718 (4)	1.1047 (2)	0.60505 (17)	0.0214 (5)

supplementary materials

O2	0.8347 (4)	0.8468 (2)	0.64018 (18)	0.0223 (5)
O3	0.6892 (4)	1.1570 (2)	0.51870 (19)	0.0252 (5)
H3A	0.547 (4)	1.129 (4)	0.548 (3)	0.038*
Zn1	1.0000	1.0000	0.5000	0.01973 (16)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0202 (15)	0.0285 (17)	0.0260 (17)	-0.0026 (13)	-0.0032 (13)	-0.0083 (13)
C2	0.0129 (13)	0.0233 (15)	0.0249 (16)	0.0025 (11)	-0.0068 (11)	-0.0075 (12)
C3	0.0172 (14)	0.0256 (16)	0.0212 (15)	0.0002 (12)	-0.0031 (12)	-0.0067 (12)
C4	0.0109 (13)	0.0223 (15)	0.0257 (16)	0.0043 (11)	-0.0019 (11)	-0.0040 (12)
C5	0.0163 (14)	0.0205 (15)	0.0227 (16)	-0.0004 (11)	-0.0018 (12)	-0.0039 (12)
C6	0.0196 (15)	0.0272 (16)	0.0301 (17)	0.0004 (13)	0.0005 (13)	-0.0024 (14)
C7	0.0196 (15)	0.0253 (16)	0.0323 (18)	-0.0009 (12)	-0.0015 (13)	-0.0020 (14)
C8	0.0251 (16)	0.0338 (18)	0.0301 (18)	-0.0146 (14)	-0.0014 (14)	-0.0048 (14)
C9	0.0271 (17)	0.0235 (16)	0.0370 (19)	0.0037 (13)	-0.0037 (14)	-0.0088 (14)
F1	0.0192 (9)	0.0440 (12)	0.0478 (13)	-0.0032 (8)	-0.0100 (9)	-0.0191 (10)
F2	0.0398 (12)	0.0472 (13)	0.0315 (11)	-0.0171 (10)	0.0070 (9)	-0.0227 (10)
F3	0.0513 (13)	0.0216 (10)	0.0473 (13)	0.0004 (9)	-0.0196 (10)	-0.0093 (9)
O1	0.0172 (10)	0.0255 (11)	0.0222 (11)	-0.0015 (8)	-0.0026 (8)	-0.0067 (9)
O2	0.0211 (10)	0.0221 (11)	0.0241 (11)	-0.0026 (8)	-0.0028 (9)	-0.0059 (9)
O3	0.0148 (10)	0.0226 (11)	0.0374 (13)	0.0007 (9)	0.0000 (9)	-0.0076 (10)
Zn1	0.0150 (3)	0.0215 (3)	0.0228 (3)	-0.00167 (19)	-0.00203 (19)	-0.0050 (2)

Geometric parameters (\AA , $^\circ$)

C1—F2	1.338 (4)	C7—H7B	0.9800
C1—F3	1.339 (4)	C7—H7C	0.9800
C1—F1	1.342 (3)	C8—H8A	0.9800
C1—C2	1.526 (4)	C8—H8B	0.9800
C2—O1	1.280 (3)	C8—H8C	0.9800
C2—C3	1.372 (4)	C9—O3	1.438 (4)
C3—C4	1.428 (4)	C9—H9A	0.9800
C3—H3	0.9500	C9—H9B	0.9800
C4—O2	1.254 (4)	C9—H9C	0.9800
C4—C5	1.536 (4)	O1—Zn1	2.054 (2)
C5—C8	1.523 (4)	O2—Zn1	2.040 (2)
C5—C6	1.536 (4)	O3—Zn1	2.161 (2)
C5—C7	1.547 (4)	O3—H3A	0.824 (18)
C6—H6A	0.9800	Zn1—O2 ⁱ	2.040 (2)
C6—H6B	0.9800	Zn1—O1 ⁱ	2.054 (2)
C6—H6C	0.9800	Zn1—O3 ⁱ	2.161 (2)
C7—H7A	0.9800		
F2—C1—F3	106.6 (3)	C5—C8—H8A	109.5
F2—C1—F1	106.0 (3)	C5—C8—H8B	109.5
F3—C1—F1	106.3 (2)	H8A—C8—H8B	109.5
F2—C1—C2	114.7 (2)	C5—C8—H8C	109.5

F3—C1—C2	111.1 (3)	H8A—C8—H8C	109.5
F1—C1—C2	111.6 (2)	H8B—C8—H8C	109.5
O1—C2—C3	130.0 (3)	O3—C9—H9A	109.5
O1—C2—C1	112.4 (2)	O3—C9—H9B	109.5
C3—C2—C1	117.7 (3)	H9A—C9—H9B	109.5
C2—C3—C4	124.6 (3)	O3—C9—H9C	109.5
C2—C3—H3	117.7	H9A—C9—H9C	109.5
C4—C3—H3	117.7	H9B—C9—H9C	109.5
O2—C4—C3	123.5 (3)	C2—O1—Zn1	119.7 (2)
O2—C4—C5	116.9 (3)	C4—O2—Zn1	126.7 (2)
C3—C4—C5	119.6 (3)	C9—O3—Zn1	122.6 (2)
C8—C5—C4	110.1 (2)	C9—O3—H3A	112 (3)
C8—C5—C6	110.5 (2)	Zn1—O3—H3A	124 (3)
C4—C5—C6	111.6 (2)	O2 ⁱ —Zn1—O2	180.0
C8—C5—C7	109.1 (3)	O2 ⁱ —Zn1—O1 ⁱ	89.64 (8)
C4—C5—C7	106.9 (2)	O2—Zn1—O1 ⁱ	90.36 (8)
C6—C5—C7	108.5 (3)	O2 ⁱ —Zn1—O1	90.36 (8)
C5—C6—H6A	109.5	O2—Zn1—O1	89.64 (8)
C5—C6—H6B	109.5	O1 ⁱ —Zn1—O1	180.00 (11)
H6A—C6—H6B	109.5	O2 ⁱ —Zn1—O3	89.95 (8)
C5—C6—H6C	109.5	O2—Zn1—O3	90.05 (8)
H6A—C6—H6C	109.5	O1 ⁱ —Zn1—O3	89.97 (8)
H6B—C6—H6C	109.5	O1—Zn1—O3	90.03 (8)
C5—C7—H7A	109.5	O2 ⁱ —Zn1—O3 ⁱ	90.05 (8)
C5—C7—H7B	109.5	O2—Zn1—O3 ⁱ	89.95 (8)
H7A—C7—H7B	109.5	O1 ⁱ —Zn1—O3 ⁱ	90.03 (8)
C5—C7—H7C	109.5	O1—Zn1—O3 ⁱ	89.97 (9)
H7A—C7—H7C	109.5	O3—Zn1—O3 ⁱ	179.999 (1)
H7B—C7—H7C	109.5		
F2—C1—C2—O1	177.3 (2)	C3—C2—O1—Zn1	20.4 (4)
F3—C1—C2—O1	56.2 (3)	C1—C2—O1—Zn1	-160.1 (2)
F1—C1—C2—O1	-62.2 (3)	C3—C4—O2—Zn1	-8.4 (4)
F2—C1—C2—C3	-3.2 (4)	C5—C4—O2—Zn1	173.6 (2)
F3—C1—C2—C3	-124.2 (3)	C4—O2—Zn1—O1 ⁱ	-159.3 (2)
F1—C1—C2—C3	117.3 (3)	C4—O2—Zn1—O1	20.7 (2)
O1—C2—C3—C4	0.2 (5)	C4—O2—Zn1—O3	-69.4 (2)
C1—C2—C3—C4	-179.2 (3)	C4—O2—Zn1—O3 ⁱ	110.6 (2)
C2—C3—C4—O2	-7.7 (5)	C2—O1—Zn1—O2 ⁱ	155.1 (2)
C2—C3—C4—C5	170.3 (3)	C2—O1—Zn1—O2	-24.9 (2)
O2—C4—C5—C8	-17.8 (4)	C2—O1—Zn1—O3	65.1 (2)
C3—C4—C5—C8	164.0 (3)	C2—O1—Zn1—O3 ⁱ	-114.9 (2)
O2—C4—C5—C6	-141.0 (3)	C9—O3—Zn1—O2 ⁱ	-43.4 (2)
C3—C4—C5—C6	40.8 (4)	C9—O3—Zn1—O2	136.6 (2)
O2—C4—C5—C7	100.5 (3)	C9—O3—Zn1—O1 ⁱ	-133.0 (2)
C3—C4—C5—C7	-77.6 (3)	C9—O3—Zn1—O1	47.0 (2)

supplementary materials

Symmetry codes: (i) $-x+2, -y+2, -z+1$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$O3-H3A\cdots O1^{ii}$	0.82 (2)	2.06 (2)	2.869 (3)	168 (4)

Symmetry codes: (ii) $x-1, y, z$.

Fig. 1

