

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-[(*Z*)-(1-Methyl-1*H*-pyrrol-2-yl)methylidene]-1*H*-1,2,4-triazol-5-amine

Zahid H. Chohan,^a Muhammad Hanif^a and M. Nawaz Tahir^{b*}

^aDepartment of Chemistry, Bahauddin Zakariya University, Multan 60800, Pakistan,and ^bDepartment of Physics, University of Sargodha, Sargodha, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 3 December 2008; accepted 4 December 2008

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.037; wR factor = 0.107; data-to-parameter ratio = 16.3.

In the title compound, $\text{C}_8\text{H}_9\text{N}_5$, a Schiff base derived from *N*-methylpyrrole-2-carbaldehyde and 3-amino-1,2,4-triazole, the $\text{C}=\text{N}$ double bond linking the two aromatic rings has a *Z* conformation. The two rings are twisted by $24.20(5)^\circ$. A chain motif results from $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonding.

Related literature

For a related structure, see: Arfan *et al.* (2008).



Experimental

Crystal data

 $\text{C}_8\text{H}_9\text{N}_5$ $M_r = 175.20$ Monoclinic, $P2_1/c$ $a = 7.2519(3)$ Å $b = 12.8616(6)$ Å $c = 9.7445(4)$ Å $\beta = 101.917(2)^\circ$ $V = 889.29(7)$ Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.09$ mm⁻¹ $T = 296(2)$ K $0.26 \times 0.20 \times 0.16$ mm

Data collection

Bruker Kappa APEXII CCD diffractometer

Absorption correction: multi-scan (*SADABS*; Bruker, 2005) $T_{\min} = 0.976$, $T_{\max} = 0.988$

10203 measured reflections

2212 independent reflections

1651 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.024$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.037$ $wR(F^2) = 0.107$ $S = 1.03$

2212 reflections

136 parameters

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.16$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.17$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N3}-\text{H3n}\cdots\text{N5}^i$	0.91 (1)	1.92 (1)	2.8225 (12)	171 (1)

Symmetry code: (i) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2003); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

The authors acknowledge the the Higher Education Commission, Islamabad, Pakistan, for funding the purchase of the diffractometer at GCU, Lahore.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2523).

References

- Arfan, M., Tahir, M. N., Khan, R. & Iqbal, M. S. (2008). *Acta Cryst.* **E64**, o1505.
 Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2009). E65, o58 [doi:10.1107/S160053680804097X]

N-[(*Z*)-(1-Methyl-1*H*-pyrrol-2-yl)methylidene]-1*H*-1,2,4-triazol-5-amine

Z. H. Chohan, M. Hanif and M. N. Tahir

Comment

1,2,4-triazole ring is a basic aromatic ring and it possess various medicinal properties. The title compound (I), has been prepared to utilize it as an intermediate ligand and for complexation with various metals.

In the crystal structure of (I), (Fig 1), the pyrrole ring A(N1,C2—C5) is connected to the 1,2,4 triazole ring B(C7/N3/N4/C8/N5) through the Schiff bond C=N. There exist an intramolecular and an intermolecular H-bond (Fig 2), Table 1. The bond distances and bond angles of ring B are compareable as observed in the same moiety of 3-(2-Benzamidophenyl)-4-(4-hydroxyphenyl)- 5-methyl-4*H*-1,2,4-triazol-1-ium chloride (Arfan *et al.*, 2008). Due to intermolecular H-bonding, the compound forms polymeric sheets. The dihedral angle between the rings A and B is 24.20 (5)°. The molecules are stabilized due to π - π interactions between the centroids CgA and CgB of rings A and B respectively. The centroid to centroid, CgA...CgBⁱ [Symmetry code: $i = 1 - x, 1 - y, 1 - z$] and CgB...CgAⁱⁱ [Symmetry code: $ii = -x, 1 - y, 1 - z$] is 3.9008 (8) and 3.9009 (8) Å, respectively.

Experimental

N-methyl pyrrole-2-carboxyaldehyde (1.047 ml, 0.01 *M*) in methanol solution (10 ml) was added to magnetically stirred methanol solution (20 ml) of 3-amino 1,2,4 triazole (0.84 g m, 0.01 *M*) and mixture refluxed for 5 h through monitoring by TLC. After completion of the reaction, the resultant mixture was cooled to room temperature, filtered and reduced nearly half of its volume by rotary. It was then allowed to stay at room temperature for 2 days which resulted in the formation of a colorless solid product. It was filtered, washed with methanol and recrystallized with a mixture of ethanol:methanol (1:1).

Refinement

H-atoms were positioned geometrically, with C—H = 0.96 Å for methyl carbon and constrained to ride on their parent atom. The coordinates of all other H-atoms were refined. The $U_{iso}(H) = xU_{eq}(C, N)$, where $x = 1.5$ for methyl H and $x = 1.2$ for all other H atoms.

Figures




Fig. 1. ORTEP-3 for Windows (Farrugia, 1997) drawing of the title compound, C₈H₉N₅, with the atom numbering scheme. The thermal ellipsoids are drawn at the 50% probability level. H-atoms are shown by small circles of arbitrary radii. The intramolecular H-bonding is shown by dashed lines.




Fig. 2. The partial unit cell packing of (I) (Spek, 2003) showing the intermolecular and intermolecular hydrogen bonding showing that polymeric sheets are formed.

***N*-[(*Z*)-(1-Methyl-1*H*-pyrrol-2-yl)methylidene]-1*H*-1,2,4-triazol-5-amine**

Crystal data

$C_8H_9N_5$	$F_{000} = 368$
$M_r = 175.20$	$D_x = 1.309 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 7.2519 (3) \text{ \AA}$	Cell parameters from 1859 reflections
$b = 12.8616 (6) \text{ \AA}$	$\theta = 2.7\text{--}28.3^\circ$
$c = 9.7445 (4) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$\beta = 101.917 (2)^\circ$	$T = 296 (2) \text{ K}$
$V = 889.29 (7) \text{ \AA}^3$	Prismatic, orange
$Z = 4$	$0.26 \times 0.20 \times 0.16 \text{ mm}$

Data collection

Bruker Kappa APEXII CCD diffractometer	2212 independent reflections
Radiation source: fine-focus sealed tube	1651 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.024$
Detector resolution: $7.4 \text{ pixels mm}^{-1}$	$\theta_{\text{max}} = 28.3^\circ$
$T = 296(2) \text{ K}$	$\theta_{\text{min}} = 2.7^\circ$
ω scans	$h = -5 \rightarrow 9$
Absorption correction: multi-scan (SADABS; Bruker, 2005)	$k = -16 \rightarrow 17$
$T_{\text{min}} = 0.976$, $T_{\text{max}} = 0.988$	$l = -12 \rightarrow 12$
10203 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.037$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.107$	$w = 1/[\sigma^2(F_o^2) + (0.0556P)^2 + 0.079P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
	$(\Delta/\sigma)_{\text{max}} = 0.001$

2212 reflections $\Delta\rho_{\max} = 0.16 \text{ e } \text{\AA}^{-3}$
 136 parameters $\Delta\rho_{\min} = -0.17 \text{ e } \text{\AA}^{-3}$
 Primary atom site location: structure-invariant direct methods Extinction correction: none

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R - factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.11672 (13)	0.58683 (8)	0.33358 (11)	0.0543 (3)
N2	0.27287 (13)	0.37308 (7)	0.38528 (9)	0.0459 (3)
N3	0.38353 (14)	0.21080 (7)	0.32861 (9)	0.0475 (3)
N4	0.44733 (16)	0.11865 (8)	0.38789 (10)	0.0552 (3)
N5	0.37569 (15)	0.22874 (7)	0.54902 (9)	0.0494 (3)
C1	0.0572 (2)	0.53330 (12)	0.20067 (14)	0.0754 (5)
C2	0.0834 (2)	0.68853 (11)	0.35562 (19)	0.0691 (5)
C3	0.1530 (2)	0.71296 (12)	0.49152 (19)	0.0719 (6)
C4	0.23118 (19)	0.62366 (11)	0.55771 (16)	0.0609 (5)
C5	0.20967 (15)	0.54493 (9)	0.45902 (12)	0.0471 (3)
C6	0.27772 (15)	0.44120 (9)	0.48170 (12)	0.0451 (3)
C7	0.34307 (15)	0.27546 (8)	0.42465 (10)	0.0415 (3)
C8	0.43865 (19)	0.13408 (10)	0.51969 (12)	0.0545 (4)
H1A	-0.03220	0.57556	0.13837	0.1132*
H1B	-0.00032	0.46826	0.21599	0.1132*
H1C	0.16470	0.52068	0.15976	0.1132*
H2	0.025 (2)	0.7315 (13)	0.2805 (17)	0.0829*
H3	0.149 (2)	0.7801 (13)	0.5321 (17)	0.0862*
H3N	0.3769 (18)	0.2235 (10)	0.2359 (14)	0.0571*
H4	0.294 (2)	0.6150 (11)	0.6550 (17)	0.0730*
H6	0.3326 (17)	0.4259 (9)	0.5798 (14)	0.0541*
H8	0.4755 (19)	0.0815 (11)	0.5904 (15)	0.0655*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0532 (5)	0.0502 (6)	0.0587 (6)	0.0034 (4)	0.0099 (4)	0.0071 (5)
N2	0.0558 (5)	0.0467 (5)	0.0355 (5)	0.0026 (4)	0.0102 (4)	0.0039 (4)
N3	0.0696 (6)	0.0446 (5)	0.0290 (4)	-0.0016 (4)	0.0116 (4)	-0.0003 (4)

supplementary materials

N4	0.0813 (7)	0.0435 (6)	0.0415 (5)	0.0025 (5)	0.0143 (5)	-0.0004 (4)
N5	0.0708 (6)	0.0486 (5)	0.0302 (4)	0.0041 (4)	0.0137 (4)	0.0046 (4)
C1	0.0920 (10)	0.0713 (10)	0.0546 (8)	0.0084 (8)	-0.0044 (7)	0.0100 (7)
C2	0.0635 (8)	0.0525 (8)	0.0924 (11)	0.0090 (6)	0.0187 (7)	0.0113 (7)
C3	0.0720 (9)	0.0503 (8)	0.0983 (12)	0.0023 (7)	0.0292 (8)	-0.0106 (8)
C4	0.0618 (8)	0.0581 (8)	0.0653 (8)	-0.0038 (6)	0.0189 (6)	-0.0101 (6)
C5	0.0455 (5)	0.0481 (6)	0.0494 (6)	-0.0023 (5)	0.0135 (4)	0.0009 (5)
C6	0.0499 (6)	0.0485 (6)	0.0373 (5)	-0.0020 (5)	0.0102 (4)	0.0035 (5)
C7	0.0515 (6)	0.0444 (6)	0.0288 (5)	-0.0021 (4)	0.0090 (4)	0.0009 (4)
C8	0.0782 (8)	0.0466 (7)	0.0388 (6)	0.0043 (6)	0.0120 (5)	0.0068 (5)

Geometric parameters (Å, °)

N1—C1	1.4515 (17)	C3—C4	1.380 (2)
N1—C2	1.3553 (18)	C4—C5	1.3830 (19)
N1—C5	1.3779 (15)	C5—C6	1.4238 (16)
N2—C6	1.2798 (14)	C1—H1A	0.9600
N2—C7	1.3792 (14)	C1—H1B	0.9600
N3—N4	1.3572 (14)	C1—H1C	0.9600
N3—C7	1.3293 (14)	C2—H2	0.945 (16)
N4—C8	1.3139 (15)	C3—H3	0.953 (17)
N5—C7	1.3294 (13)	C4—H4	0.970 (16)
N5—C8	1.3514 (16)	C6—H6	0.977 (13)
N3—H3N	0.910 (13)	C8—H8	0.963 (14)
C2—C3	1.353 (3)		
N1…N2	2.9763 (14)	C1…H3 ^{vii}	3.058 (16)
N2…N1	2.9763 (14)	C6…H1B	2.9600
N2…C1	2.9609 (17)	C7…H2 ^{iv}	3.039 (16)
N3…N5 ⁱ	2.8225 (12)	C7…H3N ⁱⁱⁱ	2.992 (13)
N3…N5	2.1725 (12)	C8…H8 ⁱⁱ	3.083 (14)
N4…C8 ⁱⁱ	3.4283 (17)	C8…H3N ⁱⁱⁱ	2.896 (13)
N4…N5	2.2532 (14)	H1A…H2	2.4200
N5…N3 ⁱⁱⁱ	2.8225 (12)	H1B…N2	2.6100
N5…N4	2.2532 (14)	H1B…C6	2.9600
N2…H1B	2.6100	H1C…N2	2.8900
N2…H1C	2.8900	H2…H1A	2.4200
N3…H2 ^{iv}	2.946 (15)	H2…N3 ^{viii}	2.946 (15)
N4…H8 ⁱⁱ	2.634 (14)	H2…C7 ^{viii}	3.039 (16)
N5…H6	2.580 (12)	H3…C1 ^{ix}	3.058 (16)
N5…H3N ⁱⁱⁱ	1.920 (13)	H3N…N5 ⁱ	1.920 (13)
C1…N2	2.9609 (17)	H3N…C7 ⁱ	2.992 (13)
C3…C7 ^v	3.5793 (19)	H3N…C8 ⁱ	2.896 (13)
C3…C8 ^v	3.576 (2)	H3N…H6 ⁱ	2.431 (18)
C4…C7 ^v	3.3206 (18)	H4…H6	2.572 (19)
C5…C5 ^{vi}	3.4963 (16)	H6…N5	2.580 (12)
C6…C6 ^v	3.5119 (16)	H6…H4	2.572 (19)

C7...C3 ^v	3.5793 (19)	H6...H3N ⁱⁱⁱ	2.431 (18)
C7...C4 ^v	3.3206 (18)	H8...N4 ⁱⁱ	2.634 (14)
C8...C3 ^v	3.576 (2)	H8...C8 ⁱⁱ	3.083 (14)
C8...N4 ⁱⁱ	3.4283 (17)		
C1—N1—C2	124.70 (12)	N3—C7—N5	109.60 (9)
C1—N1—C5	127.37 (11)	N4—C8—N5	115.42 (11)
C2—N1—C5	107.91 (11)	N1—C1—H1A	109.00
C6—N2—C7	117.80 (9)	N1—C1—H1B	109.00
N4—N3—C7	110.62 (8)	N1—C1—H1C	109.00
N3—N4—C8	101.73 (10)	H1A—C1—H1B	109.00
C7—N5—C8	102.63 (9)	H1A—C1—H1C	109.00
N4—N3—H3N	121.5 (8)	H1B—C1—H1C	109.00
C7—N3—H3N	127.8 (8)	N1—C2—H2	120.5 (10)
N1—C2—C3	109.66 (14)	C3—C2—H2	129.7 (10)
C2—C3—C4	107.35 (14)	C2—C3—H3	125.4 (10)
C3—C4—C5	107.93 (13)	C4—C3—H3	127.2 (10)
N1—C5—C4	107.15 (11)	C3—C4—H4	128.1 (9)
C4—C5—C6	126.33 (11)	C5—C4—H4	124.0 (9)
N1—C5—C6	126.48 (10)	N2—C6—H6	121.6 (7)
N2—C6—C5	124.85 (11)	C5—C6—H6	113.5 (7)
N2—C7—N3	119.75 (9)	N4—C8—H8	122.1 (9)
N2—C7—N5	130.62 (9)	N5—C8—H8	122.5 (9)
C1—N1—C5—C4	-177.86 (12)	C7—N3—N4—C8	0.72 (13)
C1—N1—C2—C3	178.37 (12)	N3—N4—C8—N5	-0.41 (15)
C5—N1—C2—C3	0.21 (16)	C8—N5—C7—N3	0.52 (13)
C2—N1—C5—C6	-177.44 (12)	C8—N5—C7—N2	178.36 (12)
C2—N1—C5—C4	0.23 (14)	C7—N5—C8—N4	-0.06 (15)
C1—N1—C5—C6	4.47 (19)	N1—C2—C3—C4	-0.57 (17)
C6—N2—C7—N5	18.65 (18)	C2—C3—C4—C5	0.70 (17)
C6—N2—C7—N3	-163.70 (11)	C3—C4—C5—C6	177.10 (12)
C7—N2—C6—C5	-179.40 (11)	C3—C4—C5—N1	-0.57 (15)
N4—N3—C7—N5	-0.82 (14)	C4—C5—C6—N2	-172.36 (12)
N4—N3—C7—N2	-178.93 (10)	N1—C5—C6—N2	4.88 (19)

Symmetry codes: (i) $x, -y+1/2, z-1/2$; (ii) $-x+1, -y, -z+1$; (iii) $x, -y+1/2, z+1/2$; (iv) $-x, y-1/2, -z+1/2$; (v) $-x+1, -y+1, -z+1$; (vi) $-x, -y+1, -z+1$; (vii) $x, -y+3/2, z-1/2$; (viii) $-x, y+1/2, -z+1/2$; (ix) $x, -y+3/2, z+1/2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N3—H3n...N5 ⁱ	0.910 (13)	1.920 (13)	2.8225 (12)	171.3 (12)
C1—H1B...N2	0.9600	2.6100	2.9609 (17)	102.00

Symmetry codes: (i) $x, -y+1/2, z-1/2$.

Fig. 1




Fig. 2

