

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bis(1-methyl-1*H*-imidazole- κ N³)[*N,N'*-*o*-phenylenebis(pyridine-2-carboxamido)- κ^4 N]manganese(II)

 Zaki N. Zahran,^a Nan Xu,^b Douglas R. Powell^b and George B. Richter-Addo^{b*}
^aChemistry Department, Faculty of Science, Tanta University, Tanta, Egypt, and

^bDepartment of Chemistry and Biochemistry, University of Oklahoma, 620

Parrington Oval, Norman, OK 73019 USA

Correspondence e-mail: grichteraddo@ou.edu


Received 16 October 2008; accepted 8 December 2008

 Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.025; wR factor = 0.062; data-to-parameter ratio = 10.4.

The title compound, $[\text{Mn}(\text{C}_{18}\text{H}_{12}\text{N}_4\text{O}_2)(\text{C}_4\text{H}_6\text{N}_2)_2]$, belongs to the family of 1,2-bis(pyridine-2-carboxamido)benzene (H_2bpb) ligated metal complexes. The manganese center is octahedrally coordinated by a bpb ligand and two axial 1-methylimidazole molecules. The axial $\text{N}-\text{Mn}-\text{N}$ group is bent with a bond angle of $151.79(7)^\circ$.

Related literature

For the structures of related Mn complexes, see Liang *et al.* (2007), Lin *et al.* (2003), and Havranek *et al.* (1999).



Experimental

Crystal data

 $[\text{Mn}(\text{C}_{18}\text{H}_{12}\text{N}_4\text{O}_2)(\text{C}_4\text{H}_6\text{N}_2)_2]$
 $M_r = 535.47$

 Orthorhombic, $Pca2_1$
 $a = 13.819(3)$ Å

 $b = 9.894(2)$ Å

 $c = 17.864(4)$ Å

 $V = 2442.5(9)$ Å³
 $Z = 4$

 Mo $K\alpha$ radiation

 $\mu = 0.58$ mm⁻¹
 $T = 100(2)$ K

 $0.54 \times 0.35 \times 0.04$ mm

Data collection

Bruker APEX diffractometer

Absorption correction: multi-scan

 (*SADABS*; Sheldrick, 2007)

 $T_{\min} = 0.744$, $T_{\max} = 0.977$

8288 measured reflections

3473 independent reflections

 3301 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.025$
 $wR(F^2) = 0.062$
 $S = 1.02$

3473 reflections

335 parameters

1 restraint

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.33$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.19$ e Å⁻³

Absolute structure: Flack (1983),

989 Friedel pairs

Flack parameter: 0.046 (17)

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1998); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

This work was supported by the National Science Foundation (CHE-0079282 and CHE-0076640). The authors thank the National Science Foundation (CHE-0130835) and the University of Oklahoma for funds to acquire the diffractometer and computers used in this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2130).

References

- Bruker (1998). *SMART* and *SAINT*. Bruker AXS, Inc., Madison, Wisconsin, USA.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Havranek, M., Singh, A. & Sames, D. (1999). *J. Am. Chem. Soc.* **121**, 8965–8966.
- Liang, L., Qu, Y.-Y., Yang, L. & Zhou, X.-G. (2007). *Acta Cryst.* **E63**, m1503–m1505.
- Lin, J., Tu, C., Lin, H., Jiang, P., Ding, J. & Guo, Z. (2003). *Inorg. Chem. Commun.* **6**, 262–265.
- Sheldrick, G. M. (2007). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, m75 [doi:10.1107/S1600536808041640]

Bis(1-methyl-1*H*-imidazole- κN^3)[*N,N'*-*o*-phenylenebis(pyridine-2-carboxamido)- $\kappa^4 N$]manganese(II)

Z. N. Zahran, N. Xu, D. R. Powell and G. B. Richter-Addo

Comment

In this paper, we report the structure of the title compound, a six-coordinate {bis(1-methylimidazole)}(bpb)manganese(II) ($H_2bpb = 1,2$ -bis(pyridine-2-carboxamido)benzene). To the best of our knowledge, this is the first reported structure of a Mn(II) complex containing ligated bpb or its derivatives. The structures of the related [Mn(bpb)(H₂O)Cl] (Lin *et al.* 2003), [Mn(bpc)(DMF)Cl] ($H_2bpc = 1,2$ -bis(pyridine-2-carboxamido)-4,5-dichlorobenzene) (Liang *et al.*, 2007) and [Mn(bpmb)(OMe)(OCOCH₃)] ($H_2bpmb = N,N'$ -bis(pyridine-2-ylcarbonyl)-4-methoxycarbonylbenzene-1,2-diamine) (Havranek *et al.*, 1999) complexes have been reported previously.

The molecular structure is shown in Fig. 1. The manganese center is six-coordinate, displaying a distorted octahedral geometry. A bpb ligand binds to the manganese through its two deprotonated amide N atoms and two pyridyl N atoms. The two axial positions are occupied by 1-methylimidazole molecules. The Mn—N(pyridyl) distances of 2.2621 (19) Å and 2.2684 (19) Å are longer than the Mn—N(amide) distances at 2.1764 (19) Å and 2.1794 (18) Å. Both of the Mn—N(pyridyl) and Mn—N(amide) distances are significantly longer than those in the related Mn complexes reported previously (Liang *et al.*, 2007; Lin *et al.*, 2003; Havranek *et al.*, 1999). In addition, the C13—C14 and C21—C22 distances of 1.513 (3) Å and 1.526 (3) Å are slightly longer than those of the other Mn complexes mentioned above. Mn—N(1-methylimidazole) distances are 2.2552 (19) Å and 2.280 (2) Å. The axial N—Mn—N linkage is bent with a bond angle of 151.79 (7)°.

Experimental

To a CH₂Cl₂ suspension (20 ml) of Mn(bpb)Cl (0.2 g, 0.49 mmol) was added excess piperidine (2 ml, 0.02 mol) (Aldrich Chemical Company, used as received) and then purged with nitric oxide (98%; Matheson Gas, purified by passing through KOH pellets and a cold trap (dry ice/acetone)) for 30 min. This resulted in the precipitation of a red-brown intermediate Mn(bpb)(NO)(pip) ($\nu_{NO} 1732\text{ cm}^{-1}$; KBr pellet) that was isolated by filtration. This intermediate was redissolved in CH₂Cl₂, and excess 1-methylimidazole (0.2 ml, 2.6 mmol) (Aldrich Chemical Company, used as received) was added. The resulting mixture was stirred for 30 min. A brown solid was obtained after removal of the solvent under vacuum. A suitable red plate-shaped crystal was grown by slow evaporation of a CH₂Cl₂ solution of the complex in the presence of excess 1-methylimidazole at room temperature under N₂.

Refinement

H atoms were positioned geometrically and refined using a riding model with C—H = 0.95 Å for aromatic carbons, 0.98 Å for methyl carbons. $U_{iso}(H)$ values were set to either 1.5 U_{eq} (RCH₃) or 1.2 U_{eq} of the attached atom.

Figures




Fig. 1. The molecular structure showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 35% probability level (H atoms omitted for clarity).

Bis(1-methyl-1H-imidazole- κ N³)[N,N'-o-phenylenebis(pyridine-2-carboxamidato)- κ ⁴N]manganese(II)

Crystal data

[Mn(C₁₈H₁₂N₄O₂)(C₄H₆N₂)₂]

$M_r = 535.47$

Orthorhombic, $Pca2_1$

Hall symbol: P 2c -2ac

$a = 13.819$ (3) Å

$b = 9.894$ (2) Å

$c = 17.864$ (4) Å

$V = 2442.5$ (9) Å³

$Z = 4$

$F_{000} = 1108$

$D_x = 1.456$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 6910 reflections

$\theta = 2.5$ – 28.3°

$\mu = 0.58$ mm⁻¹

$T = 100$ (2) K

Plate, red

$0.54 \times 0.35 \times 0.04$ mm

Data collection

Bruker APEX
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 100$ (2) K

ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 2007)

$T_{\min} = 0.744$, $T_{\max} = 0.977$

8288 measured reflections

3473 independent reflections

3301 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.022$

$\theta_{\max} = 26.0^\circ$

$\theta_{\min} = 2.1^\circ$

$h = -16 \rightarrow 17$

$k = -12 \rightarrow 7$

$l = -22 \rightarrow 13$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.025$

$wR(F^2) = 0.062$

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.038P)^2 + 0.1P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$S = 1.02$ $\Delta\rho_{\max} = 0.33 \text{ e } \text{\AA}^{-3}$
 3473 reflections $\Delta\rho_{\min} = -0.19 \text{ e } \text{\AA}^{-3}$
 335 parameters Extinction correction: none
 1 restraint Absolute structure: Flack (1983), 989 Friedel pairs
 Primary atom site location: structure-invariant direct methods Flack parameter: 0.046 (17)
 Secondary atom site location: difference Fourier map

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Mn1	0.38539 (2)	0.37818 (3)	0.51871 (2)	0.01386 (8)
O1	0.36109 (12)	0.12118 (15)	0.32716 (9)	0.0186 (3)
O2	0.65501 (11)	0.32649 (17)	0.63464 (9)	0.0236 (4)
N1	0.30418 (13)	0.22376 (19)	0.58708 (11)	0.0194 (4)
N2	0.19272 (14)	0.08610 (19)	0.63164 (11)	0.0201 (4)
N3	0.40259 (14)	0.58964 (19)	0.46979 (11)	0.0185 (4)
N4	0.41706 (15)	0.74233 (19)	0.38026 (12)	0.0204 (4)
N5	0.25565 (14)	0.37585 (17)	0.44149 (10)	0.0157 (4)
N6	0.42608 (13)	0.24628 (19)	0.42643 (11)	0.0151 (4)
N7	0.53284 (12)	0.31457 (18)	0.54386 (10)	0.0156 (4)
N8	0.42419 (13)	0.48437 (18)	0.62717 (10)	0.0156 (4)
C1	0.21158 (16)	0.1926 (2)	0.58814 (13)	0.0206 (5)
H1	0.1635	0.2405	0.5610	0.025*
C2	0.27962 (17)	0.0447 (3)	0.66076 (15)	0.0283 (6)
H2	0.2902	-0.0294	0.6935	0.034*
C3	0.34717 (18)	0.1303 (2)	0.63362 (15)	0.0260 (5)
H3	0.4143	0.1267	0.6449	0.031*
C4	0.09803 (17)	0.0263 (3)	0.64689 (15)	0.0266 (5)
H4A	0.0520	0.0549	0.6083	0.040*
H4B	0.0751	0.0564	0.6961	0.040*
H4C	0.1035	-0.0724	0.6465	0.040*
C5	0.38396 (16)	0.6197 (2)	0.39912 (15)	0.0183 (5)
H5	0.3509	0.5611	0.3656	0.022*
C6	0.45966 (18)	0.7961 (2)	0.44268 (14)	0.0239 (5)
H6	0.4896	0.8822	0.4470	0.029*
C7	0.45043 (16)	0.7012 (2)	0.49738 (13)	0.0216 (5)
H7	0.4735	0.7105	0.5472	0.026*
C8	0.4078 (2)	0.8059 (2)	0.30677 (15)	0.0271 (5)
H8A	0.3726	0.7453	0.2730	0.041*
H8B	0.4724	0.8239	0.2863	0.041*

supplementary materials

H8C	0.3723	0.8912	0.3117	0.041*
C9	0.17954 (16)	0.4603 (2)	0.44273 (13)	0.0183 (5)
H9	0.1700	0.5150	0.4858	0.022*
C10	0.11453 (15)	0.4712 (2)	0.38434 (14)	0.0200 (5)
H10	0.0623	0.5334	0.3865	0.024*
C11	0.12757 (17)	0.3891 (2)	0.32266 (15)	0.0206 (5)
H11	0.0835	0.3932	0.2819	0.025*
C12	0.20533 (16)	0.3006 (2)	0.32047 (13)	0.0188 (5)
H12	0.2149	0.2430	0.2785	0.023*
C13	0.26922 (15)	0.2974 (2)	0.38068 (13)	0.0158 (4)
C14	0.35901 (16)	0.2101 (2)	0.37724 (12)	0.0147 (4)
C15	0.51905 (15)	0.1872 (2)	0.43010 (13)	0.0151 (4)
C16	0.55961 (17)	0.1028 (2)	0.37494 (14)	0.0178 (5)
H16	0.5224	0.0787	0.3323	0.021*
C17	0.65348 (17)	0.0543 (2)	0.38227 (14)	0.0212 (5)
H17	0.6800	-0.0026	0.3447	0.025*
C18	0.70854 (17)	0.0888 (2)	0.44413 (14)	0.0216 (5)
H18	0.7723	0.0542	0.4492	0.026*
C19	0.67119 (15)	0.1736 (2)	0.49883 (13)	0.0193 (5)
H19	0.7102	0.1978	0.5405	0.023*
C20	0.57629 (15)	0.2242 (2)	0.49336 (13)	0.0163 (5)
C21	0.57588 (16)	0.3589 (2)	0.60555 (13)	0.0171 (5)
C22	0.51797 (15)	0.4679 (2)	0.64610 (12)	0.0160 (4)
C23	0.56201 (16)	0.5497 (2)	0.69906 (13)	0.0200 (5)
H23	0.6280	0.5363	0.7119	0.024*
C24	0.50893 (18)	0.6512 (2)	0.73299 (13)	0.0234 (5)
H24	0.5385	0.7100	0.7683	0.028*
C25	0.41228 (18)	0.6657 (2)	0.71479 (14)	0.0213 (5)
H25	0.3737	0.7328	0.7383	0.026*
C26	0.37272 (17)	0.5799 (2)	0.66128 (14)	0.0187 (5)
H26	0.3063	0.5898	0.6486	0.022*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mn1	0.01291 (14)	0.01638 (15)	0.01230 (15)	-0.00013 (12)	-0.00020 (14)	-0.00077 (16)
O1	0.0198 (8)	0.0179 (8)	0.0181 (9)	-0.0009 (6)	0.0002 (7)	-0.0041 (7)
O2	0.0145 (8)	0.0335 (9)	0.0227 (9)	0.0036 (7)	-0.0046 (7)	-0.0034 (8)
N1	0.0194 (10)	0.0207 (10)	0.0181 (10)	-0.0003 (8)	0.0007 (8)	0.0018 (8)
N2	0.0224 (10)	0.0204 (10)	0.0175 (10)	-0.0027 (8)	0.0029 (8)	-0.0006 (8)
N3	0.0208 (10)	0.0178 (9)	0.0170 (10)	0.0010 (7)	0.0003 (8)	0.0005 (8)
N4	0.0258 (10)	0.0165 (10)	0.0188 (10)	0.0017 (8)	0.0023 (8)	0.0012 (8)
N5	0.0172 (9)	0.0148 (9)	0.0149 (10)	-0.0009 (7)	0.0003 (8)	0.0010 (7)
N6	0.0153 (9)	0.0161 (8)	0.0138 (9)	0.0004 (7)	0.0029 (7)	0.0003 (7)
N7	0.0120 (9)	0.0188 (9)	0.0161 (9)	0.0009 (7)	0.0013 (7)	0.0014 (7)
N8	0.0171 (9)	0.0181 (9)	0.0116 (9)	-0.0019 (7)	0.0007 (7)	0.0020 (8)
C1	0.0217 (12)	0.0219 (12)	0.0181 (12)	-0.0003 (10)	0.0002 (9)	0.0028 (9)
C2	0.0294 (13)	0.0272 (13)	0.0282 (14)	0.0023 (10)	0.0008 (11)	0.0096 (11)

C3	0.0206 (12)	0.0314 (14)	0.0259 (13)	0.0031 (10)	-0.0013 (11)	0.0050 (11)
C4	0.0241 (12)	0.0291 (13)	0.0266 (13)	-0.0073 (10)	0.0035 (10)	0.0032 (11)
C5	0.0177 (12)	0.0151 (12)	0.0220 (13)	-0.0017 (8)	0.0020 (9)	-0.0010 (9)
C6	0.0309 (13)	0.0188 (11)	0.0220 (13)	-0.0047 (10)	0.0013 (10)	0.0000 (10)
C7	0.0231 (11)	0.0213 (12)	0.0203 (12)	-0.0039 (9)	0.0017 (9)	-0.0002 (9)
C8	0.0400 (15)	0.0218 (12)	0.0197 (12)	0.0026 (11)	0.0031 (11)	0.0045 (10)
C9	0.0183 (11)	0.0192 (11)	0.0174 (11)	0.0009 (9)	0.0038 (9)	-0.0002 (9)
C10	0.0158 (11)	0.0207 (11)	0.0235 (12)	0.0008 (9)	0.0015 (9)	0.0040 (10)
C11	0.0188 (12)	0.0225 (12)	0.0206 (12)	-0.0035 (9)	-0.0038 (9)	0.0054 (10)
C12	0.0213 (12)	0.0187 (11)	0.0163 (11)	-0.0027 (9)	0.0005 (9)	-0.0002 (9)
C13	0.0178 (11)	0.0124 (10)	0.0172 (11)	-0.0039 (8)	0.0033 (9)	-0.0002 (9)
C14	0.0177 (11)	0.0132 (10)	0.0131 (10)	-0.0025 (8)	0.0019 (9)	0.0041 (9)
C15	0.0161 (11)	0.0143 (10)	0.0151 (11)	-0.0006 (8)	0.0021 (9)	0.0032 (9)
C16	0.0218 (12)	0.0164 (11)	0.0153 (11)	-0.0013 (9)	0.0025 (9)	-0.0013 (9)
C17	0.0240 (12)	0.0187 (12)	0.0209 (12)	0.0018 (10)	0.0070 (10)	-0.0015 (10)
C18	0.0142 (11)	0.0224 (12)	0.0283 (13)	0.0030 (9)	0.0036 (10)	0.0029 (10)
C19	0.0145 (10)	0.0216 (11)	0.0218 (13)	0.0010 (8)	0.0004 (9)	0.0027 (9)
C20	0.0148 (10)	0.0166 (10)	0.0173 (11)	-0.0028 (8)	0.0019 (8)	0.0033 (9)
C21	0.0144 (11)	0.0212 (12)	0.0157 (11)	-0.0030 (9)	0.0007 (9)	0.0011 (9)
C22	0.0171 (10)	0.0175 (11)	0.0135 (11)	-0.0024 (8)	0.0025 (8)	0.0051 (9)
C23	0.0174 (11)	0.0227 (12)	0.0199 (12)	-0.0047 (9)	-0.0023 (9)	0.0004 (10)
C24	0.0327 (14)	0.0209 (12)	0.0168 (12)	-0.0043 (10)	-0.0057 (10)	-0.0014 (10)
C25	0.0278 (12)	0.0192 (11)	0.0168 (11)	0.0025 (9)	-0.0008 (10)	-0.0006 (10)
C26	0.0186 (11)	0.0192 (11)	0.0181 (12)	0.0010 (9)	-0.0011 (9)	0.0017 (10)

Geometric parameters (Å, °)

Mn1—N6	2.1764 (19)	C6—C7	1.361 (3)
Mn1—N7	2.1794 (18)	C6—H6	0.9500
Mn1—N1	2.2552 (19)	C7—H7	0.9500
Mn1—N5	2.2621 (19)	C8—H8A	0.9800
Mn1—N8	2.2684 (19)	C8—H8B	0.9800
Mn1—N3	2.280 (2)	C8—H8C	0.9800
O1—C14	1.255 (3)	C9—C10	1.381 (3)
O2—C21	1.252 (3)	C9—H9	0.9500
N1—C1	1.316 (3)	C10—C11	1.381 (4)
N1—C3	1.378 (3)	C10—H10	0.9500
N2—C1	1.335 (3)	C11—C12	1.387 (3)
N2—C2	1.371 (3)	C11—H11	0.9500
N2—C4	1.462 (3)	C12—C13	1.392 (3)
N3—C5	1.322 (3)	C12—H12	0.9500
N3—C7	1.378 (3)	C13—C14	1.513 (3)
N4—C5	1.340 (3)	C15—C16	1.408 (3)
N4—C6	1.369 (3)	C15—C20	1.427 (3)
N4—C8	1.461 (3)	C16—C17	1.389 (3)
N5—C9	1.343 (3)	C16—H16	0.9500
N5—C13	1.348 (3)	C17—C18	1.384 (3)
N6—C14	1.326 (3)	C17—H17	0.9500
N6—C15	1.413 (3)	C18—C19	1.387 (3)

supplementary materials

N7—C21	1.327 (3)	C18—H18	0.9500
N7—C20	1.405 (3)	C19—C20	1.407 (3)
N8—C26	1.330 (3)	C19—H19	0.9500
N8—C22	1.349 (3)	C21—C22	1.526 (3)
C1—H1	0.9500	C22—C23	1.386 (3)
C2—C3	1.350 (3)	C23—C24	1.383 (3)
C2—H2	0.9500	C23—H23	0.9500
C3—H3	0.9500	C24—C25	1.382 (3)
C4—H4A	0.9800	C24—H24	0.9500
C4—H4B	0.9800	C25—C26	1.390 (4)
C4—H4C	0.9800	C25—H25	0.9500
C5—H5	0.9500	C26—H26	0.9500
N6—Mn1—N7	75.02 (7)	N3—C7—H7	125.0
N6—Mn1—N1	97.62 (7)	N4—C8—H8A	109.5
N7—Mn1—N1	99.09 (7)	N4—C8—H8B	109.5
N6—Mn1—N5	74.74 (7)	H8A—C8—H8B	109.5
N7—Mn1—N5	149.74 (7)	N4—C8—H8C	109.5
N1—Mn1—N5	85.93 (7)	H8A—C8—H8C	109.5
N6—Mn1—N8	149.72 (7)	H8B—C8—H8C	109.5
N7—Mn1—N8	74.73 (7)	N5—C9—C10	123.1 (2)
N1—Mn1—N8	88.21 (7)	N5—C9—H9	118.5
N5—Mn1—N8	135.47 (7)	C10—C9—H9	118.5
N6—Mn1—N3	103.48 (8)	C11—C10—C9	118.2 (2)
N7—Mn1—N3	104.27 (7)	C11—C10—H10	120.9
N1—Mn1—N3	151.79 (7)	C9—C10—H10	120.9
N5—Mn1—N3	81.85 (7)	C10—C11—C12	119.7 (2)
N8—Mn1—N3	82.97 (7)	C10—C11—H11	120.2
C1—N1—C3	104.67 (19)	C12—C11—H11	120.2
C1—N1—Mn1	130.57 (15)	C11—C12—C13	119.0 (2)
C3—N1—Mn1	124.52 (15)	C11—C12—H12	120.5
C1—N2—C2	106.58 (19)	C13—C12—H12	120.5
C1—N2—C4	127.1 (2)	N5—C13—C12	121.4 (2)
C2—N2—C4	126.3 (2)	N5—C13—C14	118.38 (19)
C5—N3—C7	104.76 (19)	C12—C13—C14	120.1 (2)
C5—N3—Mn1	123.51 (16)	O1—C14—N6	130.2 (2)
C7—N3—Mn1	130.40 (16)	O1—C14—C13	116.61 (19)
C5—N4—C6	107.1 (2)	N6—C14—C13	113.10 (19)
C5—N4—C8	125.9 (2)	C16—C15—N6	125.1 (2)
C6—N4—C8	127.0 (2)	C16—C15—C20	119.1 (2)
C9—N5—C13	118.7 (2)	N6—C15—C20	115.75 (19)
C9—N5—Mn1	127.17 (15)	C17—C16—C15	120.7 (2)
C13—N5—Mn1	112.77 (14)	C17—C16—H16	119.7
C14—N6—C15	123.67 (19)	C15—C16—H16	119.7
C14—N6—Mn1	118.88 (14)	C18—C17—C16	120.2 (2)
C15—N6—Mn1	116.61 (14)	C18—C17—H17	119.9
C21—N7—C20	123.51 (18)	C16—C17—H17	119.9
C21—N7—Mn1	119.67 (14)	C17—C18—C19	120.4 (2)
C20—N7—Mn1	116.81 (14)	C17—C18—H18	119.8
C26—N8—C22	118.98 (19)	C19—C18—H18	119.8

C26—N8—Mn1	126.43 (15)	C18—C19—C20	120.9 (2)
C22—N8—Mn1	112.66 (14)	C18—C19—H19	119.6
N1—C1—N2	112.5 (2)	C20—C19—H19	119.6
N1—C1—H1	123.7	N7—C20—C19	125.5 (2)
N2—C1—H1	123.7	N7—C20—C15	115.78 (18)
C3—C2—N2	106.4 (2)	C19—C20—C15	118.7 (2)
C3—C2—H2	126.8	O2—C21—N7	130.6 (2)
N2—C2—H2	126.8	O2—C21—C22	116.22 (19)
C2—C3—N1	109.8 (2)	N7—C21—C22	113.13 (19)
C2—C3—H3	125.1	N8—C22—C23	121.5 (2)
N1—C3—H3	125.1	N8—C22—C21	118.04 (19)
N2—C4—H4A	109.5	C23—C22—C21	120.44 (19)
N2—C4—H4B	109.5	C24—C23—C22	119.4 (2)
H4A—C4—H4B	109.5	C24—C23—H23	120.3
N2—C4—H4C	109.5	C22—C23—H23	120.3
H4A—C4—H4C	109.5	C25—C24—C23	119.0 (2)
H4B—C4—H4C	109.5	C25—C24—H24	120.5
N3—C5—N4	112.2 (2)	C23—C24—H24	120.5
N3—C5—H5	123.9	C24—C25—C26	118.6 (2)
N4—C5—H5	123.9	C24—C25—H25	120.7
C7—C6—N4	106.0 (2)	C26—C25—H25	120.7
C7—C6—H6	127.0	N8—C26—C25	122.6 (2)
N4—C6—H6	127.0	N8—C26—H26	118.7
C6—C7—N3	109.9 (2)	C25—C26—H26	118.7
C6—C7—H7	125.0		
N6—Mn1—N1—C1	-92.7 (2)	Mn1—N1—C3—C2	-174.18 (17)
N7—Mn1—N1—C1	-168.6 (2)	C7—N3—C5—N4	0.2 (3)
N5—Mn1—N1—C1	-18.7 (2)	Mn1—N3—C5—N4	-167.83 (14)
N8—Mn1—N1—C1	117.1 (2)	C6—N4—C5—N3	-0.3 (3)
N3—Mn1—N1—C1	45.6 (3)	C8—N4—C5—N3	-179.6 (2)
N6—Mn1—N1—C3	80.7 (2)	C5—N4—C6—C7	0.3 (3)
N7—Mn1—N1—C3	4.8 (2)	C8—N4—C6—C7	179.6 (2)
N5—Mn1—N1—C3	154.7 (2)	N4—C6—C7—N3	-0.1 (3)
N8—Mn1—N1—C3	-69.46 (19)	C5—N3—C7—C6	0.0 (3)
N3—Mn1—N1—C3	-140.97 (19)	Mn1—N3—C7—C6	166.84 (16)
N6—Mn1—N3—C5	34.70 (19)	C13—N5—C9—C10	-0.4 (3)
N7—Mn1—N3—C5	112.38 (18)	Mn1—N5—C9—C10	165.25 (17)
N1—Mn1—N3—C5	-102.6 (2)	N5—C9—C10—C11	1.5 (3)
N5—Mn1—N3—C5	-37.38 (18)	C9—C10—C11—C12	-1.0 (3)
N8—Mn1—N3—C5	-175.39 (18)	C10—C11—C12—C13	-0.5 (3)
N6—Mn1—N3—C7	-130.0 (2)	C9—N5—C13—C12	-1.2 (3)
N7—Mn1—N3—C7	-52.4 (2)	Mn1—N5—C13—C12	-168.82 (16)
N1—Mn1—N3—C7	92.6 (2)	C9—N5—C13—C14	175.61 (19)
N5—Mn1—N3—C7	157.9 (2)	Mn1—N5—C13—C14	8.0 (2)
N8—Mn1—N3—C7	19.9 (2)	C11—C12—C13—N5	1.6 (3)
N6—Mn1—N5—C9	-166.23 (19)	C11—C12—C13—C14	-175.1 (2)
N7—Mn1—N5—C9	-164.19 (16)	C15—N6—C14—O1	1.1 (4)
N1—Mn1—N5—C9	94.73 (18)	Mn1—N6—C14—O1	-168.07 (19)
N8—Mn1—N5—C9	11.5 (2)	C15—N6—C14—C13	-174.87 (18)

supplementary materials

N3—Mn1—N5—C9	-59.78 (18)	Mn1—N6—C14—C13	16.0 (2)
N6—Mn1—N5—C13	0.09 (14)	N5—C13—C14—O1	167.57 (19)
N7—Mn1—N5—C13	2.1 (2)	C12—C13—C14—O1	-15.6 (3)
N1—Mn1—N5—C13	-98.95 (15)	N5—C13—C14—N6	-15.9 (3)
N8—Mn1—N5—C13	177.77 (13)	C12—C13—C14—N6	161.0 (2)
N3—Mn1—N5—C13	106.54 (15)	C14—N6—C15—C16	12.7 (3)
N7—Mn1—N6—C14	171.66 (18)	Mn1—N6—C15—C16	-177.88 (17)
N1—Mn1—N6—C14	74.24 (17)	C14—N6—C15—C20	-170.9 (2)
N5—Mn1—N6—C14	-9.40 (16)	Mn1—N6—C15—C20	-1.6 (2)
N8—Mn1—N6—C14	173.83 (15)	N6—C15—C16—C17	177.1 (2)
N3—Mn1—N6—C14	-86.90 (17)	C20—C15—C16—C17	0.9 (3)
N7—Mn1—N6—C15	1.75 (14)	C15—C16—C17—C18	-0.1 (3)
N1—Mn1—N6—C15	-95.67 (15)	C16—C17—C18—C19	-1.0 (4)
N5—Mn1—N6—C15	-179.31 (16)	C17—C18—C19—C20	1.2 (3)
N8—Mn1—N6—C15	3.9 (2)	C21—N7—C20—C19	-3.1 (3)
N3—Mn1—N6—C15	103.19 (15)	Mn1—N7—C20—C19	178.62 (17)
N6—Mn1—N7—C21	179.90 (17)	C21—N7—C20—C15	179.81 (19)
N1—Mn1—N7—C21	-84.58 (16)	Mn1—N7—C20—C15	1.5 (2)
N5—Mn1—N7—C21	177.87 (15)	C18—C19—C20—N7	-177.3 (2)
N8—Mn1—N7—C21	1.03 (15)	C18—C19—C20—C15	-0.3 (3)
N3—Mn1—N7—C21	79.48 (17)	C16—C15—C20—N7	176.57 (19)
N6—Mn1—N7—C20	-1.75 (14)	N6—C15—C20—N7	0.0 (3)
N1—Mn1—N7—C20	93.76 (15)	C16—C15—C20—C19	-0.7 (3)
N5—Mn1—N7—C20	-3.8 (2)	N6—C15—C20—C19	-177.28 (18)
N8—Mn1—N7—C20	179.38 (15)	C20—N7—C21—O2	-5.9 (4)
N3—Mn1—N7—C20	-102.17 (15)	Mn1—N7—C21—O2	172.31 (19)
N6—Mn1—N8—C26	168.94 (17)	C20—N7—C21—C22	173.58 (18)
N7—Mn1—N8—C26	171.1 (2)	Mn1—N7—C21—C22	-8.2 (2)
N1—Mn1—N8—C26	-88.97 (19)	C26—N8—C22—C23	-1.3 (3)
N5—Mn1—N8—C26	-6.6 (2)	Mn1—N8—C22—C23	163.93 (17)
N3—Mn1—N8—C26	64.18 (19)	C26—N8—C22—C21	-179.3 (2)
N6—Mn1—N8—C22	5.1 (2)	Mn1—N8—C22—C21	-14.1 (2)
N7—Mn1—N8—C22	7.25 (14)	O2—C21—C22—N8	-165.29 (19)
N1—Mn1—N8—C22	107.17 (15)	N7—C21—C22—N8	15.1 (3)
N5—Mn1—N8—C22	-170.47 (13)	O2—C21—C22—C23	16.7 (3)
N3—Mn1—N8—C22	-99.68 (15)	N7—C21—C22—C23	-162.9 (2)
C3—N1—C1—N2	-0.2 (3)	N8—C22—C23—C24	-0.4 (3)
Mn1—N1—C1—N2	174.15 (16)	C21—C22—C23—C24	177.6 (2)
C2—N2—C1—N1	-0.3 (3)	C22—C23—C24—C25	1.9 (3)
C4—N2—C1—N1	178.6 (2)	C23—C24—C25—C26	-1.8 (4)
C1—N2—C2—C3	0.6 (3)	C22—N8—C26—C25	1.4 (3)
C4—N2—C2—C3	-178.2 (2)	Mn1—N8—C26—C25	-161.57 (18)
N2—C2—C3—N1	-0.8 (3)	C24—C25—C26—N8	0.2 (4)
C1—N1—C3—C2	0.6 (3)		

Fig. 1

