

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3,3'-(2,2'-Bi-1*H*-imidazole-1,1'-diyl)di-propanol

Tao Zhang and Hong-Ze Liang*

State Key Laboratory Base of Novel Functional Materials and Preparation Science, Institute of Solid Materials Chemistry, Faculty of Materials Science and Chemical Engineering, Ningbo University, Ningbo 315211, People's Republic of China

Correspondence e-mail: lianghongze@nbu.edu.cn


Received 9 December 2008; accepted 19 December 2008

 Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.036; wR factor = 0.104; data-to-parameter ratio = 16.3.

In the title compound, $\text{C}_{12}\text{H}_{18}\text{N}_4\text{O}_2$, unlike other unconjugated disubstituted biimidazole derivatives reported so far, the two imidazole rings in a *trans* conformation exhibit a large planar rotation angle of $51.27(4)^\circ$, and consist of half-molecule asymmetric units related by a twofold rotation. The molecules are linked into a three-dimensional framework with a parallel laminated construction *via* $\text{O}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{O}$ interactions.

Related literature

For background to 2,2'-biimidazole derivatives, see: Forster *et al.* (2004); Fortin & Beauchamp (2000); Fu *et al.* (2007); Ion *et al.* (2007); Mao *et al.* (2003); Pereira *et al.* (2006); Xiao & Shreeve (2005); Xiao *et al.* (2004). For other unconjugated 1,1'-disubstituted compounds, see: Barnett *et al.* (1997, 2002); Secondo *et al.* (1996, 1997). For the synthesis, see: Barnett *et al.* (1999)



Experimental

Crystal data

 $\text{C}_{12}\text{H}_{18}\text{N}_4\text{O}_2$
 $M_r = 250.30$

 Monoclinic, $C2/c$
 $a = 15.812(3)$ Å
 $b = 9.5961(19)$ Å
 $c = 9.3194(19)$ Å
 $\beta = 119.44(3)^\circ$
 $V = 1231.5(6)$ Å³
 $Z = 4$

 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 295(2)$ K
 $0.56 \times 0.48 \times 0.37$ mm

Data collection

 Rigaku R-AXIS RAPID
 diffractometer
 Absorption correction: multi-scan
 (*ABSCOR*; Higashi, 1995)
 $T_{\min} = 0.948$, $T_{\max} = 0.970$

 4715 measured reflections
 1400 independent reflections
 1183 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.015$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.104$
 $S = 1.07$
 1400 reflections
 86 parameters

 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.17$ e Å⁻³
 $\Delta\rho_{\min} = -0.23$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H9}\cdots\text{N2}^i$	0.89 (2)	1.92 (2)	2.8047 (17)	175
$\text{C2}-\text{H2}\cdots\text{O1}^ii$	0.93	2.59	3.5019 (17)	166

 Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $x - \frac{1}{2}, y - \frac{1}{2}, z - 1$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Rigaku/MSc, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This project was sponsored by the Scientific Research Foundation of the State Education Ministry for Returned Overseas Chinese Scholars (2006331), the Critical Projects in Science and Technology Department of Zhejiang Province (2007 C21113), the Education Committee of Zhejiang Province (20061696), the Natural Science Foundation of Ningbo City (2007 A610021), K. C. Wong Magna Fund in Ningbo University and Ningbo University (2005062). We thank Mr W. Xu for collecting the crystal data.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2693).

References

- Barnett, W. M., Baughman, R. G., Collier, H. L. & Vizuete, W. G. (1999). *J. Chem. Crystallogr.* **29**, 765–768.
 Barnett, W. M., Baughman, R. G., Secondo, P. M. & Hermansen, C. J. (2002). *Acta Cryst.* **C58**, o565–o567.
 Barnett, W. M., Lin, G., Collier, H. L. & Baughman, R. G. (1997). *J. Chem. Crystallogr.* **27**, 423–427.
 Forster, R. J., Walsh, D. A., Mano, N., Mao, F. & Heller, A. (2004). *Langmuir*, **20**, 862–868.
 Fortin, S. & Beauchamp, A. L. (2000). *Inorg. Chem.* **39**, 4886–4893.
 Fu, Y. M., Zhao, Y. H., Lan, Y. Q., Wang, Y., Qiu, Y. Q., Shao, K. Z. & Su, Z. M. (2007). *Inorg. Chem. Commun.* **10**, 720–723.
 Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
 Ion, L., Morales, D., Nieto, S., Perez, J., Riera, L., Riera, V., Miguel, D., Kowenicki, R. A. & Mcpartlin, M. (2007). *Inorg. Chem.* **46**, 2846–2853.
 Mao, F., Mano, N. & Heller, A. (2003). *J. Am. Chem. Soc.* **125**, 4951–4957.
 Pereira, C. C. L., Costa, P. J., Calhorda, M. J., Freire, C., Rodrigues, S. S., Herdtweck, E. & Romão, C. C. (2006). *Organometallics*, **25**, 5223–5234.
 Rigaku (1998). *RAPID-AUTO*. Rigaku Cooperation, Tokyo, Japan.
 Rigaku/MSc (2002). *CrystalStructure*. Rigaku/MSc Inc., The Woodlands, Texas, USA.

Secondo, P. M., Barnett, W. M., Collier, H. L. & Baughman, R. G. (1996). *Acta Cryst.* **C52**, 2636–2638.

Secondo, P. M., Baughman, R. G. & Collier, H. L. (1997). *J. Chem. Crystallogr.* **27**, 371–375.

Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

Xiao, J. C. & Shreeve, J. M. (2005). *J. Org. Chem.* **70**, 3072–3078.

Xiao, J. C., Twamley, B. & Shreeve, J. M. (2004). *Org. Lett.* **6**, 3845–3847.

supplementary materials

Acta Cryst. (2009). E65, o213-o214 [doi:10.1107/S1600536808043377]

3,3'-(2,2'-Bi-1*H*-imidazole-1,1'-diyl)dipropanol

T. Zhang and H.-Z. Liang

Comment

2,2'-Biimidazole (H₂biim) derivatives as versatile ligands are widely used in the construction of metal complexes (Ion *et al.*, 2007; Pereira *et al.*, 2006; Fortin *et al.*, 2000). In addition to the study of metal complexes, attention has also been devoted to the polymeric systems (Fu *et al.*, 2007; Forster *et al.*, 2004; Mao *et al.*, 2003) and the ionic liquids (Xiao *et al.*, 2005, 2004). The title compound (I), as a hydroxy-terminated derivative can be easily synthesized by Michael addition of dianion biim²⁻ to alkyl acrylate (Barnett *et al.*, 1999) and subsequent mild reduction by NaBH₄ in an excellent yield.

The crystal structure of (I), shown in Fig. 1, adopts a *trans* conformation, and consists of half-molecule asymmetric units related by a twofold rotation. Exceptionally, unlike the other unconjugated 1,1'-disubstituted compounds reported before (Barnett *et al.*, 1997, 2002; Secondo *et al.*, 1996, 1996), the imidazole rings of the investigated compound (I) exhibit a rather large torsion angle of 51.27 (4)°, as shown in Fig. 1. The terminal hydroxyl groups are responsible for this noncoplanarity. The imidazole rings are forced to rotate by intermolecular interactions *via* strong O–H···N hydrogen bonds. The N1–C4–C5–C6 moiety is essentially planar with an r.m.s. deviation of 0.054 Å. This plane is at angle of 71.90 (7)° with respect to the adjacent imidazole ring, which is consistent with the reported analogous compounds referenced before. While the atom O1 does not lie in the N1–C4–C5–C6 plane, rather it lies 1.116 Å from this plane. The bond length and bond angle in (I) are within normal ranges.

The molecules of (I) are linked into a three-dimensional framework with parallel laminated construction by a combination of one strong O–H···N hydrogen bond and the other weak C–H···O hydrogen bond (Table 1). It can be analyzed in terms of thousands of one-dimensional substructures, linked by C–H···O hydrogen bonds, which generate series of two-dimensional sheets through C–H···O hydrogen bonds (Fig.2).

Experimental

1,1'-Di(ethylpropionato)-2,2'-biimidazole (0.80 g, 2.40 mmol) was dissolved in absolute ethyl alcohol (50 ml). To this mixture was added batchly NaBH₄ (0.906 g, 24.0 mmol). The mixture was heated to reflux for 3 h. Evaporating the solvent and added to 30 ml H₂O yield a transparent solution. Adjusting the PH of the solution to 8–9 with 1M HCl led to a white suspension. The mixture was filtered, then the filtrate extracted with chloroform (10 ml) three times. The combined organic layer was washed with H₂O (5 ml) two times, dried by MgSO₄. After evaporating the solvent, the residue was chromatographed on silica gel. Elution with CH₃CO₂C₂H₅/C₂H₅OH (10:1) afforded a white solid (0.52 g, 86.7%). Tetragonal crystals of the titled compound were formed by evaporating saturated petroleum ether/chloroform solution slowly.

Similarly, the title compound(I) can be synthesized by reduction of other 1,1'-di(alkylpropionato)-2,2'-biimidazole (alkyl = methyl, butyl) which was prepared according to the published procedure (Barnett *et al.*, 1999) with NaBH₄ in anhydrous ethyl alcohol or with LiAlH₄ in anhydrous ethereal solution. The yields ranges from 44 to 85°.

Refinement

All H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms (C—H = 0.93 and = 0.97 Å; O—H = 0.82 Å) and $U_{\text{iso}}(\text{H})$ values were taken to be equal to $1.2 U_{\text{eq}}(\text{C})$ and $1.5 U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. A perspective view of (I) with the atom-labelling scheme, showing that the two imidazole rings are distinctly non-coplanar. Displacement ellipsoids are drawn at the 40% probability level and H atoms are shown as small spheres of arbitrary radius.




Fig. 2. A perspective view of the three-dimensional framework with parallellaminated construction containing two sheets from [011] direction, showing the packing mode and the interactions of hydrogen bonds O—H...N (pink dashed lines in the electronic version of the paper) and C—H...O (green dashed lines in the electronic version). All H atoms not involved in the hydrogen-bond motifs have been omitted for clarity.

3,3'-(2,2'-Bi-1*H*-imidazole-1,1'-diyl)dipropanol

Crystal data

$\text{C}_{12}\text{H}_{18}\text{N}_4\text{O}_2$

$M_r = 250.30$

Monoclinic, $C2/c$

Hall symbol: $-C\ 2yc$

$a = 15.812\ (3)\ \text{\AA}$

$b = 9.5961\ (19)\ \text{\AA}$

$c = 9.3194\ (19)\ \text{\AA}$

$\beta = 119.44\ (3)^\circ$

$V = 1231.5\ (6)\ \text{\AA}^3$

$Z = 4$

$F_{000} = 536$

$D_x = 1.350\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 1400 reflections

$\theta = 3.1\text{--}27.5^\circ$

$\mu = 0.10\ \text{mm}^{-1}$

$T = 295\ (2)\ \text{K}$

Block, colourless

$0.56 \times 0.48 \times 0.37\ \text{mm}$

Data collection

Rigaku R-AXIS RAPID
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 295\ (2)\ \text{K}$

ω scans

1400 independent reflections

1183 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.015$

$\theta_{\text{max}} = 27.5^\circ$

$\theta_{\text{min}} = 3.1^\circ$

Absorption correction: multi-scan
(ABSCOR; Higashi, 1995) $h = -20 \rightarrow 20$
 $T_{\min} = 0.948$, $T_{\max} = 0.970$ $k = -12 \rightarrow 12$
 4715 measured reflections $l = -12 \rightarrow 12$

Refinement

Refinement on F^2 Secondary atom site location: difference Fourier map
 Least-squares matrix: full Hydrogen site location: inferred from neighbouring sites
 $R[F^2 > 2\sigma(F^2)] = 0.036$ H atoms treated by a mixture of independent and constrained refinement
 $wR(F^2) = 0.104$ $w = 1/[\sigma^2(F_o^2) + (0.0631P)^2 + 0.3657P]$
 $S = 1.07$ where $P = (F_o^2 + 2F_c^2)/3$
 1400 reflections $(\Delta/\sigma)_{\max} < 0.001$
 86 parameters $\Delta\rho_{\max} = 0.17 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.23 \text{ e } \text{\AA}^{-3}$
 Primary atom site location: structure-invariant direct methods Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.03257 (9)	0.22372 (12)	0.47224 (14)	0.0366 (3)
H1	0.0646	0.2434	0.4134	0.044*
C2	-0.05565 (9)	0.16313 (12)	0.41318 (14)	0.0375 (3)
H2	-0.0946	0.1336	0.3050	0.045*
C3	-0.00459 (7)	0.20468 (10)	0.66867 (13)	0.0280 (3)
C4	0.15367 (7)	0.32897 (11)	0.74530 (14)	0.0328 (3)
H3	0.1970	0.3292	0.6993	0.039*
H4	0.1869	0.2829	0.8516	0.039*
C5	0.13141 (8)	0.47728 (12)	0.76898 (17)	0.0418 (3)
H5	0.0930	0.4771	0.8241	0.050*
H6	0.0930	0.5207	0.6620	0.050*
C6	0.22260 (8)	0.56240 (12)	0.86937 (16)	0.0390 (3)
H7	0.2635	0.5562	0.8189	0.047*

supplementary materials

H8	0.2048	0.6594	0.8674	0.047*
N1	0.06543 (6)	0.25027 (9)	0.63558 (11)	0.0300 (2)
N2	-0.07917 (6)	0.15167 (10)	0.53523 (11)	0.0340 (3)
O1	0.27596 (7)	0.51756 (11)	1.03437 (12)	0.0475 (3)
H9	0.3207 (14)	0.460 (2)	1.037 (2)	0.073 (6)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0423 (6)	0.0386 (6)	0.0319 (6)	0.0015 (5)	0.0206 (5)	0.0000 (4)
C2	0.0425 (6)	0.0373 (6)	0.0258 (6)	-0.0003 (4)	0.0114 (4)	-0.0030 (4)
C3	0.0253 (5)	0.0265 (5)	0.0287 (6)	0.0014 (4)	0.0107 (4)	0.0005 (4)
C4	0.0246 (5)	0.0324 (5)	0.0387 (6)	-0.0004 (4)	0.0134 (4)	-0.0013 (4)
C5	0.0285 (6)	0.0333 (6)	0.0525 (8)	0.0020 (4)	0.0114 (5)	-0.0050 (5)
C6	0.0330 (6)	0.0316 (5)	0.0477 (7)	-0.0014 (4)	0.0163 (5)	-0.0033 (5)
N1	0.0284 (5)	0.0313 (5)	0.0295 (5)	0.0000 (3)	0.0136 (4)	-0.0014 (3)
N2	0.0318 (5)	0.0351 (5)	0.0281 (5)	-0.0035 (4)	0.0093 (4)	-0.0020 (3)
O1	0.0407 (5)	0.0592 (6)	0.0391 (6)	0.0028 (4)	0.0168 (4)	-0.0102 (4)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.3536 (18)	C4—H3	0.9700
C1—N1	1.3692 (16)	C4—H4	0.9700
C1—H1	0.9300	C5—C6	1.5151 (16)
C2—N2	1.3638 (16)	C5—H5	0.9700
C2—H2	0.9300	C5—H6	0.9700
C3—N2	1.3245 (14)	C6—O1	1.4093 (17)
C3—N1	1.3595 (14)	C6—H7	0.9700
C3—C3 ⁱ	1.450 (2)	C6—H8	0.9700
C4—N1	1.4697 (14)	O1—H9	0.89 (2)
C4—C5	1.5081 (16)		
C2—C1—N1	106.51 (11)	C4—C5—H5	109.1
C2—C1—H1	126.7	C6—C5—H5	109.1
N1—C1—H1	126.7	C4—C5—H6	109.1
C1—C2—N2	110.15 (10)	C6—C5—H6	109.1
C1—C2—H2	124.9	H5—C5—H6	107.9
N2—C2—H2	124.9	O1—C6—C5	112.75 (11)
N2—C3—N1	111.05 (10)	O1—C6—H7	109.0
N2—C3—C3 ⁱ	124.54 (11)	C5—C6—H7	109.0
N1—C3—C3 ⁱ	124.26 (11)	O1—C6—H8	109.0
N1—C4—C5	112.13 (9)	C5—C6—H8	109.0
N1—C4—H3	109.2	H7—C6—H8	107.8
C5—C4—H3	109.2	C3—N1—C1	106.62 (10)
N1—C4—H4	109.2	C3—N1—C4	127.45 (10)
C5—C4—H4	109.2	C1—N1—C4	125.51 (10)
H3—C4—H4	107.9	C3—N2—C2	105.68 (10)
C4—C5—C6	112.29 (9)	C6—O1—H9	105.3 (12)

Symmetry codes: (i) $-x, y, -z+3/2$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O1—H9···N2 ⁱⁱ	0.89 (2)	1.92 (2)	2.8047 (17)	175
C2—H2···O1 ⁱⁱⁱ	0.93	2.59	3.5019 (17)	166

Symmetry codes: (ii) $x+1/2, -y+1/2, z+1/2$; (iii) $x-1/2, y-1/2, z-1$.

Fig. 1




Fig. 2

