

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

 5-Hydroxy-2-methyl-4*H*-pyran-4-one

 Muhammad Ashraf Shaheen,^a Christian G. Hartinger,^b
 M. Nawaz Tahir,^{c*} Ahmad Awan Shafiq^d and Bernhard K.
 Keppler^b
^aDepartment of Chemistry, University of Sargodha, Sargodha, Pakistan, ^bInstitute of Inorganic Chemistry, University of Vienna, Währinger Strasse 42, A-1090 Vienna, Austria, ^cDepartment of Physics, University of Sargodha, Sargodha, Pakistan, and ^dPAEC, PO Box No. 1114, Islamabad, Pakistan
 Correspondence e-mail: dmntahir_uos@yahoo.com


Received 25 January 2009; accepted 26 January 2009

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.053; wR factor = 0.131; data-to-parameter ratio = 17.3.

The title compound, $\text{C}_6\text{H}_6\text{O}_3$, is a member of the pyrone family. The molecules are planar (r.m.s. deviation of the asymmetric unit is 0.0248 Å, whereas that of the dimer is 0.0360 Å) and they are dimerized due to intermolecular O—H \cdots O hydrogen bonds. The dimers are connected to each other through hydrogen bonds involving the CH_3 group and the hydroxy O atom. There are π – π interactions between the centroids of the pyrone rings at a distance of 3.8552 (13) Å. A C—H \cdots π interaction also exists between the carbonyl group and the centroid CgA of the pyrone ring, with O \cdots CgA = 3.65 (1) Å and C \cdots CgA = 4.363 (2) Å.

Related literature

For general background, see: Aytimir *et al.* (1999); Erol & Yulug (1999). For studies involving metal complexes of allostol, see: Ma *et al.* (2004); Shaheen *et al.* (2008, 2008a). For crystal structures of related compounds, see: Tak *et al.* (1994); Rahman *et al.* (1997).



Experimental

Crystal data

 $\text{C}_6\text{H}_6\text{O}_3$
 $M_r = 126.11$

 Triclinic, $P\bar{1}$
 $a = 5.4467$ (4) Å

 $b = 7.3301$ (5) Å
 $c = 7.6945$ (5) Å
 $\alpha = 105.354$ (3) $^\circ$
 $\beta = 98.416$ (4) $^\circ$
 $\gamma = 100.008$ (4) $^\circ$
 $V = 285.68$ (4) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.12$ mm⁻¹
 $T = 296$ (2) K
 $0.22 \times 0.20 \times 0.10$ mm

Data collection

 Bruker Kappa APEXII CCD
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 2005)
 $T_{\min} = 0.970$, $T_{\max} = 0.986$

 6426 measured reflections
 1504 independent reflections
 713 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.045$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.131$
 $S = 1.00$
 1504 reflections
 87 parameters

 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.18$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.20$ e Å⁻³

Table 1

 Hydrogen-bond geometry (Å, $^\circ$).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O2—H2 \cdots O3	0.87 (3)	2.46 (2)	2.7853 (19)	103.1 (18)
O2—H2 \cdots O3 ⁱ	0.87 (3)	1.83 (3)	2.635 (2)	152 (2)
C6—H6A \cdots O2 ⁱⁱ	0.96	2.42	3.378 (3)	173

 Symmetry codes: (i) $-x - 1, -y, -z + 1$; (ii) $x, y, z - 1$; (iii) $x - 1, y, z$.

Data collection: APEX2 (Bruker, 2007); cell refinement: APEX2; data reduction: SAINT (Bruker, 2007); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997) and PLATON (Spek, 2003); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2715).

References

- Aytimir, M. D., Uzbay, T. & Erol, D. D. (1999). *Drug Res.* **49**, 250–254.
 Bruker (2005). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Erol, D. D. & Yulug, N. (1999). *Eur. J. Med. Chem.* **29**, 893–895.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Ma, Y., Luo, W., Quinn, P. J., Liu, Z. & Hider, R. C. (2004). *J. Med. Chem.* **47**, 6349–6362.
 Rahman, A., Nasreen, A., Akhtar, F., Shekhani, M. S., Clardy, J., Parvez, M. & Choudhary, M. I. (1997). *J. Nat. Prod.* **60**, 472–474.
 Shaheen, M. A., Berger, I., Alexey, A. N., Hartinger, C. G., Koroteev, M. P., Nifant, E. E. & Keppler, B. K. (2008). *Chem. Biodivers.* **5**, 1640–1644.
 Shaheen, M. A., Kandioller, W., Maria-Grazia, M. F., Alexey, A. N., Hartinger, C. G. & Keppler, B. K. (2008a). *Chem. Biodivers.* **5**, 2060–2066.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
 Tak, H., Fronczek, F. R. & Fischer, N. H. (1994). *Spectrosc. Lett.* **27**, 1431–1435.

supporting information

Acta Cryst. (2009). E65, o437 [doi:10.1107/S1600536809003158]

5-Hydroxy-2-methyl-4*H*-pyran-4-one

Muhammad Ashraf Shaheen, Christian G. Hartinger, M. Nawaz Tahir, Ahmad Awan Shafiq and Bernhard K. Keppler

S1. Comment

A variety of compounds with a 4(1*H*)-pyridinone structure have been synthesized and their biological activities studied extensively (Aytemir *et al.*, 1999; Erol & Yulug, 1999). Allomaltol, the title compound (I), (Fig. 1) and its derivatives have been exploited as iron chelators (Ma *et al.*, 2004). Ruthenium and osmium complexes of allomaltol were found to be effective in catalyzing the hydration of chloroacetonitriles (Shaheen *et al.*, 2008, 2008*a*).

The crystal structures of 3-Hydroxy-4-pyrone (Tak *et al.*, 1994) has been published which have same heterocyclic ring as of title compound. 3-Hydroxy-2-methyl-4*H*-pyran-4-one (Rahman *et al.*, 1997) has also been published which is chemical isomer of (I) but have different position of CH₃. The title compound has been prepared for various purposes such as complexation and as an intermediate ligand.


The heterocyclic ring is not regular as it has two C—C [1.426 (3), 1.446 (3) Å], two C=C [1.323 (3), 1.334 (3) Å] and two C—O [1.352 (3), 1.358 (3) Å], bonds respectively. Due to intra as well as intermolecular H-bonds (Table 1), the molecules are dimerized with central four-membered [O···H···O···H] ring, (Fig. 2). The dimers are linked to each other through H-bond between CH₃ and hydroxy groups. The molecules may be stabilized due to π - π interaction between the centroids of the ring A (O1/C1–C5). The distance between the centroids of CgA and CgAⁱ [Symmetry code: $i = -x, 1 - y, 1 - z$] is 3.8552 (13) Å. There exist a C3=O3··· π interaction (Table 1), as well.

S2. Experimental


A mixture of 2-chloromethyl-5-hydroxy-4-pyrone (1.0 g, 0.6 mmol) and zinc dust (0.8 g, 12 mmol) in water (20 ml) was stirred for 30 min at 323 K. Concentrated HCl (6 ml) was added dropwise to dissolve the zinc dust completely and the mixture was stirred for 3 h at 353 K. The reaction mixture was transferred to ice–water and extracted with dichloromethane, dried with anhydrous Na₂SO₄ and evaporated to dryness. The crystals of the title compound were obtained by recrystallizing the crude product in isopropanol.

S3. Refinement

The coordinates of H atom of hydroxy group were refined. H atoms were positioned geometrically, with C—H = 0.93 and 0.96 Å for aromatic and methyl H, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C}, \text{O})$, where $x = 1.5$ for methyl H, and $x = 1.2$ for other H atoms.

**Figure 1**

ORTEP drawing of the title compound, with the atom numbering scheme. The thermal ellipsoids are drawn at the 30% probability level. H atoms are shown by small circles of arbitrary radii.

**Figure 2**

The packing figure (*PLATON*: Spek, 2003) which shows that the title compound is dimersed and dimers are connected through H-bonds in helical way.

(I)*Crystal data* $C_6H_6O_3$ $M_r = 126.11$ Triclinic, $P\bar{1}$ Hall symbol: $-P\ 1$ $a = 5.4467\ (4)\ \text{\AA}$ $b = 7.3301\ (5)\ \text{\AA}$ $c = 7.6945\ (5)\ \text{\AA}$ $\alpha = 105.354\ (3)^\circ$ $\beta = 98.416\ (4)^\circ$ $\gamma = 100.008\ (4)^\circ$ $V = 285.68\ (4)\ \text{\AA}^3$ $Z = 2$ $F(000) = 132$ $D_x = 1.466\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 1504 reflections
 $\theta = 2.8\text{--}29.1^\circ$
 $\mu = 0.12 \text{ mm}^{-1}$

$T = 296 \text{ K}$
 Prismatic, colourless
 $0.22 \times 0.20 \times 0.10 \text{ mm}$

Data collection

Bruker Kappa APEXII CCD
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 Detector resolution: $7.40 \text{ pixels mm}^{-1}$
 ω scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 2005)
 $T_{\min} = 0.970$, $T_{\max} = 0.986$

6426 measured reflections
 1504 independent reflections
 713 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.045$
 $\theta_{\max} = 29.1^\circ$, $\theta_{\min} = 2.8^\circ$
 $h = -7 \rightarrow 7$
 $k = -9 \rightarrow 9$
 $l = -10 \rightarrow 10$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.131$
 $S = 1.00$
 1504 reflections
 87 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0523P)^2 + 0.0219P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.18 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.20 \text{ e \AA}^{-3}$
 Extinction correction: SHELXL97 (Sheldrick,
 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.053 (14)

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.2557 (3)	0.3634 (2)	0.37275 (16)	0.0542 (5)
O2	-0.1534 (3)	0.1916 (2)	0.64758 (18)	0.0641 (6)
O3	-0.4440 (3)	0.0215 (2)	0.29282 (18)	0.0631 (6)
C1	0.1595 (4)	0.3242 (3)	0.5153 (3)	0.0554 (8)
C2	-0.0705 (4)	0.2171 (3)	0.4961 (3)	0.0467 (7)
C3	-0.2315 (4)	0.1292 (3)	0.3153 (3)	0.0455 (7)
C4	-0.1223 (4)	0.1770 (3)	0.1711 (3)	0.0486 (7)
C5	0.1096 (4)	0.2896 (3)	0.2016 (3)	0.0463 (7)
C6	0.2424 (4)	0.3492 (4)	0.0642 (3)	0.0624 (8)
H1	0.26084	0.37532	0.63280	0.0664*
H2	-0.299 (5)	0.111 (4)	0.628 (3)	0.0769*

H4	-0.21633	0.12780	0.05123	0.0584*
H6A	0.13508	0.29403	-0.05590	0.0936*
H6B	0.28060	0.48817	0.09427	0.0936*
H6C	0.39779	0.30418	0.06523	0.0936*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0501 (9)	0.0655 (10)	0.0377 (8)	-0.0005 (7)	-0.0002 (6)	0.0131 (7)
O2	0.0703 (11)	0.0723 (12)	0.0356 (8)	-0.0115 (9)	0.0030 (7)	0.0143 (8)
O3	0.0506 (10)	0.0853 (12)	0.0440 (8)	-0.0041 (9)	0.0051 (7)	0.0182 (8)
C1	0.0612 (15)	0.0605 (15)	0.0346 (10)	0.0001 (12)	-0.0004 (10)	0.0117 (10)
C2	0.0540 (14)	0.0473 (13)	0.0343 (10)	0.0071 (11)	0.0035 (9)	0.0101 (9)
C3	0.0410 (13)	0.0519 (14)	0.0391 (11)	0.0078 (11)	0.0038 (9)	0.0099 (9)
C4	0.0457 (13)	0.0612 (15)	0.0327 (9)	0.0077 (11)	0.0027 (9)	0.0089 (10)
C5	0.0454 (13)	0.0544 (14)	0.0343 (10)	0.0090 (11)	0.0033 (9)	0.0091 (9)
C6	0.0547 (14)	0.0819 (17)	0.0488 (12)	0.0074 (12)	0.0110 (10)	0.0210 (12)

Geometric parameters (Å, °)

O1—C1	1.358 (3)	C4—C5	1.334 (3)
O1—C5	1.352 (3)	C5—C6	1.480 (3)
O2—C2	1.356 (3)	C1—H1	0.9300
O3—C3	1.243 (3)	C4—H4	0.9300
O2—H2	0.87 (3)	C6—H6A	0.9600
C1—C2	1.323 (3)	C6—H6B	0.9600
C2—C3	1.446 (3)	C6—H6C	0.9600
C3—C4	1.426 (3)		
O1...O3 ⁱ	3.200 (2)	C2...O1 ⁱⁱⁱ	3.350 (3)
O1...O1 ⁱⁱ	3.078 (2)	C2...O2 ^{vi}	3.405 (3)
O1...C2 ⁱⁱⁱ	3.350 (3)	C2...C1 ⁱⁱⁱ	3.501 (3)
O2...O3	2.7853 (19)	C2...C2 ^{vi}	3.415 (3)
O2...C6 ^{iv}	3.378 (3)	C6...O2 ^{ix}	3.378 (3)
O2...O3 ^v	2.635 (2)	C3...H2 ^v	3.00 (3)
O2...C2 ^{vi}	3.405 (3)	C4...H6C ^{vii}	3.0000
O3...O2 ^v	2.635 (2)	H2...O3	2.46 (2)
O3...O1 ^{vii}	3.200 (2)	H2...O3 ^v	1.83 (3)
O3...O2	2.7853 (19)	H2...C3 ^v	3.00 (3)
O2...H6B ⁱⁱⁱ	2.9000	H4...H6A	2.4500
O2...H6A ^{iv}	2.4200	H4...O3 ^{viii}	2.8200
O3...H2	2.46 (2)	H6A...O2 ^{ix}	2.4200
O3...H2 ^v	1.83 (3)	H6A...H4	2.4500
O3...H4 ^{viii}	2.8200	H6B...O2 ⁱⁱⁱ	2.9000
C1...C1 ⁱⁱⁱ	3.387 (3)	H6C...C4 ⁱ	3.0000
C1...C2 ⁱⁱⁱ	3.501 (3)		
C1—O1—C5	118.57 (18)	O1—C5—C4	121.3 (2)

C2—O2—H2	116.2 (15)	O1—C1—H1	118.00
O1—C1—C2	123.6 (2)	C2—C1—H1	118.00
O2—C2—C3	120.55 (19)	C3—C4—H4	119.00
C1—C2—C3	120.2 (2)	C5—C4—H4	119.00
O2—C2—C1	119.3 (2)	C5—C6—H6A	109.00
O3—C3—C4	124.7 (2)	C5—C6—H6B	109.00
C2—C3—C4	113.9 (2)	C5—C6—H6C	109.00
O3—C3—C2	121.4 (2)	H6A—C6—H6B	109.00
C3—C4—C5	122.5 (2)	H6A—C6—H6C	109.00
O1—C5—C6	111.28 (19)	H6B—C6—H6C	109.00
C4—C5—C6	127.5 (2)		
C5—O1—C1—C2	0.0 (3)	C1—C2—C3—O3	177.0 (2)
C1—O1—C5—C4	-1.4 (3)	C1—C2—C3—C4	-2.6 (3)
C1—O1—C5—C6	179.1 (2)	O3—C3—C4—C5	-178.3 (2)
O1—C1—C2—O2	-177.69 (19)	C2—C3—C4—C5	1.3 (3)
O1—C1—C2—C3	2.0 (4)	C3—C4—C5—O1	0.7 (4)
O2—C2—C3—O3	-3.3 (3)	C3—C4—C5—C6	-179.9 (2)
O2—C2—C3—C4	177.2 (2)		

Symmetry codes: (i) $x+1, y, z$; (ii) $-x+1, -y+1, -z+1$; (iii) $-x, -y+1, -z+1$; (iv) $x, y, z+1$; (v) $-x-1, -y, -z+1$; (vi) $-x, -y, -z+1$; (vii) $x-1, y, z$; (viii) $-x-1, -y, -z$; (ix) $x, y, z-1$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O2—H2...O3	0.87 (3)	2.46 (2)	2.7853 (19)	103.1 (18)
O2—H2...O3 ^v	0.87 (3)	1.83 (3)	2.635 (2)	152 (2)
C6—H6A...O2 ^{ix}	0.9600	2.4200	3.378 (3)	173.00
C3—O3...CgA ^{vii}	1.24 (1)	3.65 (1)	4.363 (2)	118 (1)

Symmetry codes: (v) $-x-1, -y, -z+1$; (vii) $x-1, y, z$; (ix) $x, y, z-1$.