

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Methyl 3-carboxy-5-nitrobenzoate

 Pei Zou,^{a‡} Min-Hao Xie,^b Shi-Neng Luo,^b Ya-Ling Liu^b and Yong-Jia Shen^{a*}
^aInstitute of Fine Chemicals, East China University of Science and Technology, Shanghai 200237, People's Republic of China, and ^bJiangsu Institute of Nuclear Medicine, Wuxi 214063, People's Republic of China

Correspondence e-mail: zou-pei@163.com


Received 16 September 2008; accepted 9 January 2009

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.039; wR factor = 0.105; data-to-parameter ratio = 11.4.

The structure of the title compound, $\text{C}_9\text{H}_7\text{NO}_6$, is essentially planar [maximum deviation 0.284 (2) Å] except for the methyl H atoms. The crystal structure is stabilized by asymmetric O—H···O hydrogen bonds linking the hydrogen carboxylates into pairs around the inversion centres. There is also π – π stacking of the benzene rings [centroid–centroid distance 3.6912 (12) Å].

Related literature

The title complex is as an important intermediate for the preparation of iodinated X-ray contrast media, see: Morin *et al.* (1987); Singh & Rathore (1980); Stacul (2001); Jin & Xiao (2005).



Experimental

Crystal data

 $\text{C}_9\text{H}_7\text{NO}_6$
 $M_r = 225.16$

 Monoclinic, $P2_1/c$
 $a = 7.3450$ (15) Å
 $b = 8.9050$ (18) Å
 $c = 14.474$ (3) Å
 $\beta = 91.18$ (3)°
 $V = 946.5$ (3) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.14$ mm⁻¹
 $T = 293$ (2) K
 $0.30 \times 0.20 \times 0.10$ mm

Data collection

 Enraf–Nonius CAD-4 diffractometer
 Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\min} = 0.950$, $T_{\max} = 0.977$
 1859 measured reflections

 1717 independent reflections
 1284 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$
 3 standard reflections every 200 reflections
 intensity decay: 1.0%

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.105$
 $S = 1.03$
 1717 reflections
 150 parameters

 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.17$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.13$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O6}-\text{H6B}\cdots\text{O5}^i$	0.95 (3)	1.67 (3)	2.6206 (19)	177.9 (17)
$\text{C8}-\text{H8A}\cdots\text{O2}^{ii}$	0.93	2.48	3.406 (2)	174

 Symmetry codes: (i) $-x + 1, -y, -z + 1$; (ii) $-x + 3, -y + 1, -z + 1$.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors acknowledge financial support from the Jiangsu Institute of Nuclear Medicine.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FB2119).

References

- Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
 Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
 Jin, L.-F. & Xiao, F.-P. (2005). *Acta Cryst.* **E61**, o1276–o1277.
 Morin, J. P., Boutelet, I., Toutain, H. & Fillastre, J. P. (1987). *Pathol. Biol.* **35**, 1215–1220.
 North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Singh, G. B. & Rathore, H. G. S. (1980). *Indian Drug. Pharm. Ind.* **15**, 35–38.
 Stacul, F. (2001). *Eur. Radiol.* **11**, 690–697.

[‡] Permanent address: Jiangsu Institute of Nuclear Medicine, Wuxi 214063, People's Republic of China.

supplementary materials

Acta Cryst. (2009). E65, o335 [doi:10.1107/S1600536809001093]

Methyl 3-carboxy-5-nitrobenzoate

P. Zou, M.-H. Xie, S.-N. Luo, Y.-L. Liu and Y.-J. Shen

Comment

The molecule of the title complex (Fig.1) is useful as an important intermediate for the preparation of iodinated X-ray contrast media, such as iotalamic acid, ioxitalamic acid, and Ioxilan, which are used clinically all over the world (Morin *et al.*, 1987; Singh *et al.*, 1980; Stacul *et al.*, 2001). We report here the crystal structure of title compound. The crystal data show that the bond lengths and angles are within expected ranges. The molecule is essentially planar: the maximum deviation from the weighted least-squares plane calculated through all the non-H atoms is 0.284 (2)Å for O2. The molecules are stacked *via* π - π interactions, with the centroid-centroid distance of 3.6912 (12)Å [symmetry code(i): 2-x, 1-y, 1-z]. The stacked columns are linked together by two intermolecular hydrogen bonds, O—H \cdots O and C—H \cdots O (Tab. 1 and Fig. 2). The O—H \cdots O hydrogen bonds bind the hydrogencarboxylates into pairs.

Experimental

Dimethyl 5-nitroisophthalic acid (956 mg, 4 mmol) was dissolved in hot methanol (6 ml), then sodium hydroxide (152 mg, 3.8 mmol) in methanol (2 ml) was added and refluxed for 30 min. Methanol was distilled off. The solid residue was extracted by warm water and the undissolved diester was filtered off. The filtrate was acidified with 1 mol/l hydrochloric acid (4 ml). The precipitate was filtered and washed with cold water. The crude product was purified by recrystallization. Single crystals were grown by slow evaporation of a ethanol/water (*v/v* 1:1) solution: colourless block-shaped crystals were formed after several days.

Refinement

All the H atoms could have been discerned in the difference electron density maps. With exception of the hydrogen belonging to the hydroxyl group of the hydrogencarboxylate the hydrogens were situated into the idealized positions and refined in riding motion approximation. The hydroxyl hydrogen was refined freely. The used constraints: $C_{\text{aryl}}\text{—H} = 0.93 \text{ \AA}$, $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(C_{\text{aryl}})$; $C_{\text{methyl}}\text{—H} = 0.96 \text{ \AA}$, $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{methyl})$.

Figures




Fig. 1. A view of the title compound with the atomic numbering scheme. Displacement ellipsoids were drawn at the 30% probability level.




Fig. 2. A packing diagram viewed along the *b* axis.

Methyl 3-carboxy-5-nitrobenzoate

Crystal data

$C_9H_7NO_6$

$M_r = 225.16$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 7.3450$ (15) Å

$b = 8.9050$ (18) Å

$c = 14.474$ (3) Å

$\beta = 91.18$ (3)°

$V = 946.5$ (3) Å³

$Z = 4$

$F_{000} = 464$

$D_x = 1.580$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 25 reflections

$\theta = 10\text{--}13^\circ$

$\mu = 0.14$ mm⁻¹

$T = 293$ (2) K

Block, colourless

$0.30 \times 0.20 \times 0.10$ mm

Data collection

Enraf-Nonius CAD-4
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293$ (2) K

$\omega/2\theta$ scans

Absorption correction: ψ scan
(North *et al.*, 1968)

$T_{\min} = 0.950$, $T_{\max} = 0.977$

1859 measured reflections

1717 independent reflections

1284 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.021$

$\theta_{\max} = 25.3^\circ$

$\theta_{\min} = 2.7^\circ$

$h = 0 \rightarrow 8$

$k = 0 \rightarrow 10$

$l = -17 \rightarrow 17$

3 standard reflections

every 200 reflections

intensity decay: 1.0%

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.039$

$wR(F^2) = 0.105$

$S = 1.03$

1717 reflections

Secondary atom site location: difference Fourier map

Hydrogen site location: difference Fourier map

H atoms treated by a mixture of
independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.057P)^2 + 0.0354P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.17$ e Å⁻³

150 parameters $\Delta\rho_{\min} = -0.13 \text{ e } \text{\AA}^{-3}$
 23 constraints Extinction correction: none
 Primary atom site location: structure-invariant direct methods

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N	1.0387 (2)	0.49976 (17)	0.29780 (10)	0.0475 (4)
O1	1.30752 (17)	0.20120 (16)	0.64959 (9)	0.0588 (4)
C1	1.4451 (3)	0.2024 (3)	0.72271 (15)	0.0742 (7)
H1A	1.4327	0.1142	0.7602	0.111*
H1B	1.4301	0.2902	0.7603	0.111*
H1C	1.5638	0.2036	0.6962	0.111*
C2	1.3171 (2)	0.3110 (2)	0.58767 (12)	0.0442 (4)
O2	1.4321 (2)	0.40512 (19)	0.58894 (10)	0.0769 (5)
C3	1.1649 (2)	0.30531 (18)	0.51817 (11)	0.0375 (4)
O3	1.1685 (2)	0.58281 (18)	0.29145 (11)	0.0730 (5)
O4	0.9140 (2)	0.4969 (2)	0.24169 (11)	0.0832 (6)
C4	1.0120 (2)	0.21563 (18)	0.53024 (11)	0.0382 (4)
H4A	1.0053	0.1533	0.5816	0.046*
O5	0.70372 (17)	0.03897 (15)	0.54893 (9)	0.0561 (4)
C5	0.8694 (2)	0.21906 (18)	0.46585 (12)	0.0373 (4)
O6	0.57474 (18)	0.13777 (16)	0.42237 (9)	0.0560 (4)
H6B	0.475 (4)	0.072 (3)	0.4318 (19)	0.119 (10)*
C6	0.8793 (2)	0.31082 (18)	0.38850 (11)	0.0380 (4)
H6A	0.7850	0.3131	0.3447	0.046*
C7	1.0324 (2)	0.39840 (18)	0.37824 (11)	0.0368 (4)
C8	1.1757 (2)	0.39837 (18)	0.44108 (11)	0.0379 (4)
H8A	1.2771	0.4589	0.4323	0.046*
C9	0.7057 (2)	0.12519 (19)	0.48049 (12)	0.0396 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N	0.0515 (9)	0.0492 (9)	0.0419 (9)	-0.0049 (8)	-0.0003 (7)	0.0054 (7)

supplementary materials

O1	0.0542 (8)	0.0634 (9)	0.0579 (8)	-0.0185 (7)	-0.0205 (6)	0.0167 (7)
C1	0.0644 (13)	0.1017 (19)	0.0555 (13)	-0.0228 (13)	-0.0249 (11)	0.0225 (13)
C2	0.0414 (9)	0.0467 (11)	0.0443 (10)	-0.0094 (9)	-0.0046 (8)	0.0030 (8)
O2	0.0626 (9)	0.0911 (12)	0.0759 (11)	-0.0421 (9)	-0.0269 (8)	0.0280 (9)
C3	0.0377 (9)	0.0344 (9)	0.0404 (9)	-0.0030 (7)	-0.0009 (7)	-0.0046 (7)
O3	0.0685 (10)	0.0765 (10)	0.0737 (10)	-0.0267 (8)	-0.0043 (8)	0.0321 (8)
O4	0.0783 (11)	0.1067 (14)	0.0635 (10)	-0.0288 (10)	-0.0274 (8)	0.0353 (9)
C4	0.0408 (9)	0.0331 (9)	0.0407 (9)	-0.0040 (7)	-0.0017 (7)	-0.0005 (7)
O5	0.0499 (8)	0.0578 (8)	0.0601 (8)	-0.0174 (6)	-0.0112 (6)	0.0174 (7)
C5	0.0369 (8)	0.0324 (9)	0.0424 (9)	-0.0027 (7)	-0.0020 (7)	-0.0053 (7)
O6	0.0454 (8)	0.0569 (9)	0.0650 (9)	-0.0201 (7)	-0.0184 (7)	0.0123 (7)
C6	0.0396 (9)	0.0369 (9)	0.0373 (9)	-0.0014 (8)	-0.0049 (7)	-0.0040 (7)
C7	0.0409 (9)	0.0355 (9)	0.0342 (8)	-0.0027 (7)	0.0020 (7)	-0.0005 (7)
C8	0.0332 (8)	0.0370 (9)	0.0436 (10)	-0.0037 (7)	0.0034 (7)	-0.0050 (8)
C9	0.0420 (9)	0.0330 (9)	0.0435 (10)	-0.0055 (8)	-0.0060 (8)	-0.0006 (8)

Geometric parameters (Å, °)

N—O3	1.211 (2)	C4—C5	1.388 (2)
N—O4	1.2122 (19)	C4—H4A	0.9300
N—C7	1.475 (2)	O5—C9	1.254 (2)
O1—C2	1.329 (2)	C5—C6	1.389 (2)
O1—C1	1.448 (2)	C5—C9	1.483 (2)
C1—H1A	0.9600	O6—C9	1.270 (2)
C1—H1B	0.9600	O6—H6B	0.95 (3)
C1—H1C	0.9600	C6—C7	1.379 (2)
C2—O2	1.190 (2)	C6—H6A	0.9300
C2—C3	1.489 (2)	C7—C8	1.377 (2)
C3—C4	1.393 (2)	C8—H8A	0.9300
C3—C8	1.393 (2)		
O3—N—O4	123.18 (16)	C3—C4—H4A	119.9
O3—N—C7	118.11 (15)	C4—C5—C6	120.19 (15)
O4—N—C7	118.71 (15)	C4—C5—C9	119.62 (15)
C2—O1—C1	116.20 (15)	C6—C5—C9	120.19 (15)
O1—C1—H1A	109.5	C9—O6—H6B	115.3 (17)
O1—C1—H1B	109.5	C7—C6—C5	118.42 (15)
H1A—C1—H1B	109.5	C7—C6—H6A	120.8
O1—C1—H1C	109.5	C5—C6—H6A	120.8
H1A—C1—H1C	109.5	C8—C7—C6	122.86 (15)
H1B—C1—H1C	109.5	C8—C7—N	119.05 (15)
O2—C2—O1	123.71 (17)	C6—C7—N	118.05 (15)
O2—C2—C3	123.86 (17)	C7—C8—C3	118.28 (15)
O1—C2—C3	112.41 (15)	C7—C8—H8A	120.9
C4—C3—C8	120.06 (15)	C3—C8—H8A	120.9
C4—C3—C2	122.07 (15)	O5—C9—O6	123.82 (16)
C8—C3—C2	117.81 (15)	O5—C9—C5	118.73 (15)
C5—C4—C3	120.18 (16)	O6—C9—C5	117.45 (15)
C5—C4—H4A	119.9		
C1—O1—C2—O2	1.4 (3)	C5—C6—C7—N	-177.71 (14)

C1—O1—C2—C3	-177.00 (17)	O3—N—C7—C8	-2.1 (2)
O2—C2—C3—C4	-165.32 (18)	O4—N—C7—C8	178.43 (17)
O1—C2—C3—C4	13.1 (2)	O3—N—C7—C6	175.95 (17)
O2—C2—C3—C8	12.0 (3)	O4—N—C7—C6	-3.6 (2)
O1—C2—C3—C8	-169.58 (15)	C6—C7—C8—C3	0.1 (2)
C8—C3—C4—C5	-0.3 (2)	N—C7—C8—C3	178.04 (14)
C2—C3—C4—C5	176.95 (16)	C4—C3—C8—C7	-0.1 (2)
C3—C4—C5—C6	0.7 (2)	C2—C3—C8—C7	-177.46 (15)
C3—C4—C5—C9	-178.81 (15)	C4—C5—C9—O5	-3.4 (2)
C4—C5—C6—C7	-0.6 (2)	C6—C5—C9—O5	177.14 (16)
C9—C5—C6—C7	178.84 (15)	C4—C5—C9—O6	176.41 (16)
C5—C6—C7—C8	0.2 (2)	C6—C5—C9—O6	-3.1 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O6—H6B \cdots O5 ⁱ	0.95 (3)	1.67 (3)	2.6206 (19)	177.9 (17)
C8—H8A \cdots O2 ⁱⁱ	0.93	2.48	3.406 (2)	174

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $-x+3, -y+1, -z+1$.

Fig. 1




Fig. 2

