

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

## Poly[bis( $\mu$ -hemihydrogen 2-phenyl-quinoline-4-carboxylato- $\kappa^2N,O$ )silver(I)]. Corrigendum

Xiutang Zhang,<sup>a,b\*</sup> Peihai Wei<sup>a</sup> and Bin Li<sup>c</sup>

<sup>a</sup>Advanced Material Institute of Research, Department of Chemistry and Chemical Engineering, Shandong Institute of Education, Jinan, 250013, People's Republic of China, <sup>b</sup>College of Chemistry and Chemical Engineering, Liaocheng University, Liaocheng, 252059, People's Republic of China, and <sup>c</sup>Department of Chemistry and Chemical Engineering, Shandong Institute of Education, Jinan 250013, People's Republic of China

Correspondence e-mail: xiutangzhang@yahoo.com.cn

Received 6 February 2009; accepted 20 February 2009

The affiliation of one of the authors in the paper by Zhang, Wei & Li [*Acta Cryst.* (2009), E65, m223] is corrected.

In the paper by Zhang, Wei & Li [*Acta Cryst.* (2009), E65, m223], the affiliation of the second author was given incorrectly. The correct address is given here.

# Poly[bis( $\mu$ -hemihydrogen 2-phenylquinoline-4-carboxylato- $\kappa^2$ N,O)silver(I)]

Xiutang Zhang,<sup>a,b,\*</sup> Peihai Wei<sup>b</sup> and Bin Li<sup>c</sup>

<sup>a</sup>Advanced Material Institute of Research, Department of Chemistry and Chemical Engineering, Shandong Institute of Education, Jinan, 250013, People's Republic of China, <sup>b</sup>College of Chemistry and Chemical Engineering, Liaocheng University, Liaocheng, 252059, People's Republic of China, and <sup>c</sup>Department of Chemistry and Chemical Engineering, Shandong Institute of Education, Jinan 250013, People's Republic of China

Correspondence e-mail: xiutangzhang@yahoo.com.cn


Received 21 October 2008; accepted 10 January 2009

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.029;  $wR$  factor = 0.067; data-to-parameter ratio = 15.8.

In the title compound,  $[\text{Ag}(\text{C}_{16}\text{H}_{10.5}\text{NO}_2)_2]$ , the  $\text{Ag}^{\text{I}}$  cation (site symmetry 2) is coordinated by two N atoms in a near-linear  $\text{AgN}_2$  arrangement. Two carboxylate O atoms from two additional 2-phenylquinoline-4-carboxylate ligands form long  $\text{Ag}-\text{O}$  bonds [2.6585 (17) Å], resulting in a distorted square-planar arrangement. The bridging ligands result in infinite corrugated sheets propagating in (010). An  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bond, disordered about a twofold axis, completes the structure.

## Related literature

For the related coordination polymers containing  $\text{Mn}^{\text{II}}$ ,  $\text{Co}^{\text{II}}$  and  $\text{Cu}^{\text{II}}$ , see: Xiao *et al.* (2005); Xie *et al.* (2005) and Xie *et al.* (2006), respectively.



## Experimental

### Crystal data

$[\text{Ag}(\text{C}_{16}\text{H}_{10.5}\text{NO}_2)_2]$   
 $M_r = 605.38$   
 Monoclinic,  $C2/c$   
 $a = 7.2163$  (6) Å  
 $b = 20.5060$  (17) Å  
 $c = 16.5632$  (12) Å  
 $\beta = 97.585$  (3)°

$V = 2429.5$  (3) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.87$  mm<sup>-1</sup>  
 $T = 293$  (2) K  
 $0.20 \times 0.15 \times 0.15$  mm

### Data collection

Bruker SMART CCD diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 2001)  
 $T_{\text{min}} = 0.877$ ,  $T_{\text{max}} = 1.00$   
 (expected range = 0.769–0.877)

9257 measured reflections  
 2791 independent reflections  
 2550 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.020$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.029$   
 $wR(F^2) = 0.067$   
 $S = 1.10$   
 2791 reflections

177 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.66$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.33$  e Å<sup>-3</sup>

**Table 1**

Selected geometric parameters (Å, °).

Ag1–N1	2.2413 (15)
N1–Ag1–N1 <sup>i</sup>	177.38 (8)
Symmetry code: (i) $-x + 1, y, -z - \frac{1}{2}$ .	

**Table 2**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1–H1O <sup>ii</sup> ⋯O1 <sup>ii</sup>	0.82	1.68	2.470 (3)	162
Symmetry code: (ii) $-x + 1, y, -z + \frac{1}{2}$ .				

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2825).

## References

- Bruker (2001). *SMART, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Xiao, H. P., Li, X.-H. & Cheng, Y.-Q. (2005). *Acta Cryst.* **E61**, m158–m159.  
 Xie, G., Zeng, M.-H., Chen, S.-P. & Gao, S.-L. (2005). *Acta Cryst.* **E61**, m2273.  
 Xie, G., Zeng, M.-H., Chen, S.-P. & Gao, S.-L. (2006). *Acta Cryst.* **E62**, m397–m399.

**supplementary materials**

*Acta Cryst.* (2009). E65, m223 [ doi:10.1107/S1600536809001184 ]

## Poly[bis( $\mu$ -hemihydrogen 2-phenylquinoline-4-carboxylato- $\kappa^2N,O$ )silver(I)]

X. Zhang, P. Wei and B. Li

### Experimental

A mixture of silver nitrate (0.5 mmol) and 2-phenyl-4-quinolinecarboxylic acid (0.5 mmol) in H<sub>2</sub>O (8 ml) and ethanol (8 ml) was sealed in a 25 ml Teflon-lined stainless steel autoclave and kept at 413 K for three days. Orange prisms of (I) were obtained after cooling to room temperature with a yield of 27%. Anal. Calc. for C<sub>32</sub>H<sub>21</sub>AgN<sub>2</sub>O<sub>4</sub>: C 63.43, H 3.47, N 4.63%; Found: C 63.41, H 3.32, N 4.59%.

### Refinement

The H atoms were placed in calculated positions (C—H = 0.93 Å, O—H = 0.82 Å) and refined as riding with and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{carrier})$ .

### Figures




Fig. 1. A view of the structure of (I), expanded to show the Ag—N bonds, showing 30% probability displacement ellipsoids. Symmetry code: I: 1-x, y, -z-1/2.




Fig. 2. Part of a polymeric sheet in (I).

## Poly[bis( $\mu$ -hemihydrogen 2-phenylquinoline-4-carboxylato- $\kappa^2$ N,O)silver(I)]

### Crystal data

[Ag(C<sub>16</sub>H<sub>10.5</sub>NO<sub>2</sub>)<sub>2</sub>]

$M_r = 605.38$

Monoclinic,  $C2/c$

Hall symbol: -C 2yc

$a = 7.2163$  (6) Å

$b = 20.5060$  (17) Å

$c = 16.5632$  (12) Å

$\beta = 97.585$  (3)°

$V = 2429.5$  (3) Å<sup>3</sup>

$Z = 4$

$F_{000} = 1224$

$D_x = 1.655$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation

$\lambda = 0.71073$  Å

Cell parameters from 3290 reflections

$\theta = 2.3$ – $27.5$ °

$\mu = 0.87$  mm<sup>-1</sup>

$T = 293$  (2) K

Prism, orange

$0.20 \times 0.15 \times 0.15$  mm

### Data collection

Bruker SMART CCD  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293$ (2) K

$\omega$  scans

Absorption correction: Multi-scan  
(SADABS; Bruker, 2001)

$T_{\min} = 0.877$ ,  $T_{\max} = 1.00$

9257 measured reflections

2791 independent reflections

2550 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.020$

$\theta_{\max} = 27.5$ °

$\theta_{\min} = 2.3$ °

$h = -9 \rightarrow 9$

$k = -25 \rightarrow 26$

$l = -21 \rightarrow 20$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.029$

$wR(F^2) = 0.067$

$S = 1.10$

2791 reflections

177 parameters

Primary atom site location: structure-invariant direct  
methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring  
sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0323P)^2 + 1.551P]$$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.66$  e Å<sup>-3</sup>

$\Delta\rho_{\min} = -0.33$  e Å<sup>-3</sup>

Extinction correction: none

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Ag1	0.5000	0.224313 (10)	-0.2500	0.03976 (9)	
N1	0.4524 (2)	0.22681 (7)	-0.11903 (9)	0.0305 (3)	
O1	0.4807 (3)	0.16931 (8)	0.17478 (8)	0.0506 (4)	
H1O	0.4750	0.1750	0.2234	0.061*	0.50
O2	0.6459 (3)	0.26118 (9)	0.18437 (10)	0.0555 (4)	
C1	0.4841 (3)	0.28414 (9)	-0.07613 (11)	0.0323 (4)	
C2	0.4827 (4)	0.34294 (10)	-0.12074 (13)	0.0468 (5)	
H2A	0.4619	0.3417	-0.1773	0.056*	
C3	0.5113 (4)	0.40127 (11)	-0.08202 (16)	0.0562 (7)	
H3A	0.5094	0.4396	-0.1121	0.067*	
C4	0.5437 (4)	0.40349 (11)	0.00334 (16)	0.0546 (6)	
H4A	0.5627	0.4436	0.0294	0.065*	
C5	0.5477 (3)	0.34808 (11)	0.04857 (13)	0.0446 (5)	
H5A	0.5712	0.3506	0.1051	0.054*	
C6	0.5165 (3)	0.28627 (9)	0.01036 (11)	0.0317 (4)	
C7	0.5077 (3)	0.22584 (9)	0.05293 (11)	0.0313 (4)	
C8	0.4621 (3)	0.17105 (10)	0.00812 (11)	0.0344 (4)	
H8A	0.4467	0.1319	0.0348	0.041*	
C9	0.4376 (3)	0.17227 (9)	-0.07805 (10)	0.0302 (4)	
C10	0.3928 (3)	0.11103 (9)	-0.12467 (11)	0.0331 (4)	
C11	0.2740 (3)	0.11181 (10)	-0.19829 (12)	0.0379 (4)	
H11A	0.2219	0.1510	-0.2185	0.045*	
C12	0.2330 (3)	0.05444 (12)	-0.24147 (13)	0.0476 (5)	
H12A	0.1558	0.0555	-0.2911	0.057*	
C13	0.3063 (4)	-0.00390 (12)	-0.21119 (15)	0.0570 (7)	
H13A	0.2773	-0.0424	-0.2399	0.068*	
C14	0.4230 (4)	-0.00520 (11)	-0.13802 (15)	0.0593 (7)	
H14A	0.4731	-0.0446	-0.1177	0.071*	
C15	0.4659 (4)	0.05159 (11)	-0.09489 (12)	0.0463 (5)	
H15A	0.5442	0.0502	-0.0456	0.056*	
C16	0.5497 (3)	0.22006 (10)	0.14510 (11)	0.0356 (4)	

## supplementary materials

---

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Ag1	0.05874 (17)	0.04506 (14)	0.01656 (11)	0.000	0.00894 (9)	0.000
N1	0.0388 (9)	0.0352 (8)	0.0176 (7)	0.0017 (7)	0.0040 (6)	0.0002 (6)
O1	0.0877 (12)	0.0478 (8)	0.0152 (6)	-0.0092 (8)	0.0029 (7)	-0.0008 (6)
O2	0.0607 (11)	0.0753 (11)	0.0275 (8)	-0.0222 (9)	-0.0048 (7)	-0.0068 (7)
C1	0.0384 (10)	0.0364 (10)	0.0232 (9)	0.0005 (8)	0.0081 (8)	-0.0004 (7)
C2	0.0718 (16)	0.0412 (11)	0.0289 (10)	0.0025 (11)	0.0122 (10)	0.0043 (8)
C3	0.085 (2)	0.0358 (11)	0.0494 (14)	-0.0001 (11)	0.0162 (13)	0.0038 (10)
C4	0.0766 (18)	0.0373 (11)	0.0498 (14)	-0.0021 (11)	0.0084 (12)	-0.0110 (10)
C5	0.0549 (14)	0.0462 (12)	0.0330 (11)	0.0004 (10)	0.0066 (10)	-0.0110 (9)
C6	0.0331 (10)	0.0391 (10)	0.0236 (9)	0.0001 (8)	0.0060 (7)	-0.0047 (7)
C7	0.0313 (10)	0.0447 (10)	0.0179 (8)	0.0001 (8)	0.0036 (7)	-0.0016 (7)
C8	0.0485 (12)	0.0369 (9)	0.0179 (8)	-0.0024 (9)	0.0046 (8)	0.0019 (7)
C9	0.0364 (10)	0.0368 (9)	0.0176 (8)	0.0005 (8)	0.0039 (7)	0.0010 (7)
C10	0.0457 (12)	0.0347 (9)	0.0194 (8)	-0.0039 (8)	0.0066 (8)	0.0005 (7)
C11	0.0447 (12)	0.0434 (11)	0.0252 (9)	-0.0031 (9)	0.0035 (8)	0.0000 (8)
C12	0.0557 (14)	0.0549 (13)	0.0309 (10)	-0.0146 (11)	0.0007 (10)	-0.0074 (9)
C13	0.088 (2)	0.0440 (12)	0.0401 (12)	-0.0171 (12)	0.0113 (12)	-0.0106 (10)
C14	0.101 (2)	0.0359 (11)	0.0414 (13)	0.0018 (12)	0.0110 (13)	0.0044 (9)
C15	0.0723 (16)	0.0412 (11)	0.0243 (10)	0.0009 (10)	0.0023 (10)	0.0027 (8)
C16	0.0385 (11)	0.0488 (11)	0.0193 (8)	0.0033 (9)	0.0031 (8)	-0.0040 (8)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Ag1—N1	2.2413 (15)	C5—H5A	0.9300
Ag1—N1 <sup>i</sup>	2.2413 (15)	C6—C7	1.431 (3)
Ag1—O2 <sup>ii</sup>	2.6585 (17)	C7—C8	1.363 (3)
Ag1—O2 <sup>iii</sup>	2.6585 (17)	C7—C16	1.521 (2)
N1—C9	1.320 (2)	C8—C9	1.415 (2)
N1—C1	1.377 (2)	C8—H8A	0.9300
O1—C16	1.280 (2)	C9—C10	1.487 (3)
O1—H1O	0.8200	C10—C15	1.392 (3)
O2—C16	1.223 (3)	C10—C11	1.395 (3)
O2—Ag1 <sup>iii</sup>	2.6585 (17)	C11—C12	1.388 (3)
C1—C2	1.414 (3)	C11—H11A	0.9300
C1—C6	1.421 (3)	C12—C13	1.376 (4)
C2—C3	1.360 (3)	C12—H12A	0.9300
C2—H2A	0.9300	C13—C14	1.382 (4)
C3—C4	1.403 (4)	C13—H13A	0.9300
C3—H3A	0.9300	C14—C15	1.380 (3)
C4—C5	1.359 (3)	C14—H14A	0.9300
C4—H4A	0.9300	C15—H15A	0.9300
C5—C6	1.421 (3)		
N1—Ag1—N1 <sup>i</sup>	177.38 (8)	C8—C7—C16	118.96 (17)

N1—Ag1—O2 <sup>ii</sup>	97.54 (6)	C6—C7—C16	123.08 (16)
N1 <sup>i</sup> —Ag1—O2 <sup>ii</sup>	82.16 (6)	C7—C8—C9	121.59 (18)
N1—Ag1—O2 <sup>iii</sup>	82.16 (6)	C7—C8—H8A	119.2
N1 <sup>i</sup> —Ag1—O2 <sup>iii</sup>	97.54 (6)	C9—C8—H8A	119.2
O2 <sup>ii</sup> —Ag1—O2 <sup>iii</sup>	167.15 (8)	N1—C9—C8	121.72 (17)
C9—N1—C1	118.51 (16)	N1—C9—C10	118.39 (15)
C9—N1—Ag1	120.76 (12)	C8—C9—C10	119.90 (16)
C1—N1—Ag1	118.86 (11)	C15—C10—C11	118.69 (18)
C16—O1—H1O	109.5	C15—C10—C9	120.59 (18)
C16—O2—Ag1 <sup>iii</sup>	137.88 (16)	C11—C10—C9	120.70 (17)
N1—C1—C2	117.98 (17)	C12—C11—C10	120.3 (2)
N1—C1—C6	122.71 (17)	C12—C11—H11A	119.8
C2—C1—C6	119.30 (17)	C10—C11—H11A	119.8
C3—C2—C1	120.9 (2)	C13—C12—C11	120.3 (2)
C3—C2—H2A	119.6	C13—C12—H12A	119.9
C1—C2—H2A	119.6	C11—C12—H12A	119.9
C2—C3—C4	119.9 (2)	C12—C13—C14	119.8 (2)
C2—C3—H3A	120.0	C12—C13—H13A	120.1
C4—C3—H3A	120.0	C14—C13—H13A	120.1
C5—C4—C3	121.1 (2)	C15—C14—C13	120.5 (2)
C5—C4—H4A	119.4	C15—C14—H14A	119.8
C3—C4—H4A	119.4	C13—C14—H14A	119.8
C4—C5—C6	120.6 (2)	C14—C15—C10	120.5 (2)
C4—C5—H5A	119.7	C14—C15—H15A	119.8
C6—C5—H5A	119.7	C10—C15—H15A	119.8
C5—C6—C1	118.14 (18)	O2—C16—O1	125.35 (19)
C5—C6—C7	124.57 (18)	O2—C16—C7	120.26 (18)
C1—C6—C7	117.24 (16)	O1—C16—C7	114.37 (17)
C8—C7—C6	117.95 (17)		

Symmetry codes: (i)  $-x+1, y, -z-1/2$ ; (ii)  $x-1/2, -y+1/2, z-1/2$ ; (iii)  $-x+3/2, -y+1/2, -z$ .

*Hydrogen-bond geometry* ( $\text{\AA}, ^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1O $\cdots$ O1 <sup>iv</sup>	0.82	1.68	2.470 (3)	162

Symmetry codes: (iv)  $-x+1, y, -z+1/2$ .

Fig. 1




Fig. 2

