

1,3-Bis(2-thienylmethyl)-4,5-dihydroimidazolium trichlorido(η^6 -*p*-cymene)-ruthenate(II)

Hakan Arslan,^{a,b*} Don VanDerveer,^c İsmail Özdemir,^d Nevin Gürbüz,^d Yetkin Gök^d and Bekir Çetinkaya^e

^aDepartment of Natural Sciences, Fayetteville State University, Fayetteville, NC 28301, USA, ^bDepartment of Chemistry, Faculty of Pharmacy, Mersin University, Mersin, TR 33169, Turkey, ^cDepartment of Chemistry, Clemson University, Clemson, SC 29634, USA, ^dDepartment of Chemistry, Faculty of Science and Arts, İnönü University, Malatya, TR 44280, Turkey, and ^eDepartment of Chemistry, Faculty of Science, Ege University, Bornova-İzmir, TR 35100, Turkey
Correspondence e-mail: hakan.arslan.acad@gmail.com


Received 1 January 2009; accepted 6 January 2009

Key indicators: single-crystal X-ray study; $T = 153$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; disorder in main residue; R factor = 0.032; wR factor = 0.084; data-to-parameter ratio = 16.0.

The asymmetric unit of the title compound, $(\text{C}_{13}\text{H}_{15}\text{N}_2\text{S}_2)[\text{RuCl}_3(\text{C}_{10}\text{H}_{14})]$, contains a 1,3-(2-thienylmethyl)-4,5-dihydroimidazolium cation and a trichlorido(η^6 -*p*-cymene)-ruthenate(II) anion. The thiophene rings of the cation are disordered by an 180° rotation about the thiophene-CH₂ bonds with occupancies of 0.847 (5)/0.153 (5) and 0.700 (5)/0.300 (5), respectively. The Ru atom exhibits a distorted octahedral coordination with the benzene ring of the *p*-cymene ligand formally occupying three sites and three Cl atoms occupying the other three sites. The short C-N bond lengths in the imidazoline ring indicate partial electron delocalization within the N-C-N fragment. Cation and anions are connected through five intermolecular C-H...Cl hydrogen bonds and one C-H... π hydrogen bond, forming a three-dimensional hydrogen-bonded network.

Related literature

For the synthesis, see: Yaşar *et al.* (2008). Özdemir *et al.* (2008, 2007, 2005). For general background, see: Herrmann *et al.* (1995); Herrmann (2002); Arduengo & Krafczyk (1998). For related compounds, see: Arslan *et al.* (2007, 2005*a,b*) and references therein; Sonar *et al.* (2004, 2005*a,b*); Wagner *et al.* (2006*a,b*); Crundwell *et al.* (2002); Linehan *et al.* (2003); Liu *et al.* (2004); Navarro *et al.* (2006); Therrien *et al.* (2004). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$(\text{C}_{13}\text{H}_{15}\text{N}_2\text{S}_2)[\text{RuCl}_3(\text{C}_{10}\text{H}_{14})]$
 $M_r = 605.02$
 Triclinic, $P\bar{1}$
 $a = 9.910$ (2) Å
 $b = 11.600$ (2) Å
 $c = 12.659$ (3) Å
 $\alpha = 84.95$ (3)°
 $\beta = 67.05$ (3)°

$\gamma = 74.14$ (3)°
 $V = 1288.8$ (6) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 1.10$ mm⁻¹
 $T = 153$ (2) K
 $0.48 \times 0.19 \times 0.17$ mm

Data collection

Rigaku AFC-8S Mercury CCD diffractometer
 Absorption correction: multi-scan (*REQUAB*; Jacobson, 1998)
 $T_{\min} = 0.621$, $T_{\max} = 0.836$

11095 measured reflections
 4557 independent reflections
 4062 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.084$
 $S = 1.13$
 4557 reflections

285 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.43$ e Å⁻³
 $\Delta\rho_{\min} = -0.68$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C11}-\text{H11}\cdots\text{Cl1}^i$	0.96	2.62	3.450 (4)	144
$\text{C14}-\text{H14A}\cdots\text{Cl1}^i$	0.96	2.82	3.553 (4)	134
$\text{C19}-\text{H19A}\cdots\text{Cl2}^i$	0.96	2.81	3.671 (4)	150
$\text{C23}-\text{H23}\cdots\text{Cl1}^{ii}$	0.96	2.66	3.549 (4)	154
$\text{C14}-\text{H14B}\cdots\text{Cg2}^{iii}$	0.96	2.83	3.784 (5)	171
$\text{C19}-\text{H19B}\cdots\text{Cl1}$	0.96	2.86	3.759 (5)	157

Symmetry codes: (i) $-x, -y + 1, -z + 2$; (ii) $-x, -y + 2, -z + 2$; (iii) $-x + 1, -y + 1, -z + 2$. Cg1 is the centroid of the S2,C20-C23 thiophene ring.

Data collection: *CrystalClear* (Rigaku/MSC, 2006); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

We thank the Technological and Scientific Research Council of Turkey TÜBİTAK-CNRS [TBAG-U/181 (106 T716)] and İnönü University Research Fund (BAP: 2008-Güdümlü3) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG2465).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Arduengo, A. J. & Krafczyk, R. (1998). *Chem. Ztg*, **32**, 6–14.
- Arslan, H., VanDerveer, D., Özdemir, İ., Çetinkaya, B. & Demir, S. (2005a). *J. Chem. Crystallogr.* **35**, 491–495.
- Arslan, H., VanDerveer, D., Özdemir, İ., Yaşar, S. & Çetinkaya, B. (2005b). *Acta Cryst.* **E61**, m1873–m1875.
- Arslan, H., VanDerveer, D., Yaşar, S., Özdemir, İ. & Çetinkaya, B. (2007). *Acta Cryst.* **E63**, m1001–m1003.
- Crundwell, G., Meskill, T., Sayers, D. & Kantardjieff, K. (2002). *Acta Cryst.* **E58**, o668–o670.
- Herrmann, W. A. (2002). *Angew. Chem. Int. Ed.* **41**, 1290–1309.
- Herrmann, W. A., Elison, M., Fischer, J., Köcher, C. & Artus, G. R. J. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 2371–2374.
- Jacobson, R. (1998). *REQAB*. Molecular Structure Corporation, The Woodlands, Texas, USA.
- Linehan, J., Crundwell, G., Herron, S. R. & Kantardjieff, K. A. (2003). *Acta Cryst.* **E59**, o466–o468.
- Liu, L., Zhang, Q.-F. & Leung, W.-H. (2004). *Acta Cryst.* **E60**, m506–m508.
- Navarro, O., Marion, N., Oonishi, Y., Kelly, R. A. & Nolan, S. P. (2006). *J. Org. Chem.* **71**, 685–692.
- Özdemir, İ., Demir, S., Çetinkaya, B. & Çetinkaya, E. (2005). *J. Organomet. Chem.* **690**, 5849–5855.
- Özdemir, İ., Demir, S., Çetinkaya, B., Toupet, L., Castarlanes, R., Fischmeister, C. & Dixneuf, P. H. (2007). *Eur. J. Inorg. Chem.* **18**, 2862–2869.
- Özdemir, İ., Gürbüz, N., Gök, Y. & Çetinkaya, B. (2008). *Heteroat. Chem.* **19**, 82–86.
- Rigaku/MS (2006). *CrystalClear*. Rigaku/MS, The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sonar, V. N., Parkin, S. & Crooks, P. A. (2004). *Acta Cryst.* **C60**, o217–o218.
- Sonar, V. N., Parkin, S. & Crooks, P. A. (2005a). *Acta Cryst.* **E61**, o933–o935.
- Sonar, V. N., Parkin, S. & Crooks, P. A. (2005b). *Acta Cryst.* **C61**, o78–o80.
- Therrien, B., Frein, S. & Süß-Fink, G. (2004). *Acta Cryst.* **E60**, m1666–m1668.
- Wagner, P., Officer, D. L. & Kubicki, M. (2006a). *Acta Cryst.* **E62**, o5745–o5747.
- Wagner, P., Officer, D. L. & Kubicki, M. (2006b). *Acta Cryst.* **E62**, o5931–o5932.
- Yaşar, S., Özdemir, İ., Çetinkaya, B., Renaud, J. L. & Bruneau, C. (2008). *Eur. J. Org. Chem.* **12**, 2142–2149.

supplementary materials

Acta Cryst. (2009). E65, m165-m166 [doi:10.1107/S1600536809000464]

1,3-Bis(2-thienylmethyl)-4,5-dihydroimidazolium trichlorido(η^6 -*p*-cymene)ruthenate(II)

H. Arslan, D. VanDerveer, I. Özdemir, N. Gürbüz, Y. Gök and B. Çetinkaya

Comment

Metal-carbene compounds, such as *N*-heterocyclic carbene palladium and ruthenium complexes, are important catalysts that have a wide range of applications such as Suzuki-Miyaura, Sonogashira, Stille and Heck reactions (Herrmann 2002; Herrmann *et al.*, 1995; Navarro *et al.*, 2006; Arduengo & Krafczyk, 1998).

In previous papers, we have described the synthesis, characterization and applications of palladium, platinum and ruthenium *N*-heterocyclic carbene complexes as catalysts (Yaşar *et al.*, 2008; Arslan *et al.*, 2007, 2005*a*, 2005*b*, and references therein; Özdemir *et al.*, 2008, 2007, 2005, and references therein). In view of these important attributes of *N*-heterocyclic carbene derivatives, we report here the crystal structure of one of them. The title compound consists 1,3-*bis*(thiophen-2-ylmethyl)-4,5-dihydro-1*H*-imidazolium cation and a trichloro(η^6 -*p*-cymene) ruthenium(II) anion. The molecular structure of the title compound, (I), is depicted in Fig. 1. Cation and anion groups are connected with five intermolecular C—H \cdots Cl hydrogen bonds and one C—H \cdots π hydrogen bond, forming a three-dimensional hydrogen-bonding network (Fig. 2).

A flip disorder of both thiophene rings in 1,3-*bis*(thiophen-2-ylmethyl)-4,5-dihydro-1*H*-imidazolium cation is observed. There are two positions of both thiophene rings, rotated by 180°. The crystal structure of the cation contains four disordered atoms, S1, S2, C16, and C21. The site occupancy factors refined to 0.847 (5) and 0.153 (5) for the S1—C15—C16—C17—C18 ring, and 0.700 (5) and 0.300 (5) for the S2—C20—C21—C22—C23 ring. A similar thiophene ring disorder has been observed in some thiophene derivatives, such as (*Z*)-3-(1-methyl-1*H*-indol-3-yl)-2-(thiophen-3-yl)acrylonitrile (Sonar *et al.*, 2004), (*Z*)-2-(3-thienyl)-3-(3,4,5-trimethoxyphenyl)acrylonitrile (Sonar *et al.*, 2005*a*), (*Z*)-3-(1*H*-Indol-3-yl)-2-(3-thienyl)acrylonitrile and (*Z*)-3-[1-(4-*tert*-butylbenzyl)-1*H*-indol-3-yl]-2-(3-thienyl)acrylonitrile (Sonar *et al.*, 2005*b*), 1,2-di-3-thienyl-2-hydroxyethanone(3,3-thenoin) (Crundwell *et al.*, 2002), 3-[2-(anthracen-9-yl)ethenyl] thiophene, (Wagner *et al.*, 2006*a*), 2,5-bis(2-cyano-2-thienylvinyl)thiophene (Wagner *et al.*, 2006*b*), and 1,4-diphenyl-2,3-dithien-3-ylcyclopentadien-1-one (Linehan *et al.*, 2003). In addition, all thiophene rings in the cation are almost planar; the maximum deviations from the least squares planes are 0.019 (4) Å for C16 and 0.006 (6) Å for C22.

The coordination geometry of ruthenium is pseudooctahedral, with an average Ru—Cl bond distance of 2.430 Å. The ruthenium atom exhibits a distorted octahedral coordination with the benzene ring of the *p*-cymene ligand formally occupying three sites and three chloride atoms occupying other three sites. The distance between the centroid of the *p*-cymene ring and ruthenium is 1.6493 (15) Å, which is longer than that reported in other ruthenium compounds (Liu *et al.*, 2004; Therrien *et al.*, 2004). All the other bond lengths in (I) are in normal ranges (Allen *et al.*, 1987).

The imidazolidine ring is almost planar, the deviations from planarity of ring are N1 0.002 (3), C11 0.001 (4), N2 0.004 (3), C12 0.005 (4), C13 0.004 (4) Å. The some C—N bond lengths (N1—C11 = 1.307 (4) Å and N2—C11 = 1.302 (4) Å) for the imidazolidine ring are shorter than the average single C—N bond length of 1.48 Å, thus showing double bond character in these C—N bonds. The other CN bonds length (N1—C13 1.458 (5), N1—C14 1.462 (4), N2—C19 1.460 (4) and

supplementary materials

N2—C12 1.466 (4) Å) is agree with 1.48 Å C—N single bond lengths. This information indicates a partial electron delocalization within the N1—C11—N2 fragment.

The crystal packing is shown in Fig. 2. Five intermolecular C—H...Cl hydrogen bonds link the molecules of (I) and generate a three-dimensional hydrogen bonded framework. In addition, a C14 (x, y, z)-H... π (S2—C20—C21—C22—C23, thiophene ring; $1 - x, 1 - y, 2 - z$) hydrogen bond is observed in the title compound, Table 1.

Experimental

A suspension of 1,3-*bis*(thiophen-2ylmethyl)-4,5-dihydro-1*H*-imidazolium chloride (1.1 mmol), Cs₂CO₃ (1.2 mmol) and [RuCl₂(*p*-cymene)] (0.5 mmol) was heated under reflux in degassed toluene (20 ml) for 7 h (Fig. 3). The reaction mixture was then filtered while hot, and the volume was reduced to about 10 ml before addition of *n*-hexane (15 ml). The precipitate formed was crystallized from CH₂Cl₂: hexane (5:10 ml) to give the complex as red-brown crystals. Yields: 0.208 g, 69%. *M.p.*: 227–228 °C. ¹H NMR (CDCl₃) δ : 1.39 (d, 6H, $J = 6.9$ Hz, CH₃(C₆H₄)CH(CH₃)₂), 2.29 (s, 3H, CH₃(C₆H₄)CH(CH₃)₂), 3.21 (m, 1H, CH₃(C₆H₄)CH(CH₃)₂), 3.79 (s, 4H, NCH₂CH₂N), 4.12 (s, 4H, CH₂C₄H₃S), 5.29 and 5.58 (d, 4H, $J = 5.8$ Hz, CH₃(C₆H₄)CH(CH₃)₂), 7.03–7.68 (m, 6H, C₄H₃S), 8.99 (s, 1H, 2-CH). ¹³C NMR (CDCl₃) δ : 18.6 (CH₃(C₆H₄)CH(CH₃)₂), 22.3 (CH₃(C₆H₄)CH(CH₃)₂), 30.8 (CH₃(C₆H₄)CH(CH₃)₂), 47.0 (NCH₂CH₂N), 47.1 (CH₂C₄H₃S), 79.6, 81.8, 96.5 and 100.8 (CH₃(C₆H₄)CH(CH₃)₂), 126.7, 127.5, 129.1 and 135.2 (C₄H₃S), 159.3 (2-CH). Anal. Calc. for C₂₃H₂₉S₂N₂RuCl₃: C, 45.66; H, 4.83; N, 4.63%. Found: C, 45.71; H, 4.89; N, 4.69%.

Refinement

All H atoms attached to carbons were geometrically fixed and allowed to ride on the corresponding non-H atom with C—H = 0.96 Å, and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ of the attached C atom for methyl H atoms and $1.2U_{\text{eq}}(\text{C})$ for other H atoms.

Figures




Fig. 1. The molecular structure of the title compound, showing the atom-numbering scheme and displacement ellipsoids drawn at the 50% probability level.




Fig. 2. A packing diagram for (I). Symmetry: H19I, H14I, H11D, $-x, 1 - y, 2 - z$; H23A, $-x, 2 - y, 2 - z$; H23C, $1 + x, -1 + y, -1 + z$; H19K, H14K, H11E, $1 + x, y, -1 + z$; H19F, C11A, C12A, $1 - x, 1 - y, 1 - z$.



Fig. 3. Synthesis of the title compound.

1,3-Bis(2-thienylmethyl)-4,5-dihydroimidazolium trichlorido(η^6 -*p*-cymene)ruthenate(II)

Crystal data

(C ₁₃ H ₁₅ N ₂ S ₂)[RuCl ₃ (C ₁₀ H ₁₄)]	$Z = 2$
$M_r = 605.02$	$F_{000} = 616$
Triclinic, $P\bar{1}$	$D_x = 1.559 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation
$a = 9.910 (2) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 11.600 (2) \text{ \AA}$	Cell parameters from 5195 reflections
$c = 12.659 (3) \text{ \AA}$	$\theta = 3.3\text{--}26.4^\circ$
$\alpha = 84.95 (3)^\circ$	$\mu = 1.10 \text{ mm}^{-1}$
$\beta = 67.05 (3)^\circ$	$T = 153 (2) \text{ K}$
$\gamma = 74.14 (3)^\circ$	Rod, red
$V = 1288.8 (6) \text{ \AA}^3$	$0.48 \times 0.19 \times 0.17 \text{ mm}$

Data collection

Rigaku AFC-8S Mercury CCD diffractometer	4557 independent reflections
Radiation source: Sealed Tube	4062 reflections with $I > 2\sigma(I)$
Monochromator: Graphite Monochromator	$R_{\text{int}} = 0.019$
Detector resolution: $14.6306 \text{ pixels mm}^{-1}$	$\theta_{\text{max}} = 25.2^\circ$
$T = 153(2) \text{ K}$	$\theta_{\text{min}} = 3.3^\circ$
ω scans	$h = -11 \rightarrow 11$
Absorption correction: multi-scan (Jacobson, 1998)	$k = -11 \rightarrow 13$
$T_{\text{min}} = 0.621$, $T_{\text{max}} = 0.836$	$l = -15 \rightarrow 15$
11095 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.032$	H-atom parameters constrained
$wR(F^2) = 0.084$	$w = 1/[\sigma^2(F_o^2) + (0.0406P)^2 + 0.9404P]$
$S = 1.13$	where $P = (F_o^2 + 2F_c^2)/3$
4557 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
285 parameters	$\Delta\rho_{\text{max}} = 0.43 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.68 \text{ e \AA}^{-3}$
	Extinction correction: none

supplementary materials

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Ru1	-0.02777 (3)	0.69238 (2)	0.639788 (19)	0.03323 (9)	
C11	-0.10558 (9)	0.78990 (7)	0.82284 (6)	0.04448 (19)	
C12	0.08996 (10)	0.51572 (7)	0.72233 (7)	0.04740 (19)	
C13	-0.26047 (9)	0.62550 (8)	0.71790 (7)	0.0506 (2)	
C1	0.1893 (4)	0.7262 (3)	0.5295 (3)	0.0484 (8)	
C2	0.1573 (4)	0.6370 (3)	0.4777 (3)	0.0462 (8)	
H2	0.2310	0.5610	0.4550	0.055*	
C3	0.0226 (4)	0.6570 (4)	0.4593 (3)	0.0512 (8)	
H3	0.0058	0.5957	0.4235	0.061*	
C4	-0.0892 (4)	0.7679 (4)	0.4935 (3)	0.0534 (9)	
C5	-0.0618 (5)	0.8573 (3)	0.5467 (3)	0.0559 (9)	
H5	-0.1363	0.9327	0.5705	0.067*	
C6	0.0761 (4)	0.8346 (3)	0.5644 (3)	0.0510 (8)	
H6	0.0924	0.8953	0.6015	0.061*	
C7	0.3370 (5)	0.6951 (5)	0.5491 (4)	0.0721 (12)	
H7	0.3489	0.6158	0.5792	0.087*	
C8	0.4683 (7)	0.6885 (10)	0.4361 (5)	0.168 (4)	
H8A	0.4600	0.7659	0.4019	0.251*	
H8B	0.4671	0.6316	0.3860	0.251*	
H8C	0.5617	0.6636	0.4484	0.251*	
C9	0.3364 (6)	0.7746 (6)	0.6359 (4)	0.0889 (16)	
H9A	0.4234	0.7411	0.6554	0.133*	
H9B	0.2455	0.7811	0.7034	0.133*	
H9C	0.3398	0.8527	0.6046	0.133*	
C10	-0.2363 (5)	0.7879 (5)	0.4785 (4)	0.0864 (15)	
H10A	-0.2231	0.8116	0.4013	0.130*	
H10B	-0.3119	0.8499	0.5311	0.130*	
H10C	-0.2683	0.7151	0.4937	0.130*	
S1	0.29307 (13)	0.18189 (11)	0.86238 (11)	0.0636 (4)	0.847 (5)
C16'	0.29307 (13)	0.18189 (11)	0.86238 (11)	0.0636 (4)	0.153 (5)
H16'	0.2049	0.2487	0.8803	0.076*	0.153 (5)
S2	0.13328 (18)	0.84868 (10)	0.98665 (14)	0.0673 (5)	0.700 (5)
C21'	0.13328 (18)	0.84868 (10)	0.98665 (14)	0.0673 (5)	0.300 (5)

H21'	0.1475	0.8704	0.9088	0.081*	0.300 (5)
N1	0.4121 (3)	0.3556 (2)	0.9546 (2)	0.0456 (6)	
N2	0.2449 (3)	0.5262 (2)	0.9644 (2)	0.0396 (6)	
C11	0.2748 (3)	0.4218 (3)	1.0096 (3)	0.0374 (6)	
H11	0.2029	0.3959	1.0769	0.045*	
C12	0.3763 (4)	0.5391 (3)	0.8622 (3)	0.0561 (9)	
H12A	0.4119	0.6054	0.8711	0.067*	
H12B	0.3525	0.5501	0.7948	0.067*	
C13	0.4936 (5)	0.4194 (3)	0.8565 (3)	0.0631 (10)	
H13A	0.5238	0.3766	0.7860	0.076*	
H13B	0.5819	0.4315	0.8633	0.076*	
C14	0.4669 (4)	0.2283 (3)	0.9737 (3)	0.0473 (8)	
H14A	0.4110	0.2105	1.0515	0.057*	
H14B	0.5718	0.2112	0.9631	0.057*	
C15	0.4510 (4)	0.1486 (3)	0.8940 (3)	0.0438 (7)	
Si1'	0.5553 (4)	0.0368 (3)	0.8423 (3)	0.0759 (12)	0.153 (5)
C16	0.5553 (4)	0.0368 (3)	0.8423 (3)	0.0759 (12)	0.847 (5)
H16	0.6534	0.0029	0.8456	0.091*	0.847 (5)
C17	0.4850 (6)	-0.0147 (4)	0.7846 (4)	0.0743 (12)	
H17	0.5312	-0.0919	0.7467	0.089*	
C18	0.3504 (6)	0.0533 (4)	0.7880 (4)	0.0732 (12)	
H18	0.2925	0.0307	0.7515	0.088*	
C19	0.1011 (4)	0.6183 (3)	1.0051 (3)	0.0409 (7)	
H19A	0.0249	0.5844	1.0610	0.049*	
H19B	0.0701	0.6438	0.9419	0.049*	
C20	0.1103 (3)	0.7249 (3)	1.0576 (3)	0.0407 (7)	
S2'	0.0919 (3)	0.7370 (2)	1.18444 (19)	0.0656 (8)	0.300 (5)
C21	0.0919 (3)	0.7370 (2)	1.18444 (19)	0.0656 (8)	0.700 (5)
H21	0.0762	0.6796	1.2445	0.079*	0.700 (5)
C22	0.1046 (5)	0.8587 (6)	1.1876 (5)	0.0860 (17)	
H22	0.0993	0.8925	1.2561	0.103*	
C23	0.1235 (5)	0.9205 (3)	1.0935 (5)	0.0765 (14)	
H23	0.1310	1.0017	1.0887	0.092*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ru1	0.04018 (15)	0.03358 (14)	0.03059 (14)	-0.01658 (10)	-0.01405 (10)	0.00217 (9)
Cl1	0.0592 (5)	0.0374 (4)	0.0377 (4)	-0.0148 (3)	-0.0163 (3)	-0.0067 (3)
Cl2	0.0554 (5)	0.0370 (4)	0.0484 (4)	-0.0104 (3)	-0.0204 (4)	0.0055 (3)
Cl3	0.0483 (4)	0.0661 (5)	0.0460 (4)	-0.0327 (4)	-0.0158 (3)	0.0045 (4)
C1	0.0502 (19)	0.064 (2)	0.0352 (16)	-0.0304 (17)	-0.0113 (14)	0.0064 (15)
C2	0.0491 (18)	0.057 (2)	0.0325 (15)	-0.0231 (16)	-0.0090 (13)	-0.0027 (14)
C3	0.063 (2)	0.074 (2)	0.0285 (15)	-0.0371 (19)	-0.0185 (15)	0.0043 (15)
C4	0.059 (2)	0.072 (2)	0.0411 (17)	-0.0304 (19)	-0.0257 (16)	0.0210 (17)
C5	0.077 (3)	0.0430 (18)	0.0461 (19)	-0.0207 (17)	-0.0225 (18)	0.0171 (15)
C6	0.072 (2)	0.0484 (19)	0.0451 (18)	-0.0386 (18)	-0.0222 (17)	0.0124 (15)
C7	0.052 (2)	0.115 (4)	0.059 (2)	-0.040 (2)	-0.0186 (18)	-0.004 (2)

supplementary materials

C8	0.075 (4)	0.359 (14)	0.081 (4)	-0.099 (6)	-0.003 (3)	-0.044 (6)
C9	0.079 (3)	0.136 (5)	0.076 (3)	-0.051 (3)	-0.040 (3)	-0.002 (3)
C10	0.072 (3)	0.131 (5)	0.073 (3)	-0.032 (3)	-0.046 (2)	0.026 (3)
S1	0.0542 (7)	0.0669 (8)	0.0728 (8)	-0.0065 (5)	-0.0317 (6)	-0.0088 (6)
C16'	0.0542 (7)	0.0669 (8)	0.0728 (8)	-0.0065 (5)	-0.0317 (6)	-0.0088 (6)
S2	0.1003 (11)	0.0355 (6)	0.0966 (11)	-0.0253 (6)	-0.0681 (9)	0.0179 (6)
C21'	0.1003 (11)	0.0355 (6)	0.0966 (11)	-0.0253 (6)	-0.0681 (9)	0.0179 (6)
N1	0.0445 (15)	0.0354 (14)	0.0491 (15)	-0.0070 (12)	-0.0116 (12)	-0.0012 (12)
N2	0.0473 (14)	0.0315 (13)	0.0384 (13)	-0.0116 (11)	-0.0141 (11)	0.0016 (10)
C11	0.0430 (16)	0.0340 (15)	0.0372 (15)	-0.0124 (13)	-0.0158 (13)	0.0010 (12)
C12	0.060 (2)	0.050 (2)	0.0482 (19)	-0.0202 (17)	-0.0080 (17)	0.0082 (16)
C13	0.060 (2)	0.051 (2)	0.059 (2)	-0.0159 (18)	-0.0004 (18)	-0.0002 (18)
C14	0.0436 (17)	0.0380 (17)	0.0561 (19)	0.0002 (14)	-0.0208 (15)	-0.0042 (15)
C15	0.0430 (17)	0.0376 (16)	0.0485 (18)	-0.0077 (13)	-0.0172 (14)	0.0023 (14)
S1'	0.092 (2)	0.0512 (17)	0.090 (2)	-0.0105 (15)	-0.0439 (19)	-0.0045 (15)
C16	0.092 (2)	0.0512 (17)	0.090 (2)	-0.0105 (15)	-0.0439 (19)	-0.0045 (15)
C17	0.097 (3)	0.053 (2)	0.068 (3)	-0.021 (2)	-0.023 (2)	-0.011 (2)
C18	0.095 (3)	0.075 (3)	0.070 (3)	-0.038 (3)	-0.044 (3)	0.005 (2)
C19	0.0448 (17)	0.0319 (15)	0.0511 (18)	-0.0088 (13)	-0.0242 (14)	-0.0002 (13)
C20	0.0374 (15)	0.0326 (15)	0.0533 (18)	-0.0059 (12)	-0.0197 (14)	-0.0032 (13)
S2'	0.0721 (15)	0.0671 (15)	0.0606 (13)	-0.0293 (11)	-0.0207 (11)	0.0003 (10)
C21	0.0721 (15)	0.0671 (15)	0.0606 (13)	-0.0293 (11)	-0.0207 (11)	0.0003 (10)
C22	0.060 (3)	0.115 (4)	0.082 (3)	-0.020 (3)	-0.015 (2)	-0.053 (3)
C23	0.067 (3)	0.0316 (18)	0.133 (5)	-0.0068 (18)	-0.041 (3)	-0.013 (2)

Geometric parameters (Å, °)

Ru1—C6	2.140 (3)	C10—H10C	0.9599
Ru1—C2	2.154 (3)	S1—C15	1.700 (3)
Ru1—C5	2.173 (3)	S2—C20	1.647 (3)
Ru1—C1	2.180 (3)	N1—C11	1.307 (4)
Ru1—C3	2.191 (3)	N1—C13	1.458 (5)
Ru1—C4	2.207 (3)	N1—C14	1.462 (4)
Ru1—Cl1	2.4157 (11)	N2—C11	1.302 (4)
Ru1—Cl2	2.4329 (11)	N2—C19	1.460 (4)
Ru1—Cl3	2.4417 (11)	N2—C12	1.466 (4)
C1—C6	1.406 (5)	C11—H11	0.9600
C1—C2	1.435 (5)	C12—C13	1.534 (5)
C1—C7	1.521 (5)	C12—H12A	0.9600
C2—C3	1.398 (5)	C12—H12B	0.9600
C2—H2	0.9600	C13—H13A	0.9600
C3—C4	1.419 (6)	C13—H13B	0.9600
C3—H3	0.9600	C14—C15	1.505 (5)
C4—C5	1.422 (5)	C14—H14A	0.9600
C4—C10	1.495 (5)	C14—H14B	0.9600
C5—C6	1.421 (5)	C15—S1'	1.438 (4)
C5—H5	0.9600	S1'—H16	0.960 (3)
C6—H6	0.9600	C17—C18	1.338 (7)
C7—C9	1.492 (6)	C17—H17	0.9600

C7—C8	1.502 (7)	C18—H18	0.9600
C7—H7	0.9600	C19—C20	1.495 (4)
C8—H8A	0.9599	C19—H19A	0.9600
C8—H8B	0.9599	C19—H19B	0.9600
C8—H8C	0.9599	C20—S2'	1.558 (4)
C9—H9A	0.9599	S2'—H21	0.960 (2)
C9—H9B	0.9599	C22—C23	1.310 (7)
C9—H9C	0.9599	C22—H22	0.9600
C10—H10A	0.9599	C23—H23	0.9600
C10—H10B	0.9599		
C6—Ru1—C2	68.59 (14)	Ru1—C6—H6	128.9
C6—Ru1—C5	38.46 (15)	C9—C7—C8	113.3 (5)
C2—Ru1—C5	80.93 (15)	C9—C7—C1	113.7 (4)
C6—Ru1—C1	37.98 (14)	C8—C7—C1	109.7 (4)
C2—Ru1—C1	38.66 (13)	C9—C7—H7	106.5
C5—Ru1—C1	69.25 (15)	C8—C7—H7	106.5
C6—Ru1—C3	81.08 (14)	C1—C7—H7	106.5
C2—Ru1—C3	37.52 (13)	C7—C8—H8A	109.5
C5—Ru1—C3	68.20 (15)	C7—C8—H8B	109.5
C1—Ru1—C3	69.15 (13)	H8A—C8—H8B	109.5
C6—Ru1—C4	68.92 (14)	C7—C8—H8C	109.5
C2—Ru1—C4	68.06 (14)	H8A—C8—H8C	109.5
C5—Ru1—C4	37.88 (15)	H8B—C8—H8C	109.5
C1—Ru1—C4	81.91 (13)	C7—C9—H9A	109.5
C3—Ru1—C4	37.65 (15)	C7—C9—H9B	109.5
C6—Ru1—Cl1	86.77 (10)	H9A—C9—H9B	109.5
C2—Ru1—Cl1	144.83 (9)	C7—C9—H9C	109.5
C5—Ru1—Cl1	95.39 (11)	H9A—C9—H9C	109.5
C1—Ru1—Cl1	107.28 (10)	H9B—C9—H9C	109.5
C3—Ru1—Cl1	163.52 (11)	C4—C10—H10A	109.5
C4—Ru1—Cl1	126.94 (11)	C4—C10—H10B	109.5
C6—Ru1—Cl2	124.11 (11)	H10A—C10—H10B	109.5
C2—Ru1—Cl2	87.76 (11)	C4—C10—H10C	109.5
C5—Ru1—Cl2	162.10 (11)	H10A—C10—H10C	109.5
C1—Ru1—Cl2	93.29 (10)	H10B—C10—H10C	109.5
C3—Ru1—Cl2	110.15 (11)	C11—N1—C13	110.3 (3)
C4—Ru1—Cl2	146.86 (11)	C11—N1—C14	125.0 (3)
Cl1—Ru1—Cl2	85.88 (4)	C13—N1—C14	123.6 (3)
C6—Ru1—Cl3	147.20 (11)	C11—N2—C19	126.2 (3)
C2—Ru1—Cl3	126.23 (9)	C11—N2—C12	110.5 (3)
C5—Ru1—Cl3	110.17 (12)	C19—N2—C12	123.3 (3)
C1—Ru1—Cl3	164.68 (9)	N2—C11—N1	113.6 (3)
C3—Ru1—Cl3	96.15 (10)	N2—C11—H11	123.2
C4—Ru1—Cl3	88.90 (10)	N1—C11—H11	123.2
Cl1—Ru1—Cl3	88.04 (4)	N2—C12—C13	102.5 (3)
Cl2—Ru1—Cl3	87.71 (4)	N2—C12—H12A	111.3
C6—C1—C2	116.7 (3)	C13—C12—H12A	111.3
C6—C1—C7	124.6 (4)	N2—C12—H12B	111.3
C2—C1—C7	118.6 (4)	C13—C12—H12B	111.3

supplementary materials

C6—C1—Ru1	69.44 (19)	H12A—C12—H12B	109.2
C2—C1—Ru1	69.67 (19)	N1—C13—C12	103.2 (3)
C7—C1—Ru1	128.6 (3)	N1—C13—H13A	111.1
C3—C2—C1	122.3 (3)	C12—C13—H13A	111.1
C3—C2—Ru1	72.69 (19)	N1—C13—H13B	111.1
C1—C2—Ru1	71.67 (19)	C12—C13—H13B	111.1
C3—C2—H2	118.9	H13A—C13—H13B	109.1
C1—C2—H2	118.9	N1—C14—C15	112.6 (3)
Ru1—C2—H2	129.3	N1—C14—H14A	109.1
C2—C3—C4	120.1 (3)	C15—C14—H14A	109.1
C2—C3—Ru1	69.79 (18)	N1—C14—H14B	109.1
C4—C3—Ru1	71.78 (19)	C15—C14—H14B	109.1
C2—C3—H3	119.9	H14A—C14—H14B	107.8
C4—C3—H3	119.9	S1'—C15—C14	126.7 (3)
Ru1—C3—H3	131.2	S1'—C15—S1	112.4 (3)
C3—C4—C5	118.9 (3)	C14—C15—S1	120.8 (2)
C3—C4—C10	120.4 (4)	C15—S1'—H16	126.4 (3)
C5—C4—C10	120.7 (4)	C18—C17—H17	122.6
C3—C4—Ru1	70.57 (19)	C17—C18—H18	123.4
C5—C4—Ru1	69.77 (19)	N2—C19—C20	112.8 (3)
C10—C4—Ru1	129.5 (3)	N2—C19—H19A	109.0
C6—C5—C4	119.9 (4)	C20—C19—H19A	109.0
C6—C5—Ru1	69.50 (19)	N2—C19—H19B	109.0
C4—C5—Ru1	72.4 (2)	C20—C19—H19B	109.0
C6—C5—H5	120.1	H19A—C19—H19B	107.8
C4—C5—H5	120.1	C19—C20—S2'	125.7 (3)
Ru1—C5—H5	130.7	C19—C20—S2	122.6 (3)
C1—C6—C5	122.1 (3)	S2'—C20—S2	111.7 (2)
C1—C6—Ru1	72.58 (19)	C20—S2'—H21	129.3 (2)
C5—C6—Ru1	72.0 (2)	C23—C22—H22	121.2
C1—C6—H6	119.0	C22—C23—H23	122.8
C5—C6—H6	119.0		
C2—Ru1—C1—C6	130.7 (3)	Cl2—Ru1—C4—C5	-150.0 (2)
C5—Ru1—C1—C6	28.9 (2)	Cl3—Ru1—C4—C5	125.9 (2)
C3—Ru1—C1—C6	102.6 (2)	C6—Ru1—C4—C10	-143.0 (5)
C4—Ru1—C1—C6	65.9 (2)	C2—Ru1—C4—C10	142.5 (5)
Cl1—Ru1—C1—C6	-60.4 (2)	C5—Ru1—C4—C10	-113.6 (5)
Cl2—Ru1—C1—C6	-147.07 (19)	C1—Ru1—C4—C10	180.0 (5)
Cl3—Ru1—C1—C6	119.6 (4)	C3—Ru1—C4—C10	113.8 (5)
C6—Ru1—C1—C2	-130.7 (3)	Cl1—Ru1—C4—C10	-74.4 (5)
C5—Ru1—C1—C2	-101.8 (2)	Cl2—Ru1—C4—C10	96.4 (4)
C3—Ru1—C1—C2	-28.1 (2)	Cl3—Ru1—C4—C10	12.3 (4)
C4—Ru1—C1—C2	-64.8 (2)	C3—C4—C5—C6	0.2 (5)
Cl1—Ru1—C1—C2	168.98 (18)	C10—C4—C5—C6	177.4 (3)
Cl2—Ru1—C1—C2	82.3 (2)	Ru1—C4—C5—C6	52.7 (3)
Cl3—Ru1—C1—C2	-11.0 (5)	C3—C4—C5—Ru1	-52.5 (3)
C6—Ru1—C1—C7	118.5 (4)	C10—C4—C5—Ru1	124.7 (3)
C2—Ru1—C1—C7	-110.9 (4)	C2—Ru1—C5—C6	-66.8 (2)
C5—Ru1—C1—C7	147.4 (4)	C1—Ru1—C5—C6	-28.6 (2)

C3—Ru1—C1—C7	-138.9 (4)	C3—Ru1—C5—C6	-103.6 (2)
C4—Ru1—C1—C7	-175.6 (4)	C4—Ru1—C5—C6	-132.6 (3)
C11—Ru1—C1—C7	58.1 (4)	C11—Ru1—C5—C6	77.9 (2)
C12—Ru1—C1—C7	-28.6 (4)	C12—Ru1—C5—C6	-15.4 (5)
C13—Ru1—C1—C7	-121.9 (4)	C13—Ru1—C5—C6	167.78 (19)
C6—C1—C2—C3	1.9 (5)	C6—Ru1—C5—C4	132.6 (3)
C7—C1—C2—C3	178.4 (3)	C2—Ru1—C5—C4	65.7 (2)
Ru1—C1—C2—C3	54.6 (3)	C1—Ru1—C5—C4	104.0 (2)
C6—C1—C2—Ru1	-52.7 (3)	C3—Ru1—C5—C4	29.0 (2)
C7—C1—C2—Ru1	123.8 (3)	C11—Ru1—C5—C4	-149.5 (2)
C6—Ru1—C2—C3	-103.7 (2)	C12—Ru1—C5—C4	117.2 (3)
C5—Ru1—C2—C3	-65.8 (2)	C13—Ru1—C5—C4	-59.6 (2)
C1—Ru1—C2—C3	-133.8 (3)	C2—C1—C6—C5	-1.9 (5)
C4—Ru1—C2—C3	-28.7 (2)	C7—C1—C6—C5	-178.1 (3)
C11—Ru1—C2—C3	-152.29 (18)	Ru1—C1—C6—C5	-54.7 (3)
C12—Ru1—C2—C3	128.1 (2)	C2—C1—C6—Ru1	52.8 (3)
C13—Ru1—C2—C3	42.6 (3)	C7—C1—C6—Ru1	-123.4 (3)
C6—Ru1—C2—C1	30.1 (2)	C4—C5—C6—C1	0.9 (5)
C5—Ru1—C2—C1	68.0 (2)	Ru1—C5—C6—C1	54.9 (3)
C3—Ru1—C2—C1	133.8 (3)	C4—C5—C6—Ru1	-54.0 (3)
C4—Ru1—C2—C1	105.1 (2)	C2—Ru1—C6—C1	-30.6 (2)
C11—Ru1—C2—C1	-18.5 (3)	C5—Ru1—C6—C1	-133.4 (3)
C12—Ru1—C2—C1	-98.1 (2)	C3—Ru1—C6—C1	-67.4 (2)
C13—Ru1—C2—C1	176.40 (17)	C4—Ru1—C6—C1	-104.4 (2)
C1—C2—C3—C4	-0.9 (5)	C11—Ru1—C6—C1	123.77 (19)
Ru1—C2—C3—C4	53.2 (3)	C12—Ru1—C6—C1	41.0 (2)
C1—C2—C3—Ru1	-54.1 (3)	C13—Ru1—C6—C1	-154.91 (17)
C6—Ru1—C3—C2	66.3 (2)	C2—Ru1—C6—C5	102.8 (2)
C5—Ru1—C3—C2	104.0 (2)	C1—Ru1—C6—C5	133.4 (3)
C1—Ru1—C3—C2	28.8 (2)	C3—Ru1—C6—C5	66.0 (2)
C4—Ru1—C3—C2	133.2 (3)	C4—Ru1—C6—C5	29.0 (2)
C11—Ru1—C3—C2	109.2 (4)	C11—Ru1—C6—C5	-102.8 (2)
C12—Ru1—C3—C2	-56.9 (2)	C12—Ru1—C6—C5	174.34 (18)
C13—Ru1—C3—C2	-146.7 (2)	C13—Ru1—C6—C5	-21.5 (3)
C6—Ru1—C3—C4	-66.9 (2)	C6—C1—C7—C9	15.6 (6)
C2—Ru1—C3—C4	-133.2 (3)	C2—C1—C7—C9	-160.6 (4)
C5—Ru1—C3—C4	-29.1 (2)	Ru1—C1—C7—C9	-74.7 (5)
C1—Ru1—C3—C4	-104.3 (2)	C6—C1—C7—C8	-112.5 (6)
C11—Ru1—C3—C4	-23.9 (5)	C2—C1—C7—C8	71.4 (6)
C12—Ru1—C3—C4	169.95 (18)	Ru1—C1—C7—C8	157.2 (5)
C13—Ru1—C3—C4	80.1 (2)	C19—N2—C11—N1	178.8 (3)
C2—C3—C4—C5	-0.2 (5)	C12—N2—C11—N1	0.4 (4)
Ru1—C3—C4—C5	52.1 (3)	C13—N1—C11—N2	0.0 (4)
C2—C3—C4—C10	-177.4 (3)	C14—N1—C11—N2	-168.6 (3)
Ru1—C3—C4—C10	-125.1 (3)	C11—N2—C12—C13	-0.6 (4)
C2—C3—C4—Ru1	-52.3 (3)	C19—N2—C12—C13	-179.1 (3)
C6—Ru1—C4—C3	103.2 (2)	C11—N1—C13—C12	-0.4 (4)
C2—Ru1—C4—C3	28.6 (2)	C14—N1—C13—C12	168.4 (3)
C5—Ru1—C4—C3	132.6 (3)	N2—C12—C13—N1	0.6 (4)

supplementary materials

C1—Ru1—C4—C3	66.1 (2)	C11—N1—C14—C15	93.3 (4)
C11—Ru1—C4—C3	171.72 (16)	C13—N1—C14—C15	-73.9 (4)
C12—Ru1—C4—C3	-17.4 (3)	N1—C14—C15—S1'	143.0 (3)
C13—Ru1—C4—C3	-101.6 (2)	N1—C14—C15—S1	-40.9 (4)
C6—Ru1—C4—C5	-29.4 (2)	C11—N2—C19—C20	109.6 (4)
C2—Ru1—C4—C5	-103.9 (2)	C12—N2—C19—C20	-72.2 (4)
C1—Ru1—C4—C5	-66.4 (2)	N2—C19—C20—S2'	-84.8 (4)
C3—Ru1—C4—C5	-132.6 (3)	N2—C19—C20—S2	98.6 (3)
C11—Ru1—C4—C5	39.2 (3)		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C11—H11 \cdots Cl1 ⁱ	0.96	2.62	3.450 (4)	144
C14—H14A \cdots Cl1 ⁱ	0.96	2.82	3.553 (4)	134
C19—H19A \cdots Cl2 ⁱ	0.96	2.81	3.671 (4)	150
C23—H23 \cdots Cl1 ⁱⁱ	0.96	2.66	3.549 (4)	154
C14—H14B \cdots Cg2 ⁱⁱⁱ	0.96	2.83	3.784 (5)	171
C19—H19B \cdots Cl1	0.96	2.86	3.759 (5)	157

Symmetry codes: (i) $-x, -y+1, -z+2$; (ii) $-x, -y+2, -z+2$; (iii) $-x+1, -y+1, -z+2$.

Fig. 1




Fig. 2




Fig. 3

