

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N,N'-Bis[(*E*)-4-cyanobenzylidene]urea

De-Hong Wu* and Ling Hu

 Ordered Matter Science Research Center, College of Chemistry and Chemical Engineering, Southeast University, Nanjing 210096, People's Republic of China
 Correspondence e-mail: wudh1971@sohu.com


Received 6 January 2009; accepted 17 January 2009

 Key indicators: single-crystal X-ray study; $T = 291$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.072; wR factor = 0.164; data-to-parameter ratio = 14.0.

The molecule of the title compound, $\text{C}_{17}\text{H}_{10}\text{N}_4\text{O}$, has crystallographically imposed C_2 symmetry. The urea group and the benzene ring are nearly coplanar, the dihedral angle between them being $4.15(7)^\circ$. The crystal packing is stabilized by aromatic π - π stacking interactions, with a centroid-to-centroid separation of $3.833(4)$ Å.

Related literature

For a general background on the use of nitriles as starting materials, see: Íkizler & Sancak (1992). For the products of the condensation of urea with alkynes, see: Martínez-García *et al.* (2004).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{10}\text{N}_4\text{O}$	$V = 1481.7(9)$ Å ³
$M_r = 286.29$	$Z = 4$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 10.552(4)$ Å	$\mu = 0.08$ mm ⁻¹
$b = 11.687(5)$ Å	$T = 291(2)$ K
$c = 12.198(3)$ Å	$0.36 \times 0.30 \times 0.28$ mm
$\beta = 99.94(4)^\circ$	

Data collection

Rigaku Mercury2 diffractometer	6461 measured reflections
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku, 2005)	1423 independent reflections
$T_{\min} = 0.96$, $T_{\max} = 0.98$	1107 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.041$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.072$	1 restraint
$wR(F^2) = 0.164$	H-atom parameters constrained
$S = 1.06$	$\Delta\rho_{\text{max}} = 0.43$ e Å ⁻³
1423 reflections	$\Delta\rho_{\text{min}} = -0.27$ e Å ⁻³
102 parameters	

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank Jiangsu Planned Projects for Post-doctoral Research Funds (grant No. 0802003B) and Professor Dr Rengen Xiong.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2289).

References

- Íkizler, A. A. & Sancak, K. (1992). *Monatsh. Chem.* **123**, 257–263.
 Martínez-García, A., Ortiz, M., Martínez, R., Ortiz, P. & Reguera, E. (2004). *Ind. Crops Prod.* **19**, 99–106.
 Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2009). E65, o371 [doi:10.1107/S1600536809002244]

N,N'-Bis[(*E*)-4-cyanobenzylidene]urea

De-Hong Wu and Ling Hu

S1. Comment

Nitriles are parent compounds for the preparation of various functional organic materials having imidazole, triazole, or thiazole functionalities (İkizler & Sancak, 1992). The Schiff-base compounds derived from the condensation of urea with alkynes are very few (Martínez-García *et al.*, 2004) because of the low reactivity of the NH₂ group of urea. Here we report the crystal structure of the title compound, which was obtained by the reaction of urea with 4-cyanobenzaldehyde in acetic acid with ammonium chloride as a catalyzer.

The molecule of the title compound (Fig. 1) possesses a crystallographically imposed *C*₂ symmetry, with atoms C1 and O1 located on a two-fold axis. The urea group and the aromatic rings are nearly coplanar, forming a dihedral angle of 4.15 (7)°. In the crystal packing, molecules are linked along the *a* axis by aromatic π - π stacking interactions, with centroid-to-centroid separations of 3.833 (4) Å, perpendicular interplanar distances of 3.474 (4) Å and centroid-centroid offsets of 1.620 (3) Å.

S2. Experimental

A mixture of 4-cyanobenzaldehyde (0.53 g, 4 mmol), urea (0.36 g, 6 mmol) and NH₄Cl (0.10 g, 1.6 mmol) was heated with stirring at 100°C in 5 ml acetic acid for 5 h. After cooling, the reaction mixture was washed with cold water (3 × 50 ml) and the residue recrystallized from ethyl acetate/*n*-hexane (1:2 v/v) to afford the title compound (0.57 g, 50%). Single crystals suitable for X-ray structure analysis were obtained by the slow evaporation of an ethyl acetate solution in air.

S3. Refinement

All H atoms were placed in calculated positions and refined using a riding model approximation, with C—H = 0.93 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.




Figure 1

The molecular structure of the title compound, showing the atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. Unlabelled atoms are related to the labelled atoms by (-*x*, *y*, 1/2 - *z*).

N,N'*-Bis[(*E*)-4-cyanobenzylidene]ureaCrystal data*C₁₇H₁₀N₄O $M_r = 286.29$ Monoclinic, *C2/c*

Hall symbol: -C 2yc

 $a = 10.552 (4) \text{ \AA}$ $b = 11.687 (5) \text{ \AA}$ $c = 12.198 (3) \text{ \AA}$ $\beta = 99.94 (4)^\circ$ $V = 1481.7 (9) \text{ \AA}^3$ $Z = 4$ $F(000) = 592$ $D_x = 1.283 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1587 reflections

 $\theta = 2.6\text{--}27.4^\circ$ $\mu = 0.08 \text{ mm}^{-1}$ $T = 291 \text{ K}$

Block, yellow

 $0.36 \times 0.30 \times 0.28 \text{ mm}$ *Data collection*

Rigaku Mercury2

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 13.6612 pixels mm⁻¹

CCD profile fitting scans

Absorption correction: multi-scan

(CrystalClear; Rigaku, 2005)

 $T_{\min} = 0.96, T_{\max} = 0.98$

6461 measured reflections

1423 independent reflections

1107 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.041$ $\theta_{\max} = 26.0^\circ, \theta_{\min} = 2.9^\circ$ $h = -12 \rightarrow 12$ $k = -14 \rightarrow 14$ $l = -15 \rightarrow 15$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.072$ $wR(F^2) = 0.164$ $S = 1.06$

1423 reflections

102 parameters

1 restraint

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0535P)^2 + 1.99P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.43 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.27 \text{ e \AA}^{-3}$ *Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.0000	0.1727 (3)	0.2500	0.0459 (8)
C2	0.0655 (2)	0.1864 (2)	0.45184 (18)	0.0476 (6)
H2A	0.0645	0.1068	0.4491	0.057*
C3	0.1028 (2)	0.2366 (2)	0.56187 (18)	0.0419 (6)

C4	0.1265 (2)	0.1659 (2)	0.65419 (19)	0.0505 (6)
H4A	0.1174	0.0872	0.6445	0.061*
C5	0.1636 (2)	0.2094 (2)	0.7610 (2)	0.0526 (7)
H5A	0.1790	0.1605	0.8220	0.063*
C6	0.1773 (2)	0.3265 (2)	0.77534 (19)	0.0484 (6)
C7	0.2135 (3)	0.3729 (2)	0.8858 (2)	0.0564 (7)
C8	0.1546 (2)	0.3991 (2)	0.6841 (2)	0.0544 (7)
H8A	0.1640	0.4777	0.6942	0.065*
C9	0.1181 (2)	0.3551 (2)	0.5786 (2)	0.0525 (7)
H9A	0.1034	0.4043	0.5179	0.063*
N1	0.0325 (2)	0.2397 (2)	0.35441 (17)	0.0643 (7)
N2	0.2416 (3)	0.4101 (2)	0.9729 (2)	0.0772 (8)
O1	0.0000	0.0683 (2)	0.2500	0.0655 (8)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0458 (18)	0.055 (2)	0.0363 (17)	0.000	0.0039 (13)	0.000
C2	0.0485 (13)	0.0509 (14)	0.0428 (12)	-0.0062 (10)	0.0065 (10)	-0.0032 (9)
C3	0.0412 (12)	0.0466 (13)	0.0384 (12)	-0.0019 (9)	0.0081 (9)	0.0009 (9)
C4	0.0628 (15)	0.0472 (14)	0.0412 (13)	-0.0034 (11)	0.0085 (10)	-0.0011 (10)
C5	0.0608 (16)	0.0575 (16)	0.0394 (12)	0.0015 (12)	0.0087 (11)	0.0065 (11)
C6	0.0449 (13)	0.0595 (15)	0.0404 (13)	0.0016 (11)	0.0058 (10)	-0.0095 (11)
C7	0.0638 (16)	0.0577 (16)	0.0468 (15)	0.0041 (12)	0.0073 (12)	-0.0036 (12)
C8	0.0626 (16)	0.0484 (14)	0.0509 (14)	-0.0036 (11)	0.0063 (11)	-0.0025 (11)
C9	0.0604 (15)	0.0522 (14)	0.0435 (13)	-0.0021 (12)	0.0049 (11)	0.0055 (11)
N1	0.0696 (15)	0.0745 (17)	0.0472 (12)	0.0003 (12)	0.0055 (10)	0.0023 (10)
N2	0.106 (2)	0.0707 (17)	0.0511 (14)	0.0061 (14)	0.0034 (13)	-0.0151 (12)
O1	0.098 (2)	0.0506 (16)	0.0442 (14)	0.000	0.0009 (13)	0.000

Geometric parameters (Å, °)

C1—O1	1.221 (4)	C4—H4A	0.9300
C1—N1	1.484 (3)	C5—C6	1.384 (4)
C1—N1 ⁱ	1.484 (3)	C5—H5A	0.9300
C2—N1	1.334 (3)	C6—C8	1.387 (4)
C2—C3	1.455 (3)	C6—C7	1.442 (3)
C2—H2A	0.9300	C7—N2	1.138 (3)
C3—C4	1.384 (3)	C8—C9	1.377 (3)
C3—C9	1.405 (3)	C8—H8A	0.9300
C4—C5	1.390 (3)	C9—H9A	0.9300
O1—C1—N1	121.83 (16)	C6—C5—H5A	120.5
O1—C1—N1 ⁱ	121.83 (16)	C4—C5—H5A	120.5
N1—C1—N1 ⁱ	116.3 (3)	C5—C6—C8	120.2 (2)
N1—C2—C3	128.4 (2)	C5—C6—C7	119.7 (2)
N1—C2—H2A	115.8	C8—C6—C7	120.1 (2)
C3—C2—H2A	115.8	N2—C7—C6	179.5 (3)

C4—C3—C9	118.0 (2)	C9—C8—C6	120.2 (2)
C4—C3—C2	119.5 (2)	C9—C8—H8A	119.9
C9—C3—C2	122.5 (2)	C6—C8—H8A	119.9
C3—C4—C5	121.8 (2)	C8—C9—C3	120.7 (2)
C3—C4—H4A	119.1	C8—C9—H9A	119.7
C5—C4—H4A	119.1	C3—C9—H9A	119.7
C6—C5—C4	119.1 (2)	C2—N1—C1	120.3 (3)

Symmetry code: (i) $-x, y, -z+1/2$.