

## Praustinoid A: a meroterpene produced by *Penicillium* sp.

Stella H. Maganhi,<sup>a</sup> Taicia Pacheco Fill,<sup>a</sup> Edson Rodrigues-Fo,<sup>a</sup> Ignez Caracelli<sup>b</sup> and Julio Zukerman-Schpector<sup>a\*</sup>

<sup>a</sup>Department of Chemistry, Universidade Federal de São Carlos, 13565-905 São Carlos, SP, Brazil, and <sup>b</sup>BioMat - Physics Department, Universidade Estadual Paulista "Júlio de Mesquita Filho", UNESP, 17033-360 Bauru, SP, Brazil

Correspondence e-mail: julio@power.ufscar.br


Received 17 December 2008; accepted 20 December 2008

Key indicators: single-crystal X-ray study;  $T = 290$  K; mean  $\sigma(C-C) = 0.003$  Å;  $R$  factor = 0.036;  $wR$  factor = 0.093; data-to-parameter ratio = 9.9.

The title meroterpene praustinoid A (systematic name: methyl 15-hydroxy-2,6,6,10,13,15-hexamethyl-17-methylene-7,14,16-trioxotetracyclo[11.3.1.0<sup>2,11</sup>.0<sup>5,10</sup>]heptadecane-1-carboxylate),  $C_{26}H_{36}O_6$ , features a fused four-ring arrangement. Three rings are in different distorted chair conformations and the other is in a distorted boat conformation. The absolute configuration was established based on  $[\alpha_D] = -4.97^\circ$  ( $c = 1.10\text{ g l}^{-1}$ ,  $\text{CH}_2\text{Cl}_2$ ). In the crystal, the molecules are connected into supramolecular chains via O—H···O hydrogen bonds.

### Related literature

For related literature, see: dos Santos & Rodrigues-Fo (2002). For structure analysis, see: Cremer and Pople (1975); Iulek and Zukerman-Schpector (1997).



### Experimental

#### Crystal data

$C_{26}H_{36}O_6$	$V = 2269.43(8)$ Å <sup>3</sup>
$M_r = 444.55$	$Z = 4$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
$a = 8.5023(2)$ Å	$\mu = 0.09$ mm <sup>-1</sup>
$b = 13.5405(2)$ Å	$T = 290(2)$ K
$c = 19.7127(4)$ Å	$0.37 \times 0.28 \times 0.11$ mm

#### Data collection

Bruker APEXII CCD area-detector diffractometer	2938 independent reflections
Absorption correction: none	2677 reflections with $I > 2\sigma(I)$
27134 measured reflections	$R_{\text{int}} = 0.034$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$	297 parameters
$wR(F^2) = 0.093$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\text{max}} = 0.22$ e Å <sup>-3</sup>
2938 reflections	$\Delta\rho_{\text{min}} = -0.13$ e Å <sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O5—H1O5···O3 <sup>i</sup>	0.82	2.05	2.870 (2)	173

Symmetry code: (i)  $x + \frac{1}{2}, -y + \frac{3}{2}, -z + 2$ .

Data collection: *APEX2*, *COSMO* and *BIS* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT*; program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

We thank FAPESP, CNPq and CAPES for financial support. Professor R. A. Burrow of the Federal University of Santa Maria is gratefully acknowledged for helping with the collection of the intensity data.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2346).

### References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Bruker (2006). *APEX2*, *COSMO*, *BIS* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Iulek, J. & Zukerman-Schpector, J. (1997). *Quím. Nova*, **20**, 433–434.
- Santos, R. M. G. dos & Rodrigues-Fo, E. (2002). *Phytochemistry*, **61**, 907–912.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

# supporting information

*Acta Cryst.* (2009). E65, o221 [doi:10.1107/S1600536808043481]

## **Preaustinoid A: a meroterpene produced by *Penicillium* sp.**

**Stella H. Maganhi, Taicia Pacheco Fill, Edson Rodrigues-Fo, Ignez Caracelli and Julio Zukerman-Schpector**

### **S1. Comment**


Endophytic fungi have been a rich source of important biologically active secondary metabolites, in particular meroterpenoids, a class of complex metabolites derived from a mixed terpenoid-polyketide biosynthetic pathway. During an on-going study of substances produced by endophytic fungi, the title compound (I) was isolated and its structure postulated based on APCIMS (Atmospheric Pressure Chemical Ionization Mass Spectrometry), HREIMS (High Resolution Electrospray Mass Spectrometry) and a variety of NMR studies (dos Santos and Rodrigues-Fo, 2002). As suitable crystals were subsequently obtained, a crystal structure determination was undertaken. The four fused rings are in different distorted conformations. Rings A and C are distorted towards a half-chair conformation, ring B is distorted towards a half-boat conformation, and ring D is a boat conformation that is highly distorted towards a half-boat. The ring-puckering parameters (Cremer & Pople, 1975, Iulek & Zukerman-Shpector, 1997) for rings A,B,C, D are:  $q_2 = 0.062$  (2), 0.093 (2), 0.059 (2), 0.582 (2) Å,  $q_3 = -0.524$  (2), 0.577 (2), -0.624 (2), -0.139 (2) Å,  $Q = 0.528$  (2), 0.584 (2), 0.627 (2), 0.599 (2) $^\circ$ ,  $\varphi_2 = 90$  (2), 172 (1), -149 (2), -122.4 (2) $^\circ$ , and  $\theta_2 = 173.3$  (2), 9.1 (2), 174.6 (2), 103.4 (2) $^\circ$ , respectively. The absolute configuration was established based on the  $[\alpha_D] = -4.97^\circ$  (c 1.10 g/L, CH<sub>2</sub>Cl<sub>2</sub>) and the results reported in dos Santos & Rodrigues-Fo (2002). The molecules are linked into a supramolecular chain *via* intermolecular O-H $\cdots$ O hydrogen bonds, Table 1.

### **S2. Experimental**

Compound (I), Preaustinoid A, was produced during cultivation of *Penicillium* sp over sterilized rice, and isolated from the methanol extract of the culture. Suitable crystals were obtained, by slow evaporation, from a mixture of dichloromethane, methanol and water.

### **S3. Refinement**

In the absence of significant anomalous scattering effects, Friedel pairs were averaged in the final refinement. The H atoms were refined in the riding-model approximation with C—H = 0.93 - 0.98 Å and (0.82 for O—H), and with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{methyl-C})$  or  $1.2U_{\text{eq}}(\text{remaining-C and -O})$ .

**Figure 1**

The molecular structure of (I) showing atom labelling scheme and displacement ellipsoids at the 30% probability level (arbitrary spheres for the H atoms).

**methyl 15-hydroxy-2,6,6,10,13,15-hexamethyl-17-methylene-7,14,16-trioxotetracyclo[11.3.1.0<sup>2,11</sup>.0<sup>5,10</sup>]heptadecane-1-carboxylate**

*Crystal data*

C<sub>26</sub>H<sub>36</sub>O<sub>6</sub>  
*M*<sub>r</sub> = 444.55  
 Orthorhombic, P2<sub>1</sub>2<sub>1</sub>2<sub>1</sub>  
 Hall symbol: P 2ac 2ab  
*a* = 8.5023 (2) Å  
*b* = 13.5405 (2) Å  
*c* = 19.7127 (4) Å

*V* = 2269.43 (8) Å<sup>3</sup>  
*Z* = 4  
*F*(000) = 960  
*D*<sub>x</sub> = 1.301 Mg m<sup>-3</sup>  
 Mo *K*α radiation,  $\lambda$  = 0.71073 Å  
 Cell parameters from 22936 reflections  
 $\theta$  = 1.0–27.5°

$\mu = 0.09 \text{ mm}^{-1}$   
 $T = 290 \text{ K}$

Prism, colorless  
 $0.37 \times 0.28 \times 0.11 \text{ mm}$

#### Data collection

Bruker APEXII CCD area-detector  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\varphi$  and  $\omega$  scans  
27134 measured reflections  
2938 independent reflections

2677 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.034$   
 $\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 1.8^\circ$   
 $h = -11 \rightarrow 9$   
 $k = -17 \rightarrow 17$   
 $l = -25 \rightarrow 25$

#### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.036$   
 $wR(F^2) = 0.093$   
 $S = 1.03$   
2938 reflections  
297 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0531P)^2 + 0.3814P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} < 0.001$   
 $\Delta\rho_{\text{max}} = 0.22 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.13 \text{ e } \text{\AA}^{-3}$   
Absolute structure:  $[\alpha_D] = -4.97^\circ$  and results in  
dos Santos & Rodrigues-Fo (2002)

#### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.6598 (2)	1.07289 (12)	1.05643 (9)	0.0288 (4)
C2	0.6226 (2)	1.08110 (12)	0.97653 (9)	0.0285 (4)
C3	0.4489 (2)	1.10698 (14)	0.96387 (10)	0.0339 (4)
H3A	0.3835	1.0666	0.9931	0.041*
H3B	0.4314	1.1756	0.9760	0.041*
C4	0.3986 (2)	1.09105 (13)	0.89014 (10)	0.0335 (4)
H4A	0.4588	1.1343	0.8607	0.040*
H4B	0.2882	1.1076	0.8851	0.040*
C5	0.4252 (2)	0.98347 (13)	0.86909 (9)	0.0281 (4)
H5	0.3789	0.9444	0.9059	0.034*
C6	0.3310 (2)	0.95241 (13)	0.80444 (10)	0.0333 (4)
C7	0.3688 (3)	0.84397 (14)	0.78946 (10)	0.0375 (4)
C8	0.5381 (3)	0.81430 (17)	0.79148 (13)	0.0480 (5)

H8A	0.5925	0.8437	0.7532	0.058*
H8B	0.5456	0.7431	0.7871	0.058*
C9	0.6192 (3)	0.84641 (14)	0.85749 (10)	0.0379 (4)
H9A	0.7299	0.8294	0.8551	0.045*
H9B	0.5736	0.8103	0.8952	0.045*
C10	0.6034 (2)	0.95805 (13)	0.87113 (9)	0.0289 (4)
C11	0.6583 (2)	0.97663 (12)	0.94619 (8)	0.0266 (3)
H11	0.5967	0.9304	0.9737	0.032*
C12	0.8309 (2)	0.94807 (15)	0.95867 (10)	0.0346 (4)
H12A	0.8986	0.9948	0.9355	0.041*
H12B	0.8503	0.8833	0.9394	0.041*
C13	0.8741 (2)	0.94642 (14)	1.03528 (10)	0.0338 (4)
C14	0.7714 (3)	0.86428 (14)	1.06479 (10)	0.0372 (4)
C15	0.6277 (3)	0.89190 (13)	1.10736 (9)	0.0346 (4)
C16	0.5554 (2)	0.99228 (14)	1.08868 (9)	0.0322 (4)
C17	0.8335 (2)	1.04648 (13)	1.06521 (9)	0.0317 (4)
C18	0.6196 (2)	1.16872 (13)	1.09524 (10)	0.0341 (4)
C19	0.5811 (3)	1.23338 (17)	1.20486 (12)	0.0498 (6)
H19A	0.4780	1.2564	1.1928	0.075*
H19B	0.6558	1.2858	1.1989	0.075*
H19C	0.5811	1.2126	1.2514	0.075*
C20	0.7279 (3)	1.16296 (14)	0.94607 (10)	0.0392 (5)
H20A	0.7083	1.2241	0.9692	0.059*
H20B	0.7043	1.1706	0.8987	0.059*
H20C	0.8364	1.1450	0.9513	0.059*
C21	0.1540 (3)	0.96145 (17)	0.81845 (13)	0.0475 (5)
H21A	0.1281	0.9253	0.8588	0.071*
H21B	0.0961	0.9349	0.7808	0.071*
H21C	0.1270	1.0297	0.8245	0.071*
C22	0.3678 (3)	1.01321 (16)	0.74011 (10)	0.0448 (5)
H22A	0.3552	1.0822	0.7498	0.067*
H22B	0.2969	0.9944	0.7044	0.067*
H22C	0.4741	1.0008	0.7261	0.067*
C23	0.7051 (2)	1.01368 (17)	0.81888 (10)	0.0409 (5)
H23A	0.7011	0.9798	0.7761	0.061*
H23B	0.8119	1.0163	0.8346	0.061*
H23C	0.6656	1.0796	0.8134	0.061*
C24	1.0483 (3)	0.91813 (19)	1.04234 (13)	0.0496 (6)
H24A	1.0757	0.9149	1.0895	0.074*
H24B	1.1122	0.9669	1.0203	0.074*
H24C	1.0658	0.8549	1.0216	0.074*
C25	0.5021 (3)	0.81210 (15)	1.10693 (14)	0.0515 (6)
H25A	0.4182	0.8306	1.1369	0.077*
H25B	0.5471	0.7508	1.1218	0.077*
H25C	0.4616	0.8045	1.0618	0.077*
C26	0.9386 (3)	1.10751 (16)	1.09100 (11)	0.0429 (5)
H26A	1.0443	1.0901	1.0915	0.051*
H26B	0.9069	1.1680	1.1087	0.051*

O1	0.5918 (2)	1.24745 (10)	1.07174 (8)	0.0472 (4)
O2	0.6234 (2)	1.15134 (10)	1.16213 (7)	0.0466 (4)
O3	0.2674 (2)	0.78454 (11)	0.77637 (9)	0.0544 (4)
O4	0.8037 (2)	0.77818 (11)	1.05593 (10)	0.0617 (5)
O5	0.6898 (2)	0.90860 (11)	1.17396 (7)	0.0485 (4)
H1O5	0.7163	0.8558	1.1907	0.058*
O6	0.41897 (17)	1.00801 (12)	1.10126 (8)	0.0465 (4)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0316 (9)	0.0236 (7)	0.0311 (8)	-0.0013 (7)	-0.0004 (7)	-0.0008 (6)
C2	0.0322 (9)	0.0238 (7)	0.0295 (8)	0.0008 (7)	-0.0008 (7)	0.0006 (6)
C3	0.0352 (10)	0.0299 (8)	0.0365 (9)	0.0057 (8)	-0.0024 (8)	-0.0049 (7)
C4	0.0348 (9)	0.0282 (8)	0.0375 (9)	0.0073 (8)	-0.0057 (8)	-0.0025 (7)
C5	0.0288 (9)	0.0265 (8)	0.0291 (8)	0.0009 (7)	0.0010 (7)	0.0010 (6)
C6	0.0334 (9)	0.0303 (8)	0.0362 (9)	0.0002 (8)	-0.0034 (8)	-0.0013 (7)
C7	0.0468 (12)	0.0338 (9)	0.0319 (9)	-0.0007 (9)	-0.0064 (9)	-0.0024 (7)
C8	0.0513 (13)	0.0410 (11)	0.0516 (13)	0.0118 (10)	-0.0063 (11)	-0.0160 (10)
C9	0.0407 (11)	0.0352 (9)	0.0378 (10)	0.0111 (9)	-0.0046 (9)	-0.0064 (8)
C10	0.0282 (9)	0.0300 (8)	0.0285 (8)	0.0025 (7)	0.0025 (7)	0.0004 (7)
C11	0.0268 (8)	0.0265 (8)	0.0265 (8)	0.0016 (7)	0.0018 (7)	0.0012 (6)
C12	0.0299 (9)	0.0406 (10)	0.0331 (9)	0.0070 (8)	0.0009 (8)	-0.0012 (8)
C13	0.0309 (9)	0.0345 (9)	0.0360 (9)	0.0056 (8)	-0.0044 (8)	-0.0009 (8)
C14	0.0451 (11)	0.0336 (9)	0.0329 (9)	0.0053 (9)	-0.0056 (9)	0.0001 (7)
C15	0.0445 (11)	0.0279 (8)	0.0313 (9)	-0.0043 (8)	-0.0031 (8)	0.0014 (7)
C16	0.0373 (10)	0.0308 (9)	0.0284 (8)	-0.0032 (8)	0.0007 (8)	-0.0023 (7)
C17	0.0331 (9)	0.0318 (8)	0.0302 (8)	0.0006 (8)	-0.0011 (8)	0.0027 (7)
C18	0.0344 (10)	0.0303 (8)	0.0376 (10)	-0.0015 (8)	-0.0026 (8)	-0.0041 (7)
C19	0.0606 (15)	0.0438 (11)	0.0450 (11)	-0.0037 (11)	0.0107 (11)	-0.0153 (10)
C20	0.0450 (12)	0.0331 (9)	0.0396 (10)	-0.0096 (9)	-0.0007 (9)	0.0069 (8)
C21	0.0339 (11)	0.0461 (11)	0.0624 (14)	-0.0015 (9)	-0.0053 (10)	-0.0104 (10)
C22	0.0533 (13)	0.0453 (11)	0.0358 (10)	0.0002 (10)	-0.0068 (10)	0.0038 (9)
C23	0.0356 (10)	0.0544 (12)	0.0328 (9)	-0.0030 (9)	0.0050 (8)	0.0065 (9)
C24	0.0362 (11)	0.0583 (13)	0.0543 (13)	0.0131 (11)	-0.0111 (10)	-0.0098 (11)
C25	0.0584 (14)	0.0319 (10)	0.0642 (14)	-0.0127 (10)	-0.0048 (12)	0.0048 (10)
C26	0.0375 (11)	0.0417 (11)	0.0496 (12)	-0.0050 (9)	-0.0073 (9)	-0.0005 (9)
O1	0.0638 (10)	0.0288 (6)	0.0490 (8)	0.0075 (7)	-0.0084 (8)	-0.0050 (6)
O2	0.0698 (11)	0.0342 (7)	0.0359 (7)	0.0028 (7)	0.0037 (8)	-0.0064 (6)
O3	0.0614 (11)	0.0355 (8)	0.0664 (10)	-0.0091 (8)	-0.0145 (9)	-0.0019 (7)
O4	0.0825 (13)	0.0312 (7)	0.0716 (11)	0.0123 (8)	0.0137 (10)	-0.0002 (7)
O5	0.0746 (12)	0.0382 (7)	0.0327 (7)	0.0031 (8)	-0.0117 (7)	-0.0005 (6)
O6	0.0388 (8)	0.0467 (8)	0.0539 (9)	-0.0008 (7)	0.0106 (7)	0.0049 (7)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

C1—C17	1.529 (3)	C13—C24	1.536 (3)
C1—C16	1.544 (3)	C14—O4	1.210 (2)

C1—C18	1.545 (2)	C14—C15	1.529 (3)
C1—C2	1.610 (2)	C15—O5	1.433 (2)
C2—C3	1.538 (3)	C15—C25	1.519 (3)
C2—C20	1.546 (3)	C15—C16	1.536 (3)
C2—C11	1.566 (2)	C16—O6	1.205 (2)
C3—C4	1.530 (3)	C17—C26	1.319 (3)
C3—H3A	0.9700	C18—O1	1.186 (2)
C3—H3B	0.9700	C18—O2	1.340 (2)
C4—C5	1.531 (2)	C19—O2	1.440 (2)
C4—H4A	0.9700	C19—H19A	0.9600
C4—H4B	0.9700	C19—H19B	0.9600
C5—C10	1.554 (2)	C19—H19C	0.9600
C5—C6	1.563 (3)	C20—H20A	0.9600
C5—H5	0.9800	C20—H20B	0.9600
C6—C7	1.532 (3)	C20—H20C	0.9600
C6—C21	1.535 (3)	C21—H21A	0.9600
C6—C22	1.544 (3)	C21—H21B	0.9600
C7—O3	1.207 (3)	C21—H21C	0.9600
C7—C8	1.495 (3)	C22—H22A	0.9600
C8—C9	1.536 (3)	C22—H22B	0.9600
C8—H8A	0.9700	C22—H22C	0.9600
C8—H8B	0.9700	C23—H23A	0.9600
C9—C10	1.541 (2)	C23—H23B	0.9600
C9—H9A	0.9700	C23—H23C	0.9600
C9—H9B	0.9700	C24—H24A	0.9600
C10—C23	1.541 (3)	C24—H24B	0.9600
C10—C11	1.572 (2)	C24—H24C	0.9600
C11—C12	1.538 (3)	C25—H25A	0.9600
C11—H11	0.9800	C25—H25B	0.9600
C12—C13	1.554 (3)	C25—H25C	0.9600
C12—H12A	0.9700	C26—H26A	0.9300
C12—H12B	0.9700	C26—H26B	0.9300
C13—C17	1.518 (3)	O5—H1O5	0.8200
C13—C14	1.529 (3)		
C17—C1—C16	110.07 (14)	C17—C13—C24	114.00 (17)
C17—C1—C18	110.75 (15)	C14—C13—C24	109.54 (17)
C16—C1—C18	105.22 (15)	C17—C13—C12	108.14 (15)
C17—C1—C2	108.47 (15)	C14—C13—C12	104.21 (16)
C16—C1—C2	109.79 (14)	C24—C13—C12	108.61 (17)
C18—C1—C2	112.51 (14)	O4—C14—C13	121.1 (2)
C3—C2—C20	109.24 (15)	O4—C14—C15	119.7 (2)
C3—C2—C11	109.27 (15)	C13—C14—C15	119.16 (16)
C20—C2—C11	112.77 (15)	O5—C15—C25	112.11 (17)
C3—C2—C1	111.30 (15)	O5—C15—C14	104.29 (17)
C20—C2—C1	108.39 (14)	C25—C15—C14	112.63 (17)
C11—C2—C1	105.86 (13)	O5—C15—C16	103.14 (14)
C4—C3—C2	112.98 (16)	C25—C15—C16	110.30 (17)

C4—C3—H3A	109.0	C14—C15—C16	113.86 (16)
C2—C3—H3A	109.0	O6—C16—C15	119.46 (18)
C4—C3—H3B	109.0	O6—C16—C1	120.88 (18)
C2—C3—H3B	109.0	C15—C16—C1	119.63 (17)
H3A—C3—H3B	107.8	C26—C17—C13	123.72 (19)
C3—C4—C5	110.50 (15)	C26—C17—C1	123.45 (18)
C3—C4—H4A	109.6	C13—C17—C1	112.61 (16)
C5—C4—H4A	109.6	O1—C18—O2	123.17 (17)
C3—C4—H4B	109.6	O1—C18—C1	127.27 (18)
C5—C4—H4B	109.6	O2—C18—C1	109.54 (15)
H4A—C4—H4B	108.1	O2—C19—H19A	109.5
C4—C5—C10	110.34 (15)	O2—C19—H19B	109.5
C4—C5—C6	113.65 (15)	H19A—C19—H19B	109.5
C10—C5—C6	117.46 (15)	O2—C19—H19C	109.5
C4—C5—H5	104.6	H19A—C19—H19C	109.5
C10—C5—H5	104.6	H19B—C19—H19C	109.5
C6—C5—H5	104.6	C2—C20—H20A	109.5
C7—C6—C21	108.47 (17)	C2—C20—H20B	109.5
C7—C6—C22	108.08 (17)	H20A—C20—H20B	109.5
C21—C6—C22	107.68 (18)	C2—C20—H20C	109.5
C7—C6—C5	107.91 (15)	H20A—C20—H20C	109.5
C21—C6—C5	109.53 (17)	H20B—C20—H20C	109.5
C22—C6—C5	114.99 (16)	C6—C21—H21A	109.5
O3—C7—C8	120.95 (19)	C6—C21—H21B	109.5
O3—C7—C6	122.0 (2)	H21A—C21—H21B	109.5
C8—C7—C6	117.03 (18)	C6—C21—H21C	109.5
C7—C8—C9	112.26 (18)	H21A—C21—H21C	109.5
C7—C8—H8A	109.2	H21B—C21—H21C	109.5
C9—C8—H8A	109.2	C6—C22—H22A	109.5
C7—C8—H8B	109.2	C6—C22—H22B	109.5
C9—C8—H8B	109.2	H22A—C22—H22B	109.5
H8A—C8—H8B	107.9	C6—C22—H22C	109.5
C8—C9—C10	112.73 (17)	H22A—C22—H22C	109.5
C8—C9—H9A	109.0	H22B—C22—H22C	109.5
C10—C9—H9A	109.0	C10—C23—H23A	109.5
C8—C9—H9B	109.0	C10—C23—H23B	109.5
C10—C9—H9B	109.0	H23A—C23—H23B	109.5
H9A—C9—H9B	107.8	C10—C23—H23C	109.5
C23—C10—C9	108.29 (16)	H23A—C23—H23C	109.5
C23—C10—C5	114.92 (15)	H23B—C23—H23C	109.5
C9—C10—C5	107.32 (16)	C13—C24—H24A	109.5
C23—C10—C11	112.61 (16)	C13—C24—H24B	109.5
C9—C10—C11	107.17 (14)	H24A—C24—H24B	109.5
C5—C10—C11	106.16 (14)	C13—C24—H24C	109.5
C12—C11—C2	110.55 (15)	H24A—C24—H24C	109.5
C12—C11—C10	113.19 (15)	H24B—C24—H24C	109.5
C2—C11—C10	116.54 (14)	C15—C25—H25A	109.5
C12—C11—H11	105.1	C15—C25—H25B	109.5

C2—C11—H11	105.1	H25A—C25—H25B	109.5
C10—C11—H11	105.1	C15—C25—H25C	109.5
C11—C12—C13	112.61 (15)	H25A—C25—H25C	109.5
C11—C12—H12A	109.1	H25B—C25—H25C	109.5
C13—C12—H12A	109.1	C17—C26—H26A	120.0
C11—C12—H12B	109.1	C17—C26—H26B	120.0
C13—C12—H12B	109.1	H26A—C26—H26B	120.0
H12A—C12—H12B	107.8	C18—O2—C19	115.72 (16)
C17—C13—C14	111.82 (16)	C15—O5—H1O5	109.5
C17—C1—C2—C3	179.65 (14)	C5—C10—C11—C2	−54.13 (19)
C16—C1—C2—C3	59.35 (19)	C2—C11—C12—C13	58.8 (2)
C18—C1—C2—C3	−57.47 (19)	C10—C11—C12—C13	−168.36 (15)
C17—C1—C2—C20	−60.19 (18)	C11—C12—C13—C17	−55.3 (2)
C16—C1—C2—C20	179.50 (15)	C11—C12—C13—C14	63.8 (2)
C18—C1—C2—C20	62.69 (19)	C11—C12—C13—C24	−179.52 (17)
C17—C1—C2—C11	61.04 (17)	C17—C13—C14—O4	−167.3 (2)
C16—C1—C2—C11	−59.27 (18)	C24—C13—C14—O4	−40.0 (3)
C18—C1—C2—C11	−176.09 (15)	C12—C13—C14—O4	76.1 (2)
C20—C2—C3—C4	74.34 (19)	C17—C13—C14—C15	11.4 (2)
C11—C2—C3—C4	−49.46 (19)	C24—C13—C14—C15	138.78 (18)
C1—C2—C3—C4	−166.01 (14)	C12—C13—C14—C15	−105.16 (19)
C2—C3—C4—C5	58.4 (2)	O4—C14—C15—O5	95.3 (2)
C3—C4—C5—C10	−63.8 (2)	C13—C14—C15—O5	−83.44 (19)
C3—C4—C5—C6	161.82 (16)	O4—C14—C15—C25	−26.5 (3)
C4—C5—C6—C7	179.97 (16)	C13—C14—C15—C25	154.76 (18)
C10—C5—C6—C7	49.0 (2)	O4—C14—C15—C16	−153.0 (2)
C4—C5—C6—C21	−62.1 (2)	C13—C14—C15—C16	28.2 (2)
C10—C5—C6—C21	166.93 (17)	O5—C15—C16—O6	−91.8 (2)
C4—C5—C6—C22	59.3 (2)	C25—C15—C16—O6	28.1 (3)
C10—C5—C6—C22	−71.7 (2)	C14—C15—C16—O6	155.85 (18)
C21—C6—C7—O3	15.0 (3)	O5—C15—C16—C1	86.0 (2)
C22—C6—C7—O3	−101.4 (2)	C25—C15—C16—C1	−154.06 (18)
C5—C6—C7—O3	133.6 (2)	C14—C15—C16—C1	−26.3 (2)
C21—C6—C7—C8	−165.2 (2)	C17—C1—C16—O6	163.11 (18)
C22—C6—C7—C8	78.4 (2)	C18—C1—C16—O6	43.8 (2)
C5—C6—C7—C8	−46.6 (2)	C2—C1—C16—O6	−77.6 (2)
O3—C7—C8—C9	−128.7 (2)	C17—C1—C16—C15	−14.7 (2)
C6—C7—C8—C9	51.5 (3)	C18—C1—C16—C15	−134.06 (16)
C7—C8—C9—C10	−55.0 (3)	C2—C1—C16—C15	104.64 (18)
C8—C9—C10—C23	−70.2 (2)	C14—C13—C17—C26	129.3 (2)
C8—C9—C10—C5	54.4 (2)	C24—C13—C17—C26	4.4 (3)
C8—C9—C10—C11	168.11 (17)	C12—C13—C17—C26	−116.5 (2)
C4—C5—C10—C23	−65.9 (2)	C14—C13—C17—C1	−55.9 (2)
C6—C5—C10—C23	66.6 (2)	C24—C13—C17—C1	179.19 (18)
C4—C5—C10—C9	173.62 (15)	C12—C13—C17—C1	58.3 (2)
C6—C5—C10—C9	−53.9 (2)	C16—C1—C17—C26	−128.2 (2)
C4—C5—C10—C11	59.28 (18)	C18—C1—C17—C26	−12.3 (3)

C6—C5—C10—C11	−168.29 (14)	C2—C1—C17—C26	111.7 (2)
C3—C2—C11—C12	−179.43 (15)	C16—C1—C17—C13	57.01 (19)
C20—C2—C11—C12	58.88 (19)	C18—C1—C17—C13	172.94 (15)
C1—C2—C11—C12	−59.47 (17)	C2—C1—C17—C13	−63.13 (18)
C3—C2—C11—C10	49.48 (19)	C17—C1—C18—O1	108.2 (2)
C20—C2—C11—C10	−72.2 (2)	C16—C1—C18—O1	−132.9 (2)
C1—C2—C11—C10	169.43 (14)	C2—C1—C18—O1	−13.4 (3)
C23—C10—C11—C12	−57.4 (2)	C17—C1—C18—O2	−70.5 (2)
C9—C10—C11—C12	61.6 (2)	C16—C1—C18—O2	48.4 (2)
C5—C10—C11—C12	176.01 (14)	C2—C1—C18—O2	167.95 (16)
C23—C10—C11—C2	72.4 (2)	O1—C18—O2—C19	4.0 (3)
C9—C10—C11—C2	−168.58 (16)	C1—C18—O2—C19	−177.29 (18)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
O5—H1O5···O3 <sup>i</sup>	0.82	2.05	2.870 (2)	173
C23—H23A···O5 <sup>ii</sup>	0.96	2.68	3.173 (2)	112

Symmetry codes: (i)  $x+1/2, -y+3/2, -z+2$ ; (ii)  $-x+3/2, -y+2, z-1/2$ .