

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Hemi(4,4'-bipyridinium) hexafluoridophosphate bis(4-aminobenzoic acid) 4,4'-bipyridine monohydrate

Yi-Yi Wu,^a Chun-De Huang,^a Jie-Xuan Huang,^a Rong-Hua Zeng^{a,b*} and Yi-Fan Luo^a

^aSchool of Chemistry and Environment, South China Normal University, Guangzhou 510006, People's Republic of China, and ^bKey Laboratory of Electrochemical Technology on Energy Storage and Power Generation in Guangdong Universities, Guangzhou 510631, People's Republic of China
Correspondence e-mail: zrh321@yahoo.com.cn


Received 11 December 2008; accepted 25 December 2008

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; disorder in main residue; R factor = 0.066; wR factor = 0.205; data-to-parameter ratio = 15.7.

In the title compound, $0.5\text{C}_{10}\text{H}_{10}\text{N}_2^{2+} \cdot \text{PF}_6^- \cdot \text{C}_{10}\text{H}_8\text{N}_2 \cdot 2\text{C}_7\text{H}_7\text{NO}_2 \cdot \text{H}_2\text{O}$, the cation is located on a center of symmetry. The crystal structure is determined by a complex three-dimensional network of intermolecular $\text{O}-\text{H} \cdots \text{O}$, $\text{O}-\text{H} \cdots \text{N}$, $\text{N}-\text{H} \cdots \text{N}$ and $\text{N}-\text{H} \cdots \text{F}$ hydrogen bonds. $\pi-\pi$ stacking interactions between neighboring pyridyl rings are also present; the centroid-centroid distance is $3.643(5)$ Å. The hexafluoridophosphate anion is disordered over two positions with site-occupancy factors of *ca* 0.6 and 0.4.

Related literature

For the use of 4-aminobenzoic acid and 4,4'-bipyridine for the construction of three-dimensional network motifs, see: Hu *et al.* (2003); Yang *et al.* (2004).



Experimental

Crystal data

$0.5\text{C}_{10}\text{H}_{10}\text{N}_2^{2+} \cdot \text{PF}_6^- \cdot \text{C}_{10}\text{H}_8\text{N}_2 \cdot 2\text{C}_7\text{H}_7\text{NO}_2 \cdot \text{H}_2\text{O}$
 $M_r = 672.54$
 Triclinic, $P\bar{1}$
 $a = 10.1032(2)$ Å
 $b = 10.1142(2)$ Å
 $c = 16.8906(3)$ Å
 $\alpha = 92.557(1)^\circ$

$\beta = 98.063(1)^\circ$
 $\gamma = 117.346(1)^\circ$
 $V = 1506.23(5)$ Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.18$ mm⁻¹
 $T = 296(2)$ K
 $0.18 \times 0.15 \times 0.14$ mm

Data collection

Bruker APEXII area-detector diffractometer
 Absorption correction: none
 22618 measured reflections

7121 independent reflections
 3129 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.057$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.066$
 $wR(F^2) = 0.205$
 $S = 1.02$
 7121 reflections
 454 parameters

43 restraints
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.29$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.34$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{O1W}-\text{H2W} \cdots \text{O4}^i$	0.89	1.87	2.751 (3)	169
$\text{O1W}-\text{H1W} \cdots \text{O2}^{ii}$	0.79	2.01	2.799 (3)	174
$\text{O3}-\text{H3A} \cdots \text{N3}^{iii}$	0.82	1.87	2.686 (3)	174
$\text{O1}-\text{H1} \cdots \text{O1W}^{iv}$	0.82	1.80	2.617 (3)	173
$\text{N1}-\text{H1B} \cdots \text{F3}$	0.86	2.57	3.324 (4)	147
$\text{N5}-\text{H27} \cdots \text{N4}$	0.86	1.84	2.700 (4)	176

Symmetry codes: (i) $x-1, y, z$; (ii) $-x, -y+1, -z+1$; (iii) $x+1, y, z-1$; (iv) $x, y, z+1$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors acknowledge South China Normal University for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WN2299).

References

- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Hu, D. H., Huang, W., Gou, S. H., Fang, J. L. & Fun, H. K. (2003). *Polyhedron*, **22**, 2661–2667.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Yang, G. P., Wang, Z. Y. & Chen, J. T. (2004). *J. Mol. Struct.* **707**, 223–229.

supplementary materials

Acta Cryst. (2009). E65, o278 [doi:10.1107/S1600536808043900]

Hemi(4,4'-bipyridinium) hexafluoridophosphate bis(4-aminobenzoic acid) 4,4'-bipyridine monohydrate

Y.-Y. Wu, C.-D. Huang, J.-X. Huang, R.-H. Zeng and Y.-F. Luo

Comment

Hydrogen-bonding interactions between ligands are specific and directional. In this context, 4-aminobenzoic acid and 4,4'-bipyridine are excellent candidates for the construction of three-dimensional network motifs, which form regular hydrogen bonds, functioning as both hydrogen-bond donor and acceptor (Hu *et al.*, 2003; Yang *et al.*, 2004). Recently, we obtained the title compound under hydrothermal conditions and report its crystal structure here.

In the title compound (Fig. 1), all bond lengths and angles are unexceptional. The 4,4'-bipyridinium cation, which is located on a center of symmetry, is, on both sides, a hydrogen bond donor to unprotonated molecules of 4,4'-bipyridine. These are, in turn, hydrogen bond acceptors for one of the two independent molecules of 4-aminobenzoic acid. Thus the two 4-aminobenzoic acid molecules, the 4,4'-bipyridine and 4,4'-bipyridinium cation are connected by hydrogen bonding interactions to form a linear centrosymmetric chain. Further each of these chains is connected with each of two other chains by O—H...O hydrogen bonds (Table 1), *via* the interstitial solvent water molecules and the other independent 4-aminobenzoic acid molecules, forming, in effect, infinite chains with each repeating unit of five molecules offset against the next unit by the width of the connecting $(C_7NH_7O_2)_2 \cdot (H_2O)_2$ unit. π - π stacking interactions (the centroid-centroid distance between neighboring pyridyl rings is 3.643 Å) connect these chains to produce a three-dimensional network motif (Fig. 2).

Experimental

4-Aminobenzoic acid (1 mmol, 0.137 g), 4,4'-bipyridine (1 mmol, 0.156 g) and sodium hexafluoridophosphate (1 mmol, 0.162 g) were dissolved in hot water with stirring. Colorless single crystals were obtained at room temperature by slow evaporation of the solvent over a period of several days.

Refinement

The disordered perchlorate ion was refined over two sites, with refined occupancies of 0.625 (8) and 0.375 (9). The P...F and F...F distances were restrained to be 1.48 (1) and 2.10 (3) Å, respectively. Water H atoms were located in a difference Fourier map and refined with distance restraints of O—H = 0.86 Å and H...H = 1.39 Å. The H atom bound to the N5 nitrogen atom in the cation and the carboxylic H atoms were refined with distance restraints of N—H = 0.90 Å and O—H = 0.90 Å, respectively. All other H atoms were placed at calculated positions and treated as riding on the parent atoms, with C—H = 0.93 Å, N—H = 0.86 Å, and with $U_{iso}(H) = 1.2U_{eq}(C, N)$.

Figures




Fig. 1. The molecular structure showing the atomic-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. Hydrogen atoms are shown as spheres of arbitrary radius. Symmetry code: (i) $2 - x, -y, 1 - z$.




Fig. 2. A view of the three-dimensional supramolecular network. Hydrogen bonds are shown as dashed lines.

Hemi(4,4'-bipyridinium) hexafluoridophosphate bis(4-aminobenzoic acid) 4,4'-bipyridine monohydrate

Crystal data

$0.5\text{C}_{10}\text{H}_{10}\text{N}_2^{2+} \cdot \text{P}_1\text{F}_6^- \cdot \text{C}_{10}\text{H}_8\text{N}_2 \cdot 2\text{C}_7\text{H}_7\text{NO}_2 \cdot \text{H}_2\text{O}_1$	$Z = 2$
$M_r = 672.54$	$F_{000} = 694$
Triclinic, $P\bar{1}$	$D_x = 1.483 \text{ Mg m}^{-3}$
Hall symbol: $-P\ 1$	Mo $K\alpha$ radiation
$a = 10.1032 (2) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 10.1142 (2) \text{ \AA}$	Cell parameters from 2023 reflections
$c = 16.8906 (3) \text{ \AA}$	$\theta = 2.4\text{--}19.5^\circ$
$\alpha = 92.557 (1)^\circ$	$\mu = 0.18 \text{ mm}^{-1}$
$\beta = 98.063 (1)^\circ$	$T = 296 (2) \text{ K}$
$\gamma = 117.346 (1)^\circ$	Block, colourless
$V = 1506.23 (5) \text{ \AA}^3$	$0.18 \times 0.15 \times 0.14 \text{ mm}$

Data collection

Bruker APEXII area-detector diffractometer	3129 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.057$
Monochromator: graphite	$\theta_{\text{max}} = 27.9^\circ$
$T = 296(2) \text{ K}$	$\theta_{\text{min}} = 2.3^\circ$
φ and ω scans	$h = -13 \rightarrow 13$
Absorption correction: none	$k = -13 \rightarrow 13$
22618 measured reflections	$l = -20 \rightarrow 22$
7121 independent reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites

$$R[F^2 > 2\sigma(F^2)] = 0.066$$

$$wR(F^2) = 0.205$$

$$S = 1.02$$

7121 reflections

454 parameters

43 restraints

Primary atom site location: structure-invariant direct methods

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0875P)^2 + 0.1722P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.006$$

$$\Delta\rho_{\max} = 0.29 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.33 \text{ e } \text{\AA}^{-3}$$

Extinction correction: SHELXL97 (Sheldrick, 2008)

Extinction coefficient: 0.009 (2)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.4034 (4)	0.5337 (4)	0.7188 (2)	0.0650 (9)	
C2	0.4778 (4)	0.5250 (4)	0.7922 (2)	0.0656 (9)	
H2	0.5740	0.5331	0.7960	0.079*	
C3	0.4122 (4)	0.5047 (4)	0.8589 (2)	0.0602 (9)	
H3	0.4645	0.4991	0.9075	0.072*	
C4	0.2682 (3)	0.4922 (3)	0.8557 (2)	0.0523 (8)	
C5	0.1957 (4)	0.4726 (3)	0.9266 (2)	0.0576 (8)	
C6	0.1930 (4)	0.4994 (3)	0.7822 (2)	0.0590 (8)	
H6	0.0961	0.4896	0.7783	0.071*	
C7	0.2589 (4)	0.5208 (4)	0.7152 (2)	0.0646 (9)	
H7	0.2068	0.5267	0.6666	0.077*	
C8	0.8336 (4)	0.1574 (4)	0.3481 (2)	0.0662 (9)	
H8	0.8482	0.2431	0.3790	0.079*	
C9	0.9268 (4)	0.1700 (4)	0.2940 (2)	0.0627 (9)	
H9	1.0027	0.2642	0.2880	0.075*	
C10	0.9095 (3)	0.0437 (3)	0.24786 (18)	0.0503 (7)	
C11	1.0119 (4)	0.0620 (4)	0.19048 (19)	0.0555 (8)	
C12	0.7954 (3)	-0.0956 (4)	0.25815 (19)	0.0545 (8)	
H12	0.7825	-0.1815	0.2283	0.065*	
C13	0.7011 (4)	-0.1079 (4)	0.31200 (19)	0.0579 (8)	
H13	0.6254	-0.2022	0.3180	0.070*	
C14	0.7174 (4)	0.0183 (4)	0.35751 (19)	0.0578 (8)	
N3	0.1693 (3)	-0.0235 (3)	1.04268 (16)	0.0592 (7)	

supplementary materials

C16	0.2657 (5)	0.1119 (4)	1.0271 (2)	0.0754 (11)	
H16	0.2693	0.1953	1.0549	0.090*	
C17	0.3601 (4)	0.1349 (4)	0.9723 (2)	0.0682 (10)	
H17	0.4246	0.2321	0.9634	0.082*	
C18	0.3597 (3)	0.0145 (3)	0.93018 (17)	0.0464 (7)	
C19	0.2589 (3)	-0.1253 (4)	0.9462 (2)	0.0606 (9)	
H19	0.2522	-0.2109	0.9192	0.073*	
C20	0.1679 (4)	-0.1388 (4)	1.0023 (2)	0.0650 (9)	
H20	0.1018	-0.2346	1.0122	0.078*	
C21	0.4624 (3)	0.0328 (3)	0.87181 (17)	0.0452 (7)	
C22	0.5692 (4)	0.1715 (4)	0.8566 (2)	0.0641 (9)	
H22	0.5793	0.2585	0.8839	0.077*	
C23	0.6600 (4)	0.1817 (4)	0.8017 (2)	0.0662 (9)	
H23	0.7302	0.2765	0.7925	0.079*	
N4	0.6522 (3)	0.0623 (3)	0.76098 (15)	0.0576 (7)	
C25	0.5513 (5)	-0.0691 (4)	0.7747 (2)	0.0791 (11)	
H25	0.5432	-0.1540	0.7460	0.095*	
C26	0.4564 (4)	-0.0888 (4)	0.8291 (2)	0.0734 (11)	
H26	0.3877	-0.1852	0.8369	0.088*	
N5	0.8058 (3)	0.0432 (3)	0.64624 (16)	0.0600 (7)	
H27	0.7588	0.0538	0.6827	0.072*	
C28	0.9248 (4)	0.1591 (4)	0.6303 (2)	0.0761 (11)	
H28	0.9572	0.2528	0.6586	0.091*	
C29	1.0029 (4)	0.1461 (4)	0.5730 (2)	0.0719 (10)	
H29	1.0865	0.2306	0.5628	0.086*	
C30	0.9587 (3)	0.0097 (3)	0.53068 (17)	0.0481 (7)	
C31	0.8329 (5)	-0.1090 (4)	0.5490 (3)	0.0920 (14)	
H31	0.7977	-0.2044	0.5220	0.110*	
C32	0.7592 (5)	-0.0883 (5)	0.6064 (3)	0.0932 (13)	
H32	0.6736	-0.1700	0.6172	0.112*	
F3	0.4396 (3)	0.6418 (3)	0.46460 (18)	0.1343 (10)	
F6	0.0964 (3)	0.5330 (3)	0.3961 (2)	0.1465 (12)	
N1	0.4710 (4)	0.5607 (4)	0.6526 (2)	0.0960 (11)	
H1A	0.5617	0.5727	0.6560	0.115*	
H1B	0.4230	0.5658	0.6076	0.115*	
N2	0.6215 (3)	0.0068 (4)	0.41031 (18)	0.0812 (9)	
H2A	0.5499	-0.0797	0.4156	0.097*	
H2B	0.6331	0.0860	0.4380	0.097*	
O1	0.2766 (3)	0.4659 (3)	0.99380 (15)	0.0756 (7)	
H1	0.2309	0.4589	1.0312	0.113*	
O2	0.0709 (3)	0.4645 (3)	0.92571 (15)	0.0767 (7)	
O3	0.9955 (3)	-0.0651 (3)	0.15538 (15)	0.0728 (7)	
H3A	1.0465	-0.0476	0.1197	0.109*	
O4	1.1065 (3)	0.1827 (3)	0.17604 (15)	0.0772 (7)	
P	0.26740 (12)	0.58756 (12)	0.43141 (7)	0.0774 (4)	
O1W	0.1528 (2)	0.4502 (3)	0.12153 (13)	0.0703 (7)	
H1W	0.0862	0.4708	0.1110	0.105*	
H2W	0.1255	0.3604	0.1373	0.105*	
F1	0.2698 (15)	0.7215 (12)	0.4737 (9)	0.129 (5)	0.375 (11)

F2	0.2105 (13)	0.4994 (17)	0.5008 (6)	0.125 (5)	0.375 (11)
F4	0.2560 (8)	0.4439 (8)	0.3880 (8)	0.090 (4)	0.375 (11)
F5	0.3176 (16)	0.6648 (13)	0.3621 (5)	0.139 (5)	0.375 (11)
F1'	0.3113 (7)	0.7452 (10)	0.4047 (10)	0.192 (6)	0.625 (11)
F4'	0.2424 (12)	0.4441 (8)	0.4616 (10)	0.222 (6)	0.625 (11)
F5'	0.2826 (14)	0.534 (2)	0.3497 (4)	0.239 (7)	0.625 (11)
F2'	0.2667 (13)	0.6511 (18)	0.5141 (5)	0.217 (5)	0.625 (11)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.070 (2)	0.051 (2)	0.073 (2)	0.0229 (18)	0.030 (2)	0.0092 (17)
C2	0.056 (2)	0.062 (2)	0.084 (3)	0.0283 (18)	0.0231 (19)	0.0113 (19)
C3	0.0564 (19)	0.055 (2)	0.070 (2)	0.0254 (16)	0.0149 (17)	0.0129 (17)
C4	0.0520 (18)	0.0403 (18)	0.064 (2)	0.0199 (15)	0.0164 (16)	0.0088 (15)
C5	0.059 (2)	0.0442 (19)	0.069 (2)	0.0215 (16)	0.0211 (18)	0.0112 (16)
C6	0.0518 (18)	0.051 (2)	0.070 (2)	0.0217 (16)	0.0105 (17)	0.0028 (17)
C7	0.068 (2)	0.057 (2)	0.067 (2)	0.0268 (18)	0.0146 (18)	0.0086 (17)
C8	0.071 (2)	0.056 (2)	0.072 (2)	0.0297 (19)	0.0200 (19)	-0.0035 (18)
C9	0.062 (2)	0.049 (2)	0.072 (2)	0.0206 (17)	0.0181 (18)	0.0064 (17)
C10	0.0495 (17)	0.051 (2)	0.0510 (18)	0.0224 (15)	0.0134 (14)	0.0095 (15)
C11	0.0519 (18)	0.061 (2)	0.055 (2)	0.0264 (18)	0.0116 (16)	0.0127 (17)
C12	0.0565 (19)	0.0490 (19)	0.057 (2)	0.0225 (16)	0.0166 (16)	0.0052 (15)
C13	0.0565 (19)	0.049 (2)	0.061 (2)	0.0174 (16)	0.0175 (16)	0.0089 (16)
C14	0.0563 (19)	0.062 (2)	0.056 (2)	0.0272 (18)	0.0155 (16)	0.0050 (17)
N3	0.0575 (16)	0.069 (2)	0.0549 (17)	0.0288 (15)	0.0229 (13)	0.0115 (15)
C16	0.099 (3)	0.066 (3)	0.080 (3)	0.045 (2)	0.045 (2)	0.014 (2)
C17	0.085 (3)	0.048 (2)	0.080 (2)	0.0299 (19)	0.045 (2)	0.0172 (18)
C18	0.0454 (16)	0.0493 (19)	0.0436 (17)	0.0213 (15)	0.0086 (13)	0.0071 (14)
C19	0.0590 (19)	0.046 (2)	0.070 (2)	0.0144 (16)	0.0276 (17)	0.0035 (16)
C20	0.056 (2)	0.061 (2)	0.068 (2)	0.0151 (17)	0.0243 (17)	0.0088 (18)
C21	0.0466 (16)	0.0461 (18)	0.0424 (17)	0.0200 (14)	0.0114 (13)	0.0086 (13)
C22	0.072 (2)	0.047 (2)	0.068 (2)	0.0196 (17)	0.0305 (18)	0.0026 (16)
C23	0.067 (2)	0.053 (2)	0.068 (2)	0.0147 (17)	0.0289 (18)	0.0063 (18)
N4	0.0616 (16)	0.0588 (18)	0.0543 (17)	0.0268 (15)	0.0213 (13)	0.0086 (14)
C25	0.105 (3)	0.052 (2)	0.089 (3)	0.034 (2)	0.053 (2)	0.008 (2)
C26	0.089 (3)	0.042 (2)	0.088 (3)	0.0195 (18)	0.054 (2)	0.0121 (18)
N5	0.0648 (18)	0.070 (2)	0.0529 (16)	0.0334 (16)	0.0259 (14)	0.0092 (15)
C28	0.069 (2)	0.063 (2)	0.087 (3)	0.021 (2)	0.030 (2)	-0.011 (2)
C29	0.060 (2)	0.056 (2)	0.085 (3)	0.0109 (17)	0.0320 (19)	-0.0064 (19)
C30	0.0546 (18)	0.0451 (18)	0.0462 (17)	0.0219 (15)	0.0172 (14)	0.0127 (15)
C31	0.122 (3)	0.048 (2)	0.104 (3)	0.023 (2)	0.078 (3)	0.008 (2)
C32	0.110 (3)	0.060 (3)	0.104 (3)	0.022 (2)	0.069 (3)	0.013 (2)
F3	0.0937 (19)	0.140 (3)	0.145 (3)	0.0434 (17)	-0.0065 (17)	-0.0012 (19)
F6	0.0795 (18)	0.118 (2)	0.226 (4)	0.0400 (16)	0.007 (2)	0.001 (2)
N1	0.104 (3)	0.109 (3)	0.090 (3)	0.052 (2)	0.051 (2)	0.029 (2)
N2	0.079 (2)	0.077 (2)	0.085 (2)	0.0292 (17)	0.0385 (18)	-0.0015 (17)
O1	0.0770 (17)	0.0873 (18)	0.0663 (16)	0.0383 (15)	0.0222 (13)	0.0198 (14)

supplementary materials

O2	0.0705 (16)	0.0926 (19)	0.0860 (18)	0.0477 (15)	0.0338 (14)	0.0260 (14)
O3	0.0778 (17)	0.0681 (17)	0.0758 (17)	0.0300 (13)	0.0393 (13)	0.0101 (13)
O4	0.0750 (16)	0.0637 (17)	0.0926 (19)	0.0245 (14)	0.0391 (14)	0.0237 (14)
P	0.0796 (7)	0.0646 (7)	0.0843 (8)	0.0286 (6)	0.0226 (6)	0.0093 (6)
O1W	0.0699 (15)	0.0638 (15)	0.0736 (16)	0.0265 (12)	0.0180 (12)	0.0168 (12)
F1	0.171 (7)	0.073 (6)	0.130 (8)	0.054 (5)	0.012 (6)	-0.041 (5)
F2	0.122 (6)	0.140 (9)	0.098 (6)	0.035 (6)	0.057 (4)	0.047 (6)
F4	0.067 (4)	0.062 (5)	0.135 (8)	0.031 (3)	0.009 (5)	-0.014 (4)
F5	0.193 (8)	0.108 (8)	0.080 (6)	0.026 (6)	0.071 (5)	0.038 (5)
F1'	0.100 (4)	0.124 (7)	0.341 (14)	0.033 (4)	0.045 (6)	0.138 (8)
F4'	0.223 (9)	0.082 (5)	0.352 (18)	0.066 (5)	0.023 (10)	0.089 (8)
F5'	0.241 (11)	0.45 (2)	0.085 (5)	0.228 (14)	-0.006 (5)	-0.058 (9)
F2'	0.294 (11)	0.263 (14)	0.141 (7)	0.164 (11)	0.081 (7)	-0.032 (8)

Geometric parameters (Å, °)

C1—N1	1.364 (4)	C19—C20	1.380 (4)
C1—C2	1.387 (5)	C19—H19	0.9300
C1—C7	1.396 (5)	C20—H20	0.9300
C2—C3	1.361 (5)	C21—C26	1.370 (4)
C2—H2	0.9300	C21—C22	1.385 (4)
C3—C4	1.394 (4)	C22—C23	1.368 (4)
C3—H3	0.9300	C22—H22	0.9300
C4—C6	1.385 (4)	C23—N4	1.327 (4)
C4—C5	1.465 (4)	C23—H23	0.9300
C5—O2	1.223 (4)	N4—C25	1.306 (4)
C5—O1	1.325 (4)	C25—C26	1.376 (5)
C6—C7	1.368 (4)	C25—H25	0.9300
C6—H6	0.9300	C26—H26	0.9300
C7—H7	0.9300	N5—C32	1.304 (4)
C8—C9	1.372 (5)	N5—C28	1.310 (4)
C8—C14	1.392 (5)	N5—H27	0.8600
C8—H8	0.9300	C28—C29	1.369 (5)
C9—C10	1.391 (4)	C28—H28	0.9300
C9—H9	0.9300	C29—C30	1.367 (4)
C10—C12	1.389 (4)	C29—H29	0.9300
C10—C11	1.472 (4)	C30—C31	1.377 (5)
C11—O4	1.218 (4)	C30—C30 ⁱ	1.477 (6)
C11—O3	1.318 (4)	C31—C32	1.366 (5)
C12—C13	1.376 (4)	C31—H31	0.9300
C12—H12	0.9300	C32—H32	0.9300
C13—C14	1.390 (4)	F3—P	1.570 (3)
C13—H13	0.9300	F6—P	1.566 (3)
C14—N2	1.377 (4)	N1—H1A	0.8600
N3—C20	1.317 (4)	N1—H1B	0.8600
N3—C16	1.331 (4)	N2—H2A	0.8600
C16—C17	1.373 (5)	N2—H2B	0.8600
C16—H16	0.9300	O1—H1	0.8200
C17—C18	1.381 (4)	O3—H3A	0.8200

C17—H17	0.9300	O1W—H1W	0.7910
C18—C19	1.378 (4)	O1W—H2W	0.8864
C18—C21	1.487 (4)		
N1—C1—C2	121.3 (3)	C19—C18—C21	121.5 (3)
N1—C1—C7	120.6 (4)	C17—C18—C21	122.6 (3)
C2—C1—C7	118.0 (3)	C18—C19—C20	120.2 (3)
C3—C2—C1	121.1 (3)	C18—C19—H19	119.9
C3—C2—H2	119.5	C20—C19—H19	119.9
C1—C2—H2	119.5	N3—C20—C19	123.7 (3)
C2—C3—C4	121.2 (3)	N3—C20—H20	118.1
C2—C3—H3	119.4	C19—C20—H20	118.2
C4—C3—H3	119.4	C26—C21—C22	115.6 (3)
C6—C4—C3	117.9 (3)	C26—C21—C18	121.4 (3)
C6—C4—C5	119.7 (3)	C22—C21—C18	123.0 (3)
C3—C4—C5	122.4 (3)	C23—C22—C21	120.5 (3)
O2—C5—O1	121.4 (3)	C23—C22—H22	119.7
O2—C5—C4	123.7 (3)	C21—C22—H22	119.7
O1—C5—C4	115.0 (3)	N4—C23—C22	122.8 (3)
C7—C6—C4	121.1 (3)	N4—C23—H23	118.6
C7—C6—H6	119.5	C22—C23—H23	118.6
C4—C6—H6	119.5	C25—N4—C23	117.1 (3)
C6—C7—C1	120.8 (3)	N4—C25—C26	123.6 (3)
C6—C7—H7	119.6	N4—C25—H25	118.2
C1—C7—H7	119.6	C26—C25—H25	118.2
C9—C8—C14	120.9 (3)	C21—C26—C25	120.4 (3)
C9—C8—H8	119.5	C21—C26—H26	119.8
C14—C8—H8	119.5	C25—C26—H26	119.8
C8—C9—C10	120.9 (3)	C32—N5—C28	119.5 (3)
C8—C9—H9	119.5	C32—N5—H27	120.3
C10—C9—H9	119.5	C28—N5—H27	120.3
C12—C10—C9	118.3 (3)	N5—C28—C29	121.8 (3)
C12—C10—C11	122.5 (3)	N5—C28—H28	119.1
C9—C10—C11	119.2 (3)	C29—C28—H28	119.1
O4—C11—O3	121.3 (3)	C30—C29—C28	120.5 (3)
O4—C11—C10	124.2 (3)	C30—C29—H29	119.8
O3—C11—C10	114.5 (3)	C28—C29—H29	119.8
C13—C12—C10	120.8 (3)	C29—C30—C31	115.9 (3)
C13—C12—H12	119.6	C29—C30—C30 ⁱ	122.2 (3)
C10—C12—H12	119.6	C31—C30—C30 ⁱ	121.8 (4)
C12—C13—C14	121.0 (3)	C32—C31—C30	120.7 (4)
C12—C13—H13	119.5	C32—C31—H31	119.6
C14—C13—H13	119.5	C30—C31—H31	119.6
N2—C14—C13	121.2 (3)	N5—C32—C31	121.6 (4)
N2—C14—C8	120.7 (3)	N5—C32—H32	119.2
C13—C14—C8	118.1 (3)	C31—C32—H32	119.2
C20—N3—C16	116.4 (3)	C1—N1—H1A	120.0
N3—C16—C17	123.5 (3)	C1—N1—H1B	120.0
N3—C16—H16	118.3	H1A—N1—H1B	120.0

supplementary materials

C17—C16—H16	118.3	C14—N2—H2A	120.0
C16—C17—C18	120.3 (3)	C14—N2—H2B	120.0
C16—C17—H17	119.9	H2A—N2—H2B	120.0
C18—C17—H17	119.9	F6—P—F3	178.6 (2)
C19—C18—C17	115.9 (3)	H1W—O1W—H2W	115.1

Symmetry codes: (i) $-x+2, -y, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1W—H2W \cdots O4 ⁱⁱ	0.89	1.87	2.751 (3)	169
O1W—H1W \cdots O2 ⁱⁱⁱ	0.79	2.01	2.799 (3)	174
O3—H3A \cdots N3 ^{iv}	0.82	1.87	2.686 (3)	174
O1—H1 \cdots O1W ^v	0.82	1.80	2.617 (3)	173
N1—H1B \cdots F3	0.86	2.57	3.324 (4)	147
N5—H27 \cdots N4	0.86	1.84	2.700 (4)	176

Symmetry codes: (ii) $x-1, y, z$; (iii) $-x, -y+1, -z+1$; (iv) $x+1, y, z-1$; (v) $x, y, z+1$.

Fig. 1




Fig. 2

