

Diaquabis(1,10-phenanthroline- κ^2N,N')-zinc(II) 2-hydroxy-5-sulfonatobenzoate tetrahydrate

 Kou-Lin Zhang,^a Bo Yang^a and Seik Weng Ng^{b*}
^aCollege of Chemistry and Chemical Engineering, Yangzhou University, Yangzhou 225002, People's Republic of China, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 21 January 2009; accepted 24 January 2009

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; disorder in main residue; R factor = 0.037; wR factor = 0.095; data-to-parameter ratio = 14.3.

The water-coordinated metal centre in the title salt, $[\text{Zn}(\text{C}_{12}\text{H}_8\text{N}_2)_2(\text{H}_2\text{O})_2]\text{C}_7\text{H}_4\text{O}_6\text{S}\cdot 4\text{H}_2\text{O}$, is chelated by the two bidentate N -heterocycles, leading to an overall distorted octahedral environment. The cation, dianion and solvent water molecules interact by $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds to form a layer motif. The SO_3 group is disordered over two positions with respect to the O atoms in a 0.76 (1):0.24 (1) ratio. One of the solvent water molecules is also disordered over two positions in a 0.56 (4):0.44 (4) ratio.

Related literature

For the isostructural manganese(II), nickel(II) and cobalt(II) analogues, see: Fan *et al.* (2005); Chen *et al.* (2005); Zhu & Fan (2005).



Experimental

Crystal data

 $[\text{Zn}(\text{C}_{12}\text{H}_8\text{N}_2)_2(\text{H}_2\text{O})_2]\text{C}_7\text{H}_4\text{O}_6\text{S}\cdot 4\text{H}_2\text{O}$
 $M_r = 750.04$
 Triclinic, $P\bar{1}$
 $a = 10.075$ (1) Å

 $b = 12.263$ (1) Å
 $c = 13.927$ (1) Å
 $\alpha = 96.937$ (2)°
 $\beta = 101.495$ (2)°
 $\gamma = 98.856$ (2)°

 $V = 1645.5$ (2) Å³
 $Z = 2$
 Mo $K\alpha$ radiation

 $\mu = 0.88$ mm⁻¹
 $T = 293$ (2) K
 $0.38 \times 0.30 \times 0.22$ mm

Data collection

 Bruker APEXII area-detector diffractometer
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.36$, $T_{\max} = 0.82$

 10806 measured reflections
 7315 independent reflections
 4849 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.095$
 $S = 0.93$
 7315 reflections
 513 parameters
 106 restraints

 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.28$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.35$ e Å⁻³

Table 1
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1W—H11 ⁱ ··O1	0.83 (1)	1.81 (1)	2.632 (2)	170 (3)
O1W—H12 ⁱ ··O6 ⁱ	0.84 (1)	1.95 (1)	2.793 (4)	177 (2)
O1W—H12 ⁱ ··O6 ^{iv}	0.84 (1)	2.04 (2)	2.787 (7)	147 (2)
O2W—H21··O3W	0.85 (1)	1.87 (1)	2.714 (3)	170 (3)
O2W—H22··O6W	0.85 (1)	1.95 (1)	2.778 (6)	166 (3)
O2W—H22··O6W ⁱ	0.85 (1)	1.80 (2)	2.615 (6)	160 (3)
O3W—H31··O5W	0.85 (1)	1.91 (1)	2.754 (3)	173 (3)
O3W—H32··O5 ⁱⁱ	0.85 (1)	1.95 (1)	2.805 (4)	178 (3)
O3W—H32··O5 ⁱⁱⁱ	0.85 (1)	2.08 (2)	2.892 (10)	159 (3)
O4W—H41··O4 ⁱⁱⁱ	0.86 (1)	2.17 (3)	2.962 (5)	153 (5)
O4W—H41··O4 ^{iv}	0.86 (1)	1.74 (2)	2.598 (8)	173 (5)
O4W—H42··O5 ⁱⁱ	0.86 (1)	2.22 (1)	3.065 (5)	169 (5)
O4W—H42··O5 ⁱⁱⁱ	0.86 (1)	2.08 (2)	2.869 (15)	153 (4)
O5W—H51··O4 ⁱⁱⁱ	0.84 (1)	2.07 (1)	2.900 (4)	168 (4)
O5W—H51··O6 ⁱⁱⁱ	0.84 (1)	2.07 (2)	2.823 (8)	148 (3)
O5W—H52··O2 ^{iv}	0.85 (1)	1.94 (1)	2.792 (3)	175 (3)
O6W—H61··O6 ⁱ	0.83 (1)	2.22 (3)	2.751 (8)	122 (3)
O6W—H62··O4W	0.84 (1)	1.88 (2)	2.642 (7)	150 (4)

Symmetry codes: (i) $-x + 2, -y + 2, -z + 1$; (ii) $x - 1, y, z - 1$; (iii) $-x + 1, -y + 2, -z + 1$; (iv) $x - 1, y, z$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *S SAINT* (Bruker, 2004); data reduction: *S SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *PUBLICIF* (Westrip, 2009).

The authors thank the Fujian Institute for Research on the Structure of Matter for the diffraction measurements, and the Foundation of Jiangsu Provincial Key Programme of Physical Chemistry in Yangzhou University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2475).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.*, **1**, 189–191.
 Bruker (2004). *APEX2* and *S SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Chen, J.-M., Fan, S.-R. & Zhu, L.-G. (2005). *Acta Cryst.* **E61**, m1724–m1726.
 Fan, S.-R., Zhu, L.-G., Xiao, H.-P. & Ng, S. W. (2005). *Acta Cryst.* **E61**, m563–m565.

Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

Westrip, S. P. (2009). *pubCIF*. In preparation.
Zhu, L.-G. & Fan, S. R. (2005). *Chin. J. Chem.* **23**, 1292–1296.

supplementary materials

Acta Cryst. (2009). E65, m239-m240 [doi:10.1107/S1600536809003055]

Diaquabis(1,10-phenanthroline- κ^2N,N')zinc(II) 2-hydroxy-5-sulfonatobenzoate tetrahydrate

K.-L. Zhang, B. Yang and S. W. Ng

Experimental

1,10-Phenanthroline monohydrate (0.10 g, 0.5 mmol) was dissolved in methanol (10 ml). To this solution was added zinc nitrate hexahydrate (0.15 g, 0.5 mmol) dissolved in water (10 ml). The solution was then mixed with an aqueous solution of 5-sulfosalicylic acid (0.11 g, 0.5 mmol) and sodium hydroxide (0.04 g, 1 mmol). Crystals separated after several days. These were collected and washed with methanol; yield: 50%. CH&N elemental analysis: Calc. for $C_{31}H_{32}N_4O_{12}SZn$: C 49.63, H 4.27, N 7.47%. Found: C 49.71, H 4.31, N, 7.41%.

Refinement

The $-SO_3$ group is disordered over two positions with respect to the O atoms. The S—O distances were restrained to 0.01 Å of each other, as were the O...O distances. The anisotropic temperature factors were restrained to be nearly isotropic. The disordered refined to a 0.76 (1):0.24 ratio. One of the lattice water molecules is also disordered over two positions in a 0.56 (4):0.44 ratio. The temperature factors of the two components were restrained to be equal to each other. The anisotropic temperature factors were also restrained to be nearly isotropic.

The carbon-bound H atoms were placed in calculated positions and were allowed to ride on the parent atoms. The oxygen-bound ones were located in a difference Fourier map, and were refined with distance restraints O—H = 0.85 (1) and H...H = 1.39 (1) Å. Their temperature factors were tied by a factor of 1.5.

Figures




Fig. 1. Thermal ellipsoid plot of $[Zn(OH_2)_2(C_{12}H_8N_2)_2][C_7H_4O_6S] \cdot 4H_2O$; displacement ellipsoids are drawn at the 50% probability level, and H atoms as spheres of arbitrary radii. The disorder is not shown.

Diaquabis(1,10-phenanthroline- κ^2N,N')zinc(II) 2-hydroxy-5-sulfonatobenzoate tetrahydrate

Crystal data

$[Zn(C_{12}H_8N_2)_2(H_2O)_2][C_7H_4O_6S] \cdot 4H_2O$

$M_r = 750.04$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$Z = 2$

$F_{000} = 776$

$D_x = 1.514 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073 \text{ \AA}$

supplementary materials

$a = 10.075$ (1) Å	Cell parameters from 2300 reflections
$b = 12.263$ (1) Å	$\theta = 2.5\text{--}22.9^\circ$
$c = 13.927$ (1) Å	$\mu = 0.88$ mm ⁻¹
$\alpha = 96.937$ (2)°	$T = 293$ (2) K
$\beta = 101.495$ (2)°	Block, colourless
$\gamma = 98.856$ (2)°	$0.38 \times 0.30 \times 0.22$ mm
$V = 1645.5$ (2) Å ³	

Data collection

Bruker APEXII area-detector diffractometer	7315 independent reflections
Radiation source: fine-focus sealed tube	4849 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.020$
$T = 293$ (2) K	$\theta_{\text{max}} = 27.5^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.7^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -13 \rightarrow 12$
$T_{\text{min}} = 0.36$, $T_{\text{max}} = 0.82$	$k = -14 \rightarrow 15$
10806 measured reflections	$l = -18 \rightarrow 15$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.037$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.095$	$w = 1/[\sigma^2(F_o^2) + (0.0512P)^2]$
$S = 0.93$	where $P = (F_o^2 + 2F_c^2)/3$
7315 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
513 parameters	$\Delta\rho_{\text{max}} = 0.28$ e Å ⁻³
106 restraints	$\Delta\rho_{\text{min}} = -0.35$ e Å ⁻³
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Zn1	0.50705 (3)	0.73388 (2)	0.26312 (2)	0.03952 (10)	
S1	1.09039 (9)	0.91846 (6)	0.82131 (5)	0.0656 (2)	
N1	0.40757 (18)	0.83635 (15)	0.35340 (14)	0.0376 (4)	
N2	0.5761 (2)	0.69351 (16)	0.40972 (14)	0.0417 (5)	
N3	0.5781 (2)	0.59926 (17)	0.18305 (14)	0.0439 (5)	
N4	0.32841 (19)	0.59452 (15)	0.22700 (14)	0.0402 (5)	
O1	0.9002 (2)	0.85085 (16)	0.42946 (13)	0.0634 (5)	
O2	0.9822 (2)	0.70605 (17)	0.36560 (14)	0.0701 (6)	

O3	1.1172 (3)	0.59708 (19)	0.47739 (18)	0.0844 (7)	
H3	1.064 (4)	0.613 (4)	0.428 (2)	0.127*	
O4	0.9776 (4)	0.9750 (3)	0.7937 (2)	0.0854 (15)	0.757 (7)
O5	1.0879 (5)	0.8618 (3)	0.9044 (3)	0.0724 (11)	0.757 (7)
O6	1.2238 (5)	0.9995 (3)	0.8373 (3)	0.1028 (16)	0.757 (7)
O4'	0.9391 (7)	0.8922 (10)	0.8218 (8)	0.104 (5)	0.243 (7)
O5'	1.1646 (14)	0.8780 (11)	0.9059 (9)	0.101 (5)	0.243 (7)
O6'	1.1326 (10)	1.0279 (5)	0.8124 (7)	0.054 (3)	0.243 (7)
O1W	0.69078 (17)	0.84787 (16)	0.28148 (14)	0.0517 (4)	
H11	0.757 (2)	0.841 (2)	0.3255 (15)	0.078*	
H12	0.713 (3)	0.8936 (19)	0.2443 (16)	0.078*	
O2W	0.42757 (19)	0.79208 (16)	0.13328 (14)	0.0555 (5)	
H21	0.3404 (11)	0.779 (2)	0.114 (2)	0.083*	
H22	0.459 (2)	0.8582 (13)	0.127 (2)	0.083*	
O3W	0.1536 (2)	0.73101 (18)	0.05280 (15)	0.0658 (5)	
H31	0.103 (3)	0.742 (3)	0.0938 (18)	0.099*	
H32	0.134 (3)	0.770 (2)	0.0068 (16)	0.099*	
O4W	0.2383 (3)	1.0300 (3)	0.0895 (2)	0.1163 (9)	
H41	0.182 (4)	1.052 (4)	0.123 (3)	0.174*	
H42	0.190 (4)	0.990 (4)	0.035 (2)	0.174*	
O5W	-0.0022 (2)	0.78419 (18)	0.18724 (16)	0.0698 (6)	
H51	-0.008 (4)	0.8523 (12)	0.190 (2)	0.105*	
H52	-0.010 (4)	0.763 (2)	0.2422 (14)	0.105*	
O6W	0.5032 (7)	1.0198 (5)	0.1338 (4)	0.0941 (15)	0.54 (6)
O6W'	0.4960 (7)	0.9808 (5)	0.0703 (5)	0.0941 (15)	0.46
H61	0.5655 (17)	1.030 (3)	0.1029 (18)	0.141*	
H62	0.429 (2)	1.038 (4)	0.105 (3)	0.141*	
C1	0.4359 (2)	0.82539 (18)	0.45066 (17)	0.0364 (5)	
C2	0.5265 (2)	0.74973 (19)	0.48134 (17)	0.0387 (5)	
C3	0.6591 (3)	0.6232 (2)	0.4370 (2)	0.0579 (7)	
H3A	0.6932	0.5842	0.3887	0.069*	
C4	0.6975 (3)	0.6053 (3)	0.5344 (3)	0.0764 (9)	
H4	0.7556	0.5550	0.5505	0.092*	
C5	0.6491 (3)	0.6622 (3)	0.6058 (2)	0.0751 (9)	
H5	0.6747	0.6512	0.6713	0.090*	
C6	0.5605 (3)	0.7376 (2)	0.58146 (19)	0.0545 (7)	
C7	0.5041 (3)	0.8011 (3)	0.6521 (2)	0.0681 (8)	
H7	0.5275	0.7939	0.7188	0.082*	
C8	0.4186 (3)	0.8704 (2)	0.6232 (2)	0.0589 (8)	
H8	0.3832	0.9101	0.6703	0.071*	
C9	0.3807 (2)	0.88451 (19)	0.52230 (19)	0.0443 (6)	
C10	0.2908 (3)	0.9557 (2)	0.4877 (2)	0.0531 (7)	
H10	0.2512	0.9960	0.5317	0.064*	
C11	0.2621 (3)	0.9652 (2)	0.3902 (2)	0.0542 (7)	
H11A	0.2022	1.0116	0.3669	0.065*	
C12	0.3231 (2)	0.90484 (19)	0.3246 (2)	0.0461 (6)	
H12A	0.3035	0.9132	0.2581	0.055*	
C13	0.4829 (3)	0.50289 (19)	0.15053 (16)	0.0411 (6)	
C14	0.3499 (2)	0.50073 (19)	0.17326 (16)	0.0407 (5)	

supplementary materials

C15	0.2076 (3)	0.5927 (2)	0.25028 (19)	0.0496 (6)
H15	0.1927	0.6562	0.2873	0.060*
C16	0.1008 (3)	0.4998 (2)	0.2220 (2)	0.0602 (7)
H16	0.0175	0.5020	0.2407	0.072*
C17	0.1194 (3)	0.4070 (2)	0.1673 (2)	0.0624 (8)
H17	0.0484	0.3453	0.1472	0.075*
C18	0.2475 (3)	0.4041 (2)	0.14070 (18)	0.0516 (7)
C19	0.2778 (4)	0.3086 (2)	0.0854 (2)	0.0712 (9)
H19	0.2102	0.2448	0.0633	0.085*
C20	0.4028 (4)	0.3099 (2)	0.0649 (2)	0.0696 (9)
H20	0.4206	0.2463	0.0299	0.084*
C21	0.5102 (3)	0.4076 (2)	0.09595 (18)	0.0533 (7)
C22	0.6413 (3)	0.4162 (3)	0.0741 (2)	0.0641 (8)
H22A	0.6641	0.3555	0.0382	0.077*
C23	0.7349 (3)	0.5124 (3)	0.1051 (2)	0.0644 (8)
H23	0.8214	0.5180	0.0901	0.077*
C24	0.7002 (3)	0.6032 (2)	0.15969 (19)	0.0551 (7)
H24	0.7652	0.6687	0.1805	0.066*
C25	0.9671 (3)	0.7750 (2)	0.4384 (2)	0.0512 (6)
C26	1.0362 (2)	0.7593 (2)	0.54018 (19)	0.0449 (6)
C27	1.0324 (2)	0.8337 (2)	0.62312 (18)	0.0442 (6)
H27	0.9882	0.8942	0.6145	0.053*
C28	1.0928 (3)	0.8198 (2)	0.71780 (19)	0.0484 (6)
C29	1.1577 (3)	0.7286 (3)	0.7311 (2)	0.0644 (8)
H29	1.1969	0.7177	0.7947	0.077*
C30	1.1638 (3)	0.6547 (3)	0.6508 (3)	0.0714 (9)
H30	1.2075	0.5941	0.6603	0.086*
C31	1.1054 (3)	0.6693 (2)	0.5552 (2)	0.0573 (7)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.04032 (17)	0.04133 (17)	0.03760 (16)	0.01290 (12)	0.00851 (12)	0.00213 (11)
S1	0.0877 (6)	0.0620 (5)	0.0438 (4)	0.0094 (4)	0.0046 (4)	0.0170 (4)
N1	0.0359 (11)	0.0355 (10)	0.0403 (11)	0.0103 (8)	0.0060 (9)	0.0005 (9)
N2	0.0415 (11)	0.0416 (11)	0.0453 (12)	0.0148 (9)	0.0107 (9)	0.0085 (9)
N3	0.0423 (12)	0.0524 (12)	0.0394 (11)	0.0176 (10)	0.0097 (9)	0.0041 (9)
N4	0.0406 (11)	0.0406 (11)	0.0409 (11)	0.0097 (9)	0.0107 (9)	0.0062 (9)
O1	0.0622 (12)	0.0758 (13)	0.0498 (11)	0.0292 (11)	-0.0026 (9)	0.0050 (10)
O2	0.0863 (15)	0.0732 (13)	0.0538 (12)	0.0241 (11)	0.0205 (11)	0.0014 (10)
O3	0.116 (2)	0.0723 (14)	0.0890 (18)	0.0524 (14)	0.0495 (16)	0.0187 (13)
O4	0.123 (3)	0.084 (3)	0.0535 (18)	0.055 (2)	0.0025 (17)	0.0085 (16)
O5	0.106 (3)	0.068 (2)	0.0442 (17)	0.009 (2)	0.0146 (19)	0.0256 (15)
O6	0.113 (3)	0.083 (2)	0.088 (2)	-0.029 (2)	-0.010 (2)	0.0347 (19)
O4'	0.100 (7)	0.099 (8)	0.115 (8)	-0.011 (5)	0.075 (6)	-0.024 (6)
O5'	0.139 (10)	0.096 (8)	0.065 (7)	0.025 (7)	0.007 (7)	0.026 (5)
O6'	0.066 (6)	0.046 (4)	0.064 (5)	0.021 (4)	0.025 (4)	0.025 (4)
O1W	0.0425 (10)	0.0623 (12)	0.0495 (11)	0.0061 (9)	0.0048 (8)	0.0192 (9)

O2W	0.0574 (11)	0.0635 (12)	0.0460 (11)	0.0180 (10)	0.0058 (10)	0.0111 (10)
O3W	0.0660 (13)	0.0783 (14)	0.0580 (13)	0.0261 (11)	0.0105 (10)	0.0185 (11)
O4W	0.145 (3)	0.096 (2)	0.099 (2)	0.0271 (19)	0.0184 (19)	-0.0103 (16)
O5W	0.0734 (14)	0.0677 (13)	0.0767 (15)	0.0189 (12)	0.0283 (12)	0.0169 (11)
O6W	0.115 (3)	0.091 (3)	0.078 (3)	0.013 (2)	0.024 (3)	0.023 (3)
O6W'	0.115 (3)	0.091 (3)	0.078 (3)	0.013 (2)	0.024 (3)	0.023 (3)
C1	0.0338 (12)	0.0333 (12)	0.0398 (13)	0.0011 (10)	0.0098 (10)	0.0001 (10)
C2	0.0371 (13)	0.0393 (13)	0.0390 (13)	0.0047 (10)	0.0080 (10)	0.0070 (10)
C3	0.0621 (18)	0.0594 (17)	0.0632 (19)	0.0307 (15)	0.0179 (15)	0.0210 (14)
C4	0.084 (2)	0.083 (2)	0.080 (2)	0.0452 (19)	0.0197 (19)	0.0421 (19)
C5	0.087 (2)	0.094 (2)	0.0559 (19)	0.032 (2)	0.0137 (17)	0.0404 (18)
C6	0.0597 (17)	0.0634 (17)	0.0434 (15)	0.0095 (14)	0.0134 (13)	0.0190 (13)
C7	0.084 (2)	0.084 (2)	0.0392 (16)	0.0082 (18)	0.0221 (15)	0.0144 (15)
C8	0.0627 (19)	0.0658 (18)	0.0488 (17)	0.0029 (15)	0.0277 (15)	-0.0032 (14)
C9	0.0404 (14)	0.0418 (13)	0.0487 (15)	0.0005 (11)	0.0170 (12)	-0.0036 (11)
C10	0.0445 (15)	0.0436 (14)	0.072 (2)	0.0066 (12)	0.0261 (14)	-0.0097 (13)
C11	0.0415 (15)	0.0450 (15)	0.076 (2)	0.0176 (12)	0.0113 (14)	0.0008 (14)
C12	0.0420 (14)	0.0425 (14)	0.0535 (16)	0.0158 (11)	0.0050 (12)	0.0060 (12)
C13	0.0552 (15)	0.0397 (13)	0.0308 (12)	0.0174 (12)	0.0069 (11)	0.0088 (10)
C14	0.0516 (15)	0.0383 (13)	0.0314 (12)	0.0094 (11)	0.0043 (11)	0.0095 (10)
C15	0.0439 (15)	0.0537 (16)	0.0527 (16)	0.0095 (12)	0.0134 (12)	0.0091 (12)
C16	0.0488 (17)	0.0710 (19)	0.0586 (18)	0.0008 (14)	0.0124 (14)	0.0139 (15)
C17	0.0621 (19)	0.0613 (18)	0.0537 (18)	-0.0128 (15)	0.0043 (14)	0.0145 (15)
C18	0.0693 (19)	0.0425 (14)	0.0380 (14)	0.0025 (13)	0.0045 (13)	0.0095 (11)
C19	0.106 (3)	0.0404 (16)	0.059 (2)	0.0000 (16)	0.0144 (19)	0.0007 (14)
C20	0.114 (3)	0.0418 (16)	0.0536 (18)	0.0264 (17)	0.0167 (19)	-0.0018 (13)
C21	0.083 (2)	0.0474 (15)	0.0351 (14)	0.0336 (15)	0.0106 (13)	0.0069 (12)
C22	0.092 (2)	0.073 (2)	0.0411 (16)	0.0530 (19)	0.0180 (16)	0.0075 (14)
C23	0.0593 (18)	0.101 (2)	0.0461 (17)	0.0459 (18)	0.0169 (14)	0.0138 (16)
C24	0.0457 (15)	0.0748 (19)	0.0482 (16)	0.0234 (14)	0.0118 (13)	0.0053 (14)
C25	0.0460 (15)	0.0575 (16)	0.0501 (16)	0.0075 (13)	0.0133 (13)	0.0064 (13)
C26	0.0374 (13)	0.0493 (14)	0.0539 (16)	0.0117 (11)	0.0170 (12)	0.0151 (12)
C27	0.0372 (13)	0.0491 (14)	0.0500 (15)	0.0132 (11)	0.0106 (11)	0.0139 (12)
C28	0.0452 (15)	0.0537 (15)	0.0515 (16)	0.0091 (12)	0.0146 (12)	0.0216 (13)
C29	0.0629 (19)	0.078 (2)	0.066 (2)	0.0272 (16)	0.0183 (15)	0.0413 (17)
C30	0.082 (2)	0.071 (2)	0.088 (2)	0.0460 (18)	0.0382 (19)	0.0433 (19)
C31	0.0589 (17)	0.0553 (16)	0.070 (2)	0.0202 (14)	0.0297 (15)	0.0212 (15)

Geometric parameters (Å, °)

Zn1—O2W	2.0810 (18)	C4—C5	1.358 (4)
Zn1—O1W	2.0914 (17)	C4—H4	0.9300
Zn1—N1	2.1615 (17)	C5—C6	1.406 (4)
Zn1—N2	2.164 (2)	C5—H5	0.9300
Zn1—N3	2.1720 (18)	C6—C7	1.438 (4)
Zn1—N4	2.2130 (19)	C7—C8	1.341 (4)
S1—O6'	1.372 (5)	C7—H7	0.9300
S1—O5	1.422 (3)	C8—C9	1.420 (4)
S1—O4	1.433 (3)	C8—H8	0.9300

supplementary materials

S1—O5'	1.448 (6)	C9—C10	1.409 (4)
S1—O6	1.503 (3)	C10—C11	1.354 (4)
S1—O4'	1.511 (6)	C10—H10	0.9300
S1—C28	1.773 (3)	C11—C12	1.400 (3)
N1—C12	1.325 (3)	C11—H11A	0.9300
N1—C1	1.355 (3)	C12—H12A	0.9300
N2—C3	1.327 (3)	C13—C21	1.410 (3)
N2—C2	1.363 (3)	C13—C14	1.434 (3)
N3—C24	1.330 (3)	C14—C18	1.407 (3)
N3—C13	1.367 (3)	C15—C16	1.398 (4)
N4—C15	1.318 (3)	C15—H15	0.9300
N4—C14	1.365 (3)	C16—C17	1.351 (4)
O1—C25	1.234 (3)	C16—H16	0.9300
O2—C25	1.288 (3)	C17—C18	1.417 (4)
O3—C31	1.351 (3)	C17—H17	0.9300
O3—H3	0.853 (11)	C18—C19	1.430 (4)
O1W—H11	0.832 (10)	C19—C20	1.344 (4)
O1W—H12	0.843 (10)	C19—H19	0.9300
O2W—H21	0.849 (10)	C20—C21	1.443 (4)
O2W—H22	0.845 (10)	C20—H20	0.9300
O3W—H31	0.849 (10)	C21—C22	1.405 (4)
O3W—H32	0.853 (10)	C22—C23	1.356 (4)
O4W—H41	0.859 (11)	C22—H22A	0.9300
O4W—H42	0.858 (11)	C23—C24	1.400 (4)
O5W—H51	0.844 (10)	C23—H23	0.9300
O5W—H52	0.851 (10)	C24—H24	0.9300
O6W—O6W'	0.937 (6)	C25—C26	1.498 (4)
O6W—H61	0.831 (11)	C26—C27	1.394 (3)
O6W—H62	0.843 (8)	C26—C31	1.408 (3)
O6W'—H61	0.857 (11)	C27—C28	1.380 (3)
O6W'—H62	1.17 (3)	C27—H27	0.9300
C1—C9	1.415 (3)	C28—C29	1.393 (4)
C1—C2	1.441 (3)	C29—C30	1.369 (4)
C2—C6	1.400 (3)	C29—H29	0.9300
C3—C4	1.388 (4)	C30—C31	1.388 (4)
C3—H3A	0.9300	C30—H30	0.9300
O2W—Zn1—O1W	89.83 (7)	C2—C6—C5	116.4 (2)
O2W—Zn1—N1	93.87 (7)	C2—C6—C7	119.3 (3)
O1W—Zn1—N1	97.42 (7)	C5—C6—C7	124.3 (3)
O2W—Zn1—N2	171.00 (7)	C8—C7—C6	121.0 (3)
O1W—Zn1—N2	90.11 (7)	C8—C7—H7	119.5
N1—Zn1—N2	77.21 (7)	C6—C7—H7	119.5
O2W—Zn1—N3	92.55 (7)	C7—C8—C9	121.5 (2)
O1W—Zn1—N3	94.26 (8)	C7—C8—H8	119.2
N1—Zn1—N3	166.67 (7)	C9—C8—H8	119.2
N2—Zn1—N3	96.43 (7)	C10—C9—C1	116.7 (2)
O2W—Zn1—N4	89.26 (7)	C10—C9—C8	124.0 (2)
O1W—Zn1—N4	170.12 (7)	C1—C9—C8	119.4 (2)
N1—Zn1—N4	92.46 (7)	C11—C10—C9	119.9 (2)

N2—Zn1—N4	92.32 (7)	C11—C10—H10	120.0
N3—Zn1—N4	75.96 (7)	C9—C10—H10	120.0
O6'—S1—O5	133.1 (4)	C10—C11—C12	119.6 (2)
O6'—S1—O4	67.2 (4)	C10—C11—H11A	120.2
O5—S1—O4	115.8 (2)	C12—C11—H11A	120.2
O6'—S1—O5'	116.8 (5)	N1—C12—C11	122.8 (2)
O5—S1—O5'	30.8 (5)	N1—C12—H12A	118.6
O4—S1—O5'	140.9 (7)	C11—C12—H12A	118.6
O6'—S1—O6	42.2 (3)	N3—C13—C21	122.9 (2)
O5—S1—O6	110.5 (2)	N3—C13—C14	117.6 (2)
O4—S1—O6	109.3 (2)	C21—C13—C14	119.5 (2)
O5'—S1—O6	82.5 (5)	N4—C14—C18	122.4 (2)
O6'—S1—O4'	112.4 (5)	N4—C14—C13	117.5 (2)
O5—S1—O4'	76.4 (5)	C18—C14—C13	120.0 (2)
O4—S1—O4'	46.8 (5)	N4—C15—C16	123.2 (2)
O5'—S1—O4'	107.2 (5)	N4—C15—H15	118.4
O6—S1—O4'	151.4 (4)	C16—C15—H15	118.4
O6'—S1—C28	114.8 (3)	C17—C16—C15	119.5 (3)
O5—S1—C28	108.38 (19)	C17—C16—H16	120.2
O4—S1—C28	108.01 (15)	C15—C16—H16	120.2
O5'—S1—C28	104.5 (6)	C16—C17—C18	119.7 (3)
O6—S1—C28	104.2 (2)	C16—C17—H17	120.2
O4'—S1—C28	99.3 (4)	C18—C17—H17	120.2
C12—N1—C1	118.06 (19)	C14—C18—C17	117.0 (2)
C12—N1—Zn1	128.06 (16)	C14—C18—C19	119.4 (3)
C1—N1—Zn1	113.88 (14)	C17—C18—C19	123.6 (3)
C3—N2—C2	117.8 (2)	C20—C19—C18	120.8 (3)
C3—N2—Zn1	128.40 (17)	C20—C19—H19	119.6
C2—N2—Zn1	113.79 (15)	C18—C19—H19	119.6
C24—N3—C13	118.0 (2)	C19—C20—C21	121.7 (3)
C24—N3—Zn1	126.88 (18)	C19—C20—H20	119.2
C13—N3—Zn1	115.07 (15)	C21—C20—H20	119.2
C15—N4—C14	118.1 (2)	C22—C21—C13	116.6 (3)
C15—N4—Zn1	128.09 (16)	C22—C21—C20	124.7 (2)
C14—N4—Zn1	113.80 (15)	C13—C21—C20	118.6 (3)
C31—O3—H3	104 (3)	C23—C22—C21	120.4 (2)
Zn1—O1W—H11	117.2 (17)	C23—C22—H22A	119.8
Zn1—O1W—H12	129.0 (17)	C21—C22—H22A	119.8
H11—O1W—H12	112.9 (17)	C22—C23—C24	119.5 (3)
Zn1—O2W—H21	116 (2)	C22—C23—H23	120.3
Zn1—O2W—H22	118 (2)	C24—C23—H23	120.3
H21—O2W—H22	109.6 (16)	N3—C24—C23	122.6 (3)
H31—O3W—H32	108.2 (16)	N3—C24—H24	118.7
H41—O4W—H42	107.2 (18)	C23—C24—H24	118.7
H51—O5W—H52	109.5 (17)	O1—C25—O2	124.6 (3)
O6W'—O6W—H61	57.6 (9)	O1—C25—C26	119.1 (2)
O6W'—O6W—H62	82 (3)	O2—C25—C26	116.2 (2)
H61—O6W—H62	113.0 (18)	C27—C26—C31	118.1 (2)
O6W—O6W'—H61	54.9 (9)	C27—C26—C25	120.4 (2)

supplementary materials

O6W—O6W'—H62	45.5 (13)	C31—C26—C25	121.5 (2)
H61—O6W'—H62	86 (2)	C28—C27—C26	121.5 (2)
N1—C1—C9	123.0 (2)	C28—C27—H27	119.2
N1—C1—C2	117.85 (19)	C26—C27—H27	119.2
C9—C1—C2	119.2 (2)	C27—C28—C29	119.3 (3)
N2—C2—C6	123.1 (2)	C27—C28—S1	120.31 (19)
N2—C2—C1	117.3 (2)	C29—C28—S1	120.4 (2)
C6—C2—C1	119.6 (2)	C30—C29—C28	120.3 (3)
N2—C3—C4	123.0 (3)	C30—C29—H29	119.9
N2—C3—H3A	118.5	C28—C29—H29	119.9
C4—C3—H3A	118.5	C29—C30—C31	120.7 (3)
C5—C4—C3	119.0 (3)	C29—C30—H30	119.6
C5—C4—H4	120.5	C31—C30—H30	119.6
C3—C4—H4	120.5	O3—C31—C30	119.2 (3)
C4—C5—C6	120.6 (3)	O3—C31—C26	120.8 (3)
C4—C5—H5	119.7	C30—C31—C26	120.0 (3)
C6—C5—H5	119.7		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1W—H11 \cdots O1	0.83 (1)	1.81 (1)	2.632 (2)	170 (3)
O1W—H12 \cdots O6 ⁱ	0.84 (1)	1.95 (1)	2.793 (4)	177 (2)
O1W—H12 \cdots O6 ⁱ	0.84 (1)	2.04 (2)	2.787 (7)	147 (2)
O2W—H21 \cdots O3W	0.85 (1)	1.87 (1)	2.714 (3)	170 (3)
O2W—H22 \cdots O6W	0.85 (1)	1.95 (1)	2.778 (6)	166 (3)
O2W—H22 \cdots O6W'	0.85 (1)	1.80 (2)	2.615 (6)	160 (3)
O3W—H31 \cdots O5W	0.85 (1)	1.91 (1)	2.754 (3)	173 (3)
O3W—H32 \cdots O5 ⁱⁱ	0.85 (1)	1.95 (1)	2.805 (4)	178 (3)
O3W—H32 \cdots O5 ⁱⁱⁱ	0.85 (1)	2.08 (2)	2.892 (10)	159 (3)
O4W—H41 \cdots O4 ⁱⁱⁱ	0.86 (1)	2.17 (3)	2.962 (5)	153 (5)
O4W—H41 \cdots O4 ⁱⁱⁱ	0.86 (1)	1.74 (2)	2.598 (8)	173 (5)
O4W—H42 \cdots O5 ⁱⁱ	0.86 (1)	2.22 (1)	3.065 (5)	169 (5)
O4W—H42 \cdots O5 ⁱⁱⁱ	0.86 (1)	2.08 (2)	2.869 (15)	153 (4)
O5W—H51 \cdots O4 ⁱⁱⁱ	0.84 (1)	2.07 (1)	2.900 (4)	168 (4)
O5W—H51 \cdots O6 ⁱⁱⁱ	0.84 (1)	2.07 (2)	2.823 (8)	148 (3)
O5W—H52 \cdots O2 ^{iv}	0.85 (1)	1.94 (1)	2.792 (3)	175 (3)
O6W—H61 \cdots O6 ⁱ	0.83 (1)	2.22 (3)	2.751 (8)	122 (3)
O6W—H62 \cdots O4W	0.84 (1)	1.88 (2)	2.642 (7)	150 (4)

Symmetry codes: (i) $-x+2, -y+2, -z+1$; (ii) $x-1, y, z-1$; (iii) $-x+1, -y+2, -z+1$; (iv) $x-1, y, z$.

Fig. 1

