

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# 3,3'-Dibenzoyl-1,1'-(3,6-dioxaoctane-1,8-diyl)dithiourea

Mouhamadou Moustapha Sow,<sup>a</sup> Ousmane Diouf,<sup>a</sup>  
Aliou Hamady Barry,<sup>b</sup> Mohamed Gaye<sup>a\*</sup> and  
Abdou Salam Sall<sup>a</sup>

<sup>a</sup>Département de Chimie, Faculté des Sciences et Techniques, Université Cheikh Anta Diop, Dakar, Senegal, and <sup>b</sup>Département de Chimie, Faculté des Sciences, Université de Nouakchott, Nouakchott, Mauritanie  
Correspondence e-mail: mlgayeastou@yahoo.fr


Received 15 January 2009; accepted 17 February 2009

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.045;  $wR$  factor = 0.128; data-to-parameter ratio = 13.8.

In the molecule of the title compound,  $\text{C}_{22}\text{H}_{26}\text{N}_4\text{O}_4\text{S}_2$ , the central  $\text{O}-\text{CH}_2-\text{CH}_2-\text{O}$  chain adopts a synclinal conformation [torsion angle =  $65.0(2)^\circ$ ]. The crystal structure is stabilized by intramolecular  $\text{N}-\text{H}\cdots\text{O}=\text{C}$  and intermolecular  $\text{N}-\text{H}\cdots\text{O}-\text{C}$  hydrogen bonds.

## Related literature

For related structures, see: Avşar *et al.* (2003); Arslan *et al.* (2004); Du & Du (2008); Ding *et al.* (2008).



## Experimental

### Crystal data

$\text{C}_{22}\text{H}_{26}\text{N}_4\text{O}_4\text{S}_2$   
 $M_r = 474.59$   
Triclinic,  $P\bar{1}$   
 $a = 7.9718(2)$  Å  
 $b = 9.2177(3)$  Å

$c = 16.4106(5)$  Å  
 $\alpha = 81.018(2)^\circ$   
 $\beta = 83.364(2)^\circ$   
 $\gamma = 80.450(2)^\circ$   
 $V = 1169.60(6)$  Å<sup>3</sup>

$Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.26$  mm<sup>-1</sup>

$T = 293$  K  
 $0.10 \times 0.10 \times 0.10$  mm

### Data collection

Nonius Kappa CCD diffractometer  
Absorption correction: none  
7989 measured reflections

4211 independent reflections  
2737 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.030$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.128$   
 $S = 1.02$   
4211 reflections  
305 parameters

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.41$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.50$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}2-\text{H}2\cdots\text{O}1$	0.82 (3)	1.99 (3)	2.648 (3)	136 (2)
$\text{N}3-\text{H}3\cdots\text{O}4$	0.83 (3)	2.01 (3)	2.632 (3)	132 (3)
$\text{N}4-\text{H}4\cdots\text{O}2^i$	0.82 (3)	2.48 (3)	3.290 (3)	170 (2)

 Symmetry code: (i)  $-x + 2, -y, -z + 1$ .

Data collection: *COLLECT* (Nonius, 1998); cell refinement: *DENZO* and *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* and *SCALEPACK*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

The authors thank the Agence Universitaire de la Francophonie for financial support (AUF-PSCI No.6314PS804).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2217).

## References

- Arslan, H., Flörke, U. & Külcü, N. (2004). *Acta Chim. Slov.* **51**, 787–792.  
Avşar, G., Arslan, H., Haupt, H.-J. & Külcü, N. (2003). *Turk. J. Chem.* **27**, 281–285.  
Ding, Y.-J., Chang, X.-B., Yang, X.-Q. & Dong, W.-K. (2008). *Acta Cryst.* **E64**, o658.  
Du, H.-T. & Du, H.-J. (2008). *Acta Cryst.* **E64**, o1632–o1633.  
Nonius (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.  
Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

**supplementary materials**

*Acta Cryst.* (2009). E65, o569 [ doi:10.1107/S1600536809005662 ]

### 3,3'-Dibenzoyl-1,1'-(3,6-dioxaoctane-1,8-diyl)dithiourea

M. M. Sow, O. Diouf, A. H. Barry, M. Gaye and A. S. Sall

#### Comment

The title compound, C<sub>22</sub>H<sub>26</sub>N<sub>4</sub>O<sub>4</sub>S<sub>2</sub>, was characterized by <sup>1</sup>H and <sup>13</sup>C NMR, solid-state IR and X-ray crystallographic techniques. The X-ray structure determination reveals that the compound crystallizes in the triclinic space group *P* $\bar{1}$  with one molecule in the asymmetric unit. The molecular geometry is illustrated in Fig. 1. The C—S bond lengths of 1.665 (3) Å and 1.659 (2) Å and the C—O bond lengths of 1.220 (3) Å and 1.222 (3) Å are double bonds character and are comparable to those observed for 1-(biphenyl-4-carbonyl)-3-*p*-tolyl-thiourea [1.647 (3) Å for C—S, 1.217 (3) and 1.224 (3) Å for C—O respectively (Arslan *et al.*, 2004)]. The C—N bond lengths are in the range 1.310 (3)–1.451 (3) Å, and are shorter than the normal single C—N bond length (Avşar *et al.*, 2003). The carbonyl group forms an intramolecular hydrogen bonds with the N2—H2 and the N3—H3 groups, which forms two six-membered rings (C8/N1/C7/O1/H2/N2 and C15/N4/C16/O4/H3/N3) structure (Fig. 2); H2···O1 and the H3···O4 separations are respectively 1.99 (3) Å and 2.01 (3) Å. There is an intermolecular hydrogen bonding between N4—H4 and the O atom of the ethoxy group of a symmetry-related molecule, the H4···O2 (-x + 2, -y, -z + 1) separation being 2.48 (3) Å (Table 1). The structure of the title compound is related to other thiourea derivatives (*e.g.* Ding *et al.*, 2008; Du & Du, 2008).

#### Experimental

Benzoyl chloride (7.10 g, 50 mmol) was reacted with potassium thiocyanate (4.86 g, 50 mmol) in CH<sub>3</sub>OCH<sub>3</sub> (50 ml) solution, to give the corresponding benzoyl isothiocyanate after one hour under refluxing. After cooling to room temperature, a solution of 2-(2-(2-aminoethoxy)ethoxy)ethanamine (3.70 g, 25 mmol) in CH<sub>3</sub>OCH<sub>3</sub> (20 ml) was added dropwise to benzoyl isothiocyanate. After three hours under stirring, 200 ml of HCl 1 *M* was added. A yellow oil was isolated and treated with diethyl ether to give the title compound which is washed with diethyl ether twice. Yield: 55.9%. m.p. 415–419 K. Anal. Calc. for C<sub>22</sub>H<sub>26</sub>N<sub>4</sub>O<sub>4</sub>S<sub>2</sub>: C 55.68, H 5.52, N 11.81%. Found: C 55.70, H 5.45, N 11.65%. Selected IR data (cm<sup>-1</sup>, KBr pellet): 3424, 3218 (ν NH), 1667 (ν C?O), 1160 (ν C?S). <sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>, δ, p.p.m.): 3.45 (t, 4H, N—CH<sub>2</sub>); 3.63 (s, 4H, O—CH<sub>2</sub>); 3.70 (t, 4H, O—CH<sub>2</sub>); 7.21–7.92 (m, 10H, C<sub>6</sub>H<sub>5</sub>); 11.01 (s, 2H, NH); 11.41 (s, 2H, NH). <sup>13</sup>C-NMR (50 MHz, DMSO-d<sub>6</sub>, δ, p.p.m.): 45.21 (N—CH<sub>2</sub>); 68.09 (O—CH<sub>2</sub>); 70.12 (O—CH<sub>2</sub>); 133.31–127.62 (C<sub>6</sub>H<sub>5</sub>); 1168.60 (C?O); 180.81 (C?S). A CH<sub>3</sub>Cl solution of the title compound was mixed with ethanol (1/1). After several days, colorless block-shaped single crystals suitable for X-ray crystallographic analysis were obtained.

#### Refinement

H atoms of NH groups were located in a difference map and refined freely. Others H atoms were placed geometrically and refined with a riding model. *U*<sub>iso</sub>(H) for H was calculated as 1.2 *U*<sub>eq</sub> of the carrier atom.

## Figures



Fig. 1. An *ORTEP* view of the asymmetric unit of the title compound, showing the atom-numbering scheme. Displacement ellipsoids are plotted at the 50% probability level.




Fig. 2. Molecular representation of the compound showing hydrogen bonds (dashed lines).

## 3,3'-Dibenzoyl-1,1'-(3,6-dioxaoctane-1,8-diyl)dithiourea

### Crystal data

$C_{22}H_{26}N_4O_4S_2$

$M_r = 474.59$

Triclinic,  $P\bar{1}$

Hall symbol: -P 1

$a = 7.9718$  (2) Å

$b = 9.2177$  (3) Å

$c = 16.4106$  (5) Å

$\alpha = 81.018$  (2)°

$\beta = 83.364$  (2)°

$\gamma = 80.450$  (2)°

$V = 1169.60$  (6) Å<sup>3</sup>

$Z = 2$

$F_{000} = 500$

$D_x = 1.347$  Mg m<sup>-3</sup>

Melting point: 415 K

Mo  $K\alpha$  radiation

$\lambda = 0.71073$  Å

Cell parameters from 4206 reflections

$\theta = 1.0$ – $25.4$ °

$\mu = 0.26$  mm<sup>-1</sup>

$T = 293$  K

Prism, yellow

$0.10 \times 0.10 \times 0.10$  mm

### Data collection

Nonius KappaCCD diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293$  K

$\omega$  scans

Absorption correction: none

7989 measured reflections

4211 independent reflections

2737 reflections with  $I > 2\sigma(I)$

$R_{int} = 0.030$

$\theta_{max} = 25.2$ °

$\theta_{min} = 2.3$ °

$h = -9 \rightarrow 9$

$k = -11 \rightarrow 11$

$l = -19 \rightarrow 19$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.045$

$wR(F^2) = 0.128$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0614P)^2 + 0.2609P]$

$S = 1.02$

4211 reflections

305 parameters

Primary atom site location: structure-invariant direct methods

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.008$

$\Delta\rho_{\max} = 0.41 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.50 \text{ e } \text{\AA}^{-3}$

Extinction correction: none

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.37094 (10)	0.16706 (9)	0.86251 (5)	0.0713 (3)
S2	1.26452 (13)	-0.27342 (8)	0.52516 (4)	0.0790 (3)
O1	0.7377 (3)	0.4586 (2)	0.91685 (11)	0.0711 (6)
O2	0.81088 (19)	0.39803 (17)	0.67050 (9)	0.0482 (4)
O3	1.12041 (19)	0.24744 (17)	0.60508 (10)	0.0472 (4)
O4	1.2463 (3)	0.1608 (2)	0.35665 (11)	0.0740 (6)
N1	0.5641 (3)	0.2839 (3)	0.94678 (14)	0.0527 (6)
H1	0.530 (3)	0.224 (3)	0.9815 (16)	0.049 (8)*
N2	0.5680 (3)	0.3729 (2)	0.80784 (13)	0.0492 (5)
H2	0.639 (3)	0.422 (3)	0.8172 (16)	0.056 (8)*
N3	1.2498 (3)	0.0165 (3)	0.50822 (13)	0.0506 (5)
H3	1.236 (3)	0.099 (3)	0.4798 (17)	0.063 (9)*
N4	1.2073 (3)	-0.0786 (2)	0.39123 (12)	0.0465 (5)
H4	1.190 (3)	-0.154 (3)	0.3741 (15)	0.052 (8)*
C1	0.7116 (3)	0.3433 (3)	1.05616 (14)	0.0456 (6)
C2	0.8101 (3)	0.4370 (3)	1.08034 (16)	0.0539 (6)
H2A	0.8497	0.5115	1.0416	0.065*
C3	0.8498 (4)	0.4211 (3)	1.16088 (18)	0.0639 (7)
H3A	0.9169	0.4840	1.1763	0.077*
C4	0.7902 (4)	0.3120 (3)	1.21864 (18)	0.0692 (8)
H4A	0.8165	0.3012	1.2732	0.083*
C5	0.6923 (4)	0.2194 (4)	1.19573 (19)	0.0771 (9)
H5	0.6513	0.1464	1.2350	0.093*
C6	0.6537 (4)	0.2335 (3)	1.11472 (17)	0.0649 (8)
H6	0.5885	0.1690	1.0995	0.078*
C7	0.6744 (3)	0.3674 (3)	0.96796 (15)	0.0488 (6)
C8	0.5078 (3)	0.2817 (3)	0.86956 (15)	0.0477 (6)
C9	0.5275 (3)	0.3838 (3)	0.72324 (15)	0.0537 (6)
H9A	0.4095	0.4293	0.7189	0.064*
H9B	0.5406	0.2851	0.7075	0.064*
C10	0.6419 (3)	0.4745 (3)	0.66582 (15)	0.0516 (6)
H10A	0.6101	0.4859	0.6096	0.062*
H10B	0.6331	0.5723	0.6822	0.062*
C11	0.9346 (3)	0.4733 (3)	0.61860 (16)	0.0522 (6)
H11A	0.9278	0.5730	0.6320	0.063*
H11B	0.9133	0.4804	0.5610	0.063*
C12	1.1067 (3)	0.3882 (3)	0.63208 (16)	0.0506 (6)
H12A	1.1936	0.4426	0.6014	0.061*

## supplementary materials

---

H12B	1.1243	0.3751	0.6904	0.061*
C13	1.2832 (3)	0.1611 (3)	0.61572 (16)	0.0529 (6)
H13A	1.3092	0.1538	0.6727	0.063*
H13B	1.3705	0.2077	0.5799	0.063*
C14	1.2803 (3)	0.0091 (3)	0.59436 (14)	0.0504 (6)
H14A	1.3887	-0.0529	0.6045	0.061*
H14B	1.1912	-0.0359	0.6297	0.061*
C15	1.2389 (3)	-0.1016 (3)	0.47501 (14)	0.0447 (6)
C16	1.2118 (3)	0.0483 (3)	0.33592 (14)	0.0454 (6)
C17	1.1796 (3)	0.0433 (2)	0.24907 (14)	0.0419 (5)
C18	1.2227 (3)	0.1595 (3)	0.19018 (15)	0.0527 (6)
H18	1.2684	0.2361	0.2062	0.063*
C19	1.1984 (3)	0.1624 (3)	0.10851 (16)	0.0610 (7)
H19	1.2299	0.2398	0.0693	0.073*
C20	1.1281 (3)	0.0520 (3)	0.08436 (16)	0.0618 (7)
H20	1.1120	0.0546	0.0289	0.074*
C21	1.0811 (3)	-0.0632 (3)	0.14224 (16)	0.0605 (7)
H21	1.0324	-0.1378	0.1260	0.073*
C22	1.1069 (3)	-0.0672 (3)	0.22467 (15)	0.0507 (6)
H22	1.0751	-0.1446	0.2638	0.061*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0807 (5)	0.0781 (5)	0.0659 (5)	-0.0445 (4)	-0.0032 (4)	-0.0115 (4)
S2	0.1467 (8)	0.0464 (4)	0.0494 (4)	-0.0296 (5)	-0.0275 (5)	0.0059 (3)
O1	0.0906 (14)	0.0822 (14)	0.0503 (11)	-0.0502 (12)	-0.0123 (10)	0.0050 (10)
O2	0.0498 (10)	0.0443 (9)	0.0461 (9)	-0.0077 (8)	0.0014 (8)	0.0036 (7)
O3	0.0497 (10)	0.0457 (9)	0.0501 (10)	-0.0119 (8)	-0.0053 (7)	-0.0140 (8)
O4	0.1328 (18)	0.0430 (11)	0.0518 (11)	-0.0288 (11)	-0.0176 (11)	-0.0010 (9)
N1	0.0584 (14)	0.0572 (14)	0.0453 (13)	-0.0237 (11)	0.0023 (11)	-0.0054 (11)
N2	0.0468 (12)	0.0626 (14)	0.0431 (12)	-0.0214 (11)	0.0000 (10)	-0.0112 (10)
N3	0.0716 (15)	0.0397 (13)	0.0409 (12)	-0.0077 (11)	-0.0081 (10)	-0.0055 (10)
N4	0.0666 (14)	0.0372 (12)	0.0385 (11)	-0.0162 (10)	-0.0079 (9)	-0.0024 (9)
C1	0.0450 (13)	0.0445 (14)	0.0455 (14)	-0.0027 (11)	-0.0034 (11)	-0.0053 (11)
C2	0.0562 (15)	0.0556 (16)	0.0514 (15)	-0.0127 (13)	-0.0023 (12)	-0.0093 (12)
C3	0.0698 (18)	0.0677 (18)	0.0579 (17)	-0.0074 (15)	-0.0134 (14)	-0.0177 (15)
C4	0.076 (2)	0.078 (2)	0.0507 (17)	0.0048 (17)	-0.0169 (15)	-0.0081 (16)
C5	0.087 (2)	0.080 (2)	0.0591 (19)	-0.0181 (18)	-0.0171 (16)	0.0192 (16)
C6	0.0748 (19)	0.0598 (17)	0.0612 (18)	-0.0203 (15)	-0.0184 (15)	0.0077 (14)
C7	0.0483 (14)	0.0492 (15)	0.0496 (15)	-0.0124 (12)	-0.0012 (12)	-0.0062 (12)
C8	0.0461 (13)	0.0507 (15)	0.0486 (15)	-0.0121 (11)	0.0023 (11)	-0.0135 (12)
C9	0.0498 (15)	0.0667 (17)	0.0491 (15)	-0.0132 (13)	-0.0048 (12)	-0.0167 (13)
C10	0.0562 (15)	0.0521 (15)	0.0451 (14)	-0.0026 (12)	-0.0084 (12)	-0.0056 (12)
C11	0.0639 (16)	0.0404 (14)	0.0514 (15)	-0.0188 (12)	0.0041 (12)	0.0006 (11)
C12	0.0578 (15)	0.0444 (14)	0.0527 (15)	-0.0208 (12)	0.0023 (12)	-0.0082 (12)
C13	0.0527 (15)	0.0607 (16)	0.0485 (15)	-0.0100 (13)	-0.0065 (12)	-0.0144 (12)
C14	0.0612 (16)	0.0531 (15)	0.0379 (13)	-0.0040 (12)	-0.0102 (11)	-0.0100 (11)

C15	0.0512 (14)	0.0442 (14)	0.0399 (13)	-0.0115 (11)	-0.0051 (11)	-0.0044 (11)
C16	0.0543 (14)	0.0392 (14)	0.0426 (13)	-0.0097 (11)	-0.0041 (11)	-0.0023 (11)
C17	0.0440 (13)	0.0396 (13)	0.0399 (13)	-0.0052 (10)	-0.0027 (10)	-0.0009 (10)
C18	0.0591 (16)	0.0488 (15)	0.0492 (15)	-0.0153 (12)	-0.0033 (12)	0.0029 (12)
C19	0.0667 (18)	0.0656 (18)	0.0458 (16)	-0.0163 (15)	-0.0043 (13)	0.0132 (13)
C20	0.0631 (17)	0.080 (2)	0.0399 (14)	-0.0112 (15)	-0.0098 (13)	0.0033 (14)
C21	0.0691 (17)	0.0629 (17)	0.0527 (16)	-0.0161 (14)	-0.0134 (13)	-0.0059 (13)
C22	0.0572 (15)	0.0500 (15)	0.0434 (14)	-0.0135 (12)	-0.0065 (12)	0.0052 (11)

*Geometric parameters (Å, °)*

S1—C8	1.665 (3)	C5—H5	0.9300
S2—C15	1.659 (2)	C6—H6	0.9300
O1—C7	1.220 (3)	C9—C10	1.492 (3)
O2—C10	1.417 (3)	C9—H9A	0.9700
O2—C11	1.424 (3)	C9—H9B	0.9700
O3—C12	1.419 (3)	C10—H10A	0.9700
O3—C13	1.419 (3)	C10—H10B	0.9700
O4—C16	1.222 (3)	C11—C12	1.485 (3)
N1—C7	1.366 (3)	C11—H11A	0.9700
N1—C8	1.396 (3)	C11—H11B	0.9700
N1—H1	0.79 (3)	C12—H12A	0.9700
N2—C8	1.311 (3)	C12—H12B	0.9700
N2—C9	1.446 (3)	C13—C14	1.501 (3)
N2—H2	0.82 (3)	C13—H13A	0.9700
N3—C15	1.310 (3)	C13—H13B	0.9700
N3—C14	1.451 (3)	C14—H14A	0.9700
N3—H3	0.83 (3)	C14—H14B	0.9700
N4—C16	1.367 (3)	C16—C17	1.486 (3)
N4—C15	1.404 (3)	C17—C22	1.382 (3)
N4—H4	0.82 (3)	C17—C18	1.386 (3)
C1—C6	1.382 (3)	C18—C19	1.372 (4)
C1—C2	1.387 (3)	C18—H18	0.9300
C1—C7	1.486 (3)	C19—C20	1.369 (4)
C2—C3	1.375 (4)	C19—H19	0.9300
C2—H2A	0.9300	C20—C21	1.380 (4)
C3—C4	1.376 (4)	C20—H20	0.9300
C3—H3A	0.9300	C21—C22	1.386 (4)
C4—C5	1.366 (4)	C21—H21	0.9300
C4—H4A	0.9300	C22—H22	0.9300
C5—C6	1.381 (4)		
C10—O2—C11	113.12 (18)	H10A—C10—H10B	108.5
C12—O3—C13	112.15 (18)	O2—C11—C12	108.43 (19)
C7—N1—C8	129.4 (2)	O2—C11—H11A	110.0
C7—N1—H1	117.6 (19)	C12—C11—H11A	110.0
C8—N1—H1	112.9 (19)	O2—C11—H11B	110.0
C8—N2—C9	124.1 (2)	C12—C11—H11B	110.0
C8—N2—H2	117.9 (18)	H11A—C11—H11B	108.4
C9—N2—H2	117.9 (19)	O3—C12—C11	109.5 (2)

## supplementary materials

C15—N3—C14	122.6 (2)	O3—C12—H12A	109.8
C15—N3—H3	119 (2)	C11—C12—H12A	109.8
C14—N3—H3	118 (2)	O3—C12—H12B	109.8
C16—N4—C15	127.9 (2)	C11—C12—H12B	109.8
C16—N4—H4	118.1 (18)	H12A—C12—H12B	108.2
C15—N4—H4	113.8 (18)	O3—C13—C14	108.6 (2)
C6—C1—C2	118.7 (2)	O3—C13—H13A	110.0
C6—C1—C7	123.7 (2)	C14—C13—H13A	110.0
C2—C1—C7	117.6 (2)	O3—C13—H13B	110.0
C3—C2—C1	120.8 (3)	C14—C13—H13B	110.0
C3—C2—H2A	119.6	H13A—C13—H13B	108.4
C1—C2—H2A	119.6	N3—C14—C13	111.0 (2)
C2—C3—C4	119.8 (3)	N3—C14—H14A	109.4
C2—C3—H3A	120.1	C13—C14—H14A	109.4
C4—C3—H3A	120.1	N3—C14—H14B	109.4
C5—C4—C3	119.9 (3)	C13—C14—H14B	109.4
C5—C4—H4A	120.0	H14A—C14—H14B	108.0
C3—C4—H4A	120.0	N3—C15—N4	116.7 (2)
C4—C5—C6	120.5 (3)	N3—C15—S2	124.06 (19)
C4—C5—H5	119.7	N4—C15—S2	119.18 (18)
C6—C5—H5	119.7	O4—C16—N4	121.1 (2)
C5—C6—C1	120.2 (3)	O4—C16—C17	121.1 (2)
C5—C6—H6	119.9	N4—C16—C17	117.8 (2)
C1—C6—H6	119.9	C22—C17—C18	119.0 (2)
O1—C7—N1	121.3 (2)	C22—C17—C16	124.1 (2)
O1—C7—C1	121.7 (2)	C18—C17—C16	116.9 (2)
N1—C7—C1	117.0 (2)	C19—C18—C17	120.5 (3)
N2—C8—N1	116.1 (2)	C19—C18—H18	119.8
N2—C8—S1	125.2 (2)	C17—C18—H18	119.8
N1—C8—S1	118.66 (18)	C20—C19—C18	120.4 (2)
N2—C9—C10	110.6 (2)	C20—C19—H19	119.8
N2—C9—H9A	109.5	C18—C19—H19	119.8
C10—C9—H9A	109.5	C19—C20—C21	120.1 (3)
N2—C9—H9B	109.5	C19—C20—H20	120.0
C10—C9—H9B	109.5	C21—C20—H20	120.0
H9A—C9—H9B	108.1	C20—C21—C22	119.7 (3)
O2—C10—C9	107.14 (19)	C20—C21—H21	120.1
O2—C10—H10A	110.3	C22—C21—H21	120.1
C9—C10—H10A	110.3	C17—C22—C21	120.3 (2)
O2—C10—H10B	110.3	C17—C22—H22	119.8
C9—C10—H10B	110.3	C21—C22—H22	119.8
O2—C11—C12—O3	65.0 (2)		

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H2 $\cdots$ O1	0.82 (3)	1.99 (3)	2.648 (3)	136 (2)
N3—H3 $\cdots$ O4	0.83 (3)	2.01 (3)	2.632 (3)	132 (3)
N4—H4 $\cdots$ O2 <sup>i</sup>	0.82 (3)	2.48 (3)	3.290 (3)	170 (2)

Symmetry codes: (i)  $-x+2, -y, -z+1$ .

Fig. 1




Fig. 2

