

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-(4-Carboxybenzyl)-1-methylpiperazin-1-ium picrate

Hongqi Li,^{a*} Q. N. M. Hakim Al-arique,^b
H. S. Yathirajan,^b B. Narayana^c and A. R. Ramesha^d^aKey Laboratory of Science and Technology of Eco-Textiles, Ministry of Education, College of Chemistry, Chemical Engineering and Biotechnology, Donghua University, Shanghai 201620, People's Republic of China, ^bDepartment of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India,^cDepartment of Studies in Chemistry, Mangalore University, Mangalagangotri 574 199, India, and ^dRL Fine Chem, Bangalore 560 064, India

Correspondence e-mail: hongqili@dhu.edu.cn


Received 27 December 2008; accepted 7 February 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.040; wR factor = 0.104; data-to-parameter ratio = 11.7.

The title compound, $\text{C}_{13}\text{H}_{19}\text{N}_2\text{O}_2^+ \cdot \text{C}_6\text{H}_2\text{N}_3\text{O}_7^-$, is a salt obtained by cocrystallization of 4-[(4-methylpiperazin-1-yl)methyl]benzoic acid and picric acid. The cations adopt an 'L-shaped' conformation and are linked into chains along [010] by $\text{O}-\text{H} \cdots \text{N}$ hydrogen bonds. The NH group of each piperazinium ring forms a hydrogen bond to the phenolate O atom of a picrate anion, and the picrate anions form face-to-face contacts with an interplanar separation of 3.023 (1) Å.

Related literature

For general background, see: Druker *et al.* (2001). For related structures, see: Swamy *et al.* (2007); Bindya *et al.* (2007); Sarojini *et al.* (2007); Wang & Jia (2008).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{19}\text{N}_2\text{O}_2^+ \cdot \text{C}_6\text{H}_2\text{N}_3\text{O}_7^-$
 $M_r = 463.41$
 Triclinic, $P\bar{1}$
 $a = 7.3020$ (12) Å

$b = 9.5993$ (16) Å
 $c = 15.131$ (3) Å
 $\alpha = 86.448$ (2)°
 $\beta = 79.145$ (2)°

$\gamma = 79.950$ (2)°
 $V = 1025.2$ (3) Å³
 $Z = 2$
 Mo $K\alpha$ radiation

$\mu = 0.12$ mm⁻¹
 $T = 296$ K
 $0.30 \times 0.20 \times 0.20$ mm

Data collection

Bruker SMART CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 2004)
 $T_{\min} = 0.965$, $T_{\max} = 0.976$

5373 measured reflections
 3565 independent reflections
 2745 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.016$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.104$
 $S = 1.06$
 3565 reflections
 305 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.18$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.17$ e Å⁻³

Table 1
 Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{O8}-\text{H8} \cdots \text{N4}^{\text{i}}$	0.82	1.79	2.6006 (19)	172
$\text{N5}-\text{H5A} \cdots \text{O1}^{\text{ii}}$	0.89 (2)	1.89 (2)	2.734 (2)	156.9 (18)

Symmetry codes: (i) $x, y + 1, z$; (ii) $-x + 1, -y + 2, -z + 1$.

Data collection: SMART (Bruker, 2001); cell refinement: SAINT (Bruker, 2001); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

QNMHA thanks R. L. Fine Chem, Bangalore, for a gift sample of 4-[(4-methylpiperazin-1-yl)methyl]benzoic acid. HSY thanks the University of Mysore for research facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: B12341).

References

- Bindya, S., Wong, W.-T., Ashok, M. A., Yathirajan, H. S. & Rathore, R. S. (2007). *Acta Cryst.* **C63**, o546–o548.
 Bruker (2001). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Druker, B. J., Sawyers, C. L., Kantarjian, H., Resta, D. J., Reese, S. F., Ford, J. M., Capdeville, R. & Talpaz, M. N. (2001). *N. Engl. J. Med.* **344**, 1038–1042.
 Sarojini, B. K., Yathirajan, H. S., Mustafa, K., Sarfraz, H. & Bolte, M. (2007). *Acta Cryst.* **E63**, o4477.
 Sheldrick, G. M. (2004). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Swamy, M. T., Ashok, M. A., Yathirajan, H. S., Narayana, B. & Bolte, M. (2007). *Acta Cryst.* **E63**, o4919.
 Wang, Z.-L. & Jia, L.-H. (2008). *Acta Cryst.* **E64**, o665–o666.

supplementary materials

Acta Cryst. (2009). E65, o518 [doi:10.1107/S1600536809004474]

4-(4-Carboxybenzyl)-1-methylpiperazin-1-ium picrate

H. Li, Q. N. M. Hakim Al-arique, H. S. Yathirajan, B. Narayana and A. R. Ramesha

Comment

4-[(4-Methylpiperazin-1-yl)methyl]benzoic acid is an intermediate in the synthesis of Imatinib, a drug used to treat certain types of cancer. Its mesylate salt is currently marketed by Novartis as Gleevec (Druker *et al.*, 2001). Picric acid forms salts or charge-transfer complexes with many organic compounds. Crystal structures of picrate complexes with organic compounds of pharmaceutical importance *viz.*, desipraminium picrate (Swamy *et al.*, 2007) and amitriptylinium picrate (Bindya *et al.*, 2007) have been reported. A three-dimensional network in piperazine-1,4-dium-picrate-piperazine (1/2/1) is reported recently (Wang & Jia, 2008).

Experimental

The title compound was prepared by taking equimolar quantities of picric acid (0.92 g, 2 mmol) and 4-[(4-methylpiperazin-1-yl)methyl]benzoic acid (0.47 g, 2 mmol) and dissolving them in water. The solution was stirred well at room temperature and slow evaporation of the solution resulted in the formation of the yellow coloured salt (yield 95%). Crystals suitable for single-crystal X-ray diffraction were grown from dimethyl formamide solvent.

Refinement

H atoms bound to C atoms were placed at calculated positions and refined using a riding model, with C—H = 0.93–0.97 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{C})$ for methyl H atoms. The H atom of the OH group was also placed geometrically and allowed to ride with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. The H atom of the NH group was located in a difference Fourier map and refined isotropically without restraint.

Figures




Fig. 1. Molecular structure with displacement ellipsoids drawn at the 30% probability level for non-H atoms.

4-(4-Carboxybenzyl)-1-methylpiperazin-1-ium picrate

Crystal data

$\text{C}_{13}\text{H}_{19}\text{N}_2\text{O}_2^+ \cdot \text{C}_6\text{H}_2\text{N}_3\text{O}_7^-$
 $M_r = 463.41$

$Z = 2$
 $F_{000} = 484$

supplementary materials

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 7.3020$ (12) Å

$b = 9.5993$ (16) Å

$c = 15.131$ (3) Å

$\alpha = 86.448$ (2)°

$\beta = 79.145$ (2)°

$\gamma = 79.950$ (2)°

$V = 1025.2$ (3) Å³

$D_x = 1.501$ Mg m⁻³

Melting point = 510–504 K

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 1962 reflections

$\theta = 2.5$ – 26.9 °

$\mu = 0.12$ mm⁻¹

$T = 296$ K

Block, colourless

$0.30 \times 0.20 \times 0.20$ mm

Data collection

Bruker SMART CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 296$ K

φ and ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 2004)

$T_{\min} = 0.965$, $T_{\max} = 0.976$

5373 measured reflections

3565 independent reflections

2745 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.016$

$\theta_{\max} = 25.0$ °

$\theta_{\min} = 2.2$ °

$h = -8$ → 6

$k = -11$ → 9

$l = -18$ → 18

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.040$

$wR(F^2) = 0.104$

$S = 1.06$

3565 reflections

305 parameters

Primary atom site location: structure-invariant direct
methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring
sites

H atoms treated by a mixture of
independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0447P)^2 + 0.2218P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.18$ e Å⁻³

$\Delta\rho_{\min} = -0.17$ e Å⁻³

Extinction correction: SHELXL97 (Sheldrick, 2008),
 $F_c^* = kFc[1 + 0.001x Fc^2\lambda^3/\sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.023 (2)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.5333 (3)	1.0778 (2)	0.14228 (14)	0.0426 (5)
C2	0.6431 (3)	1.0691 (2)	0.05175 (13)	0.0438 (5)
C3	0.7751 (3)	0.9547 (2)	0.02127 (13)	0.0430 (5)
H3	0.8395	0.9546	-0.0379	0.052*
C4	0.8126 (3)	0.8398 (2)	0.07793 (12)	0.0379 (4)
C5	0.7175 (2)	0.83776 (19)	0.16598 (12)	0.0359 (4)
H5	0.7422	0.7595	0.2039	0.043*
C6	0.5870 (3)	0.9521 (2)	0.19627 (12)	0.0372 (4)
C7	0.1636 (3)	1.33630 (19)	0.60024 (12)	0.0366 (4)
C8	0.1280 (2)	1.18702 (18)	0.60753 (11)	0.0320 (4)
C9	0.2178 (3)	1.09572 (19)	0.53938 (12)	0.0380 (4)
H9	0.2943	1.1292	0.4892	0.046*
C10	0.1949 (3)	0.95535 (19)	0.54527 (12)	0.0391 (4)
H10	0.2557	0.8954	0.4989	0.047*
C11	0.0818 (2)	0.90303 (18)	0.61993 (12)	0.0349 (4)
C12	-0.0108 (2)	0.99528 (18)	0.68722 (12)	0.0361 (4)
H12	-0.0897	0.9622	0.7367	0.043*
C13	0.0123 (2)	1.13606 (18)	0.68187 (12)	0.0348 (4)
H13	-0.0494	1.1964	0.7279	0.042*
C14	0.0608 (3)	0.74891 (19)	0.62770 (13)	0.0413 (5)
H14A	-0.0640	0.7413	0.6616	0.050*
H14B	0.0677	0.7156	0.5677	0.050*
C15	0.3984 (3)	0.6541 (2)	0.62183 (12)	0.0399 (5)
H15A	0.4281	0.7488	0.6208	0.048*
H15B	0.4063	0.6282	0.5601	0.048*
C16	0.5410 (3)	0.5522 (2)	0.66450 (12)	0.0415 (5)
H16A	0.5196	0.4562	0.6598	0.050*
H16B	0.6674	0.5591	0.6324	0.050*
C17	0.3292 (3)	0.5816 (2)	0.80965 (12)	0.0402 (5)
H17A	0.3189	0.6046	0.8721	0.048*
H17B	0.2985	0.4875	0.8083	0.048*
C18	0.1914 (3)	0.68674 (19)	0.76667 (12)	0.0375 (4)
H18A	0.0639	0.6846	0.7991	0.045*
H18B	0.2190	0.7813	0.7701	0.045*
C19	0.6653 (3)	0.4834 (2)	0.80485 (15)	0.0598 (6)
H19A	0.6595	0.5121	0.8651	0.090*
H19B	0.7903	0.4841	0.7710	0.090*
H19C	0.6355	0.3896	0.8065	0.090*
N1	0.6141 (3)	1.1856 (2)	-0.01420 (15)	0.0587 (5)
N2	0.9524 (2)	0.71935 (19)	0.04516 (11)	0.0471 (4)

supplementary materials

N3	0.4925 (2)	0.9427 (2)	0.28970 (11)	0.0469 (4)
N4	0.2038 (2)	0.65332 (14)	0.67160 (9)	0.0340 (4)
N5	0.5264 (2)	0.58329 (17)	0.76126 (10)	0.0375 (4)
O1	0.4059 (2)	1.17657 (15)	0.17135 (11)	0.0614 (4)
O2	0.5114 (3)	1.2956 (2)	0.00975 (14)	0.0921 (6)
O3	0.6952 (3)	1.16659 (19)	-0.09220 (12)	0.0799 (6)
O4	1.0379 (2)	0.72647 (18)	-0.03240 (10)	0.0680 (5)
O5	0.9804 (2)	0.61680 (16)	0.09632 (11)	0.0649 (5)
O6	0.4639 (2)	0.82634 (18)	0.32167 (10)	0.0633 (4)
O7	0.4470 (2)	1.05126 (19)	0.33248 (10)	0.0707 (5)
O8	0.1067 (2)	1.40458 (13)	0.67570 (8)	0.0445 (4)
H8	0.1397	1.4825	0.6691	0.067*
O9	0.2400 (2)	1.38891 (15)	0.53152 (9)	0.0605 (4)
H5A	0.551 (3)	0.669 (2)	0.7674 (13)	0.045 (6)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0386 (11)	0.0359 (11)	0.0586 (13)	-0.0106 (8)	-0.0166 (9)	-0.0054 (9)
C2	0.0462 (12)	0.0407 (11)	0.0508 (12)	-0.0163 (9)	-0.0195 (10)	0.0092 (9)
C3	0.0449 (11)	0.0537 (13)	0.0350 (10)	-0.0208 (10)	-0.0079 (9)	0.0021 (9)
C4	0.0359 (10)	0.0421 (11)	0.0355 (10)	-0.0087 (8)	-0.0032 (8)	-0.0036 (8)
C5	0.0377 (10)	0.0382 (10)	0.0345 (10)	-0.0097 (8)	-0.0102 (8)	-0.0007 (8)
C6	0.0368 (10)	0.0426 (11)	0.0348 (10)	-0.0100 (8)	-0.0079 (8)	-0.0059 (8)
C7	0.0393 (10)	0.0345 (10)	0.0353 (10)	-0.0079 (8)	-0.0036 (8)	-0.0002 (8)
C8	0.0339 (9)	0.0298 (9)	0.0334 (9)	-0.0060 (7)	-0.0086 (8)	-0.0001 (7)
C9	0.0457 (11)	0.0366 (10)	0.0307 (9)	-0.0071 (8)	-0.0045 (8)	0.0000 (8)
C10	0.0491 (11)	0.0327 (10)	0.0349 (10)	-0.0028 (8)	-0.0081 (9)	-0.0067 (8)
C11	0.0360 (10)	0.0296 (10)	0.0423 (10)	-0.0070 (7)	-0.0141 (8)	0.0004 (8)
C12	0.0311 (10)	0.0339 (10)	0.0415 (10)	-0.0064 (7)	-0.0026 (8)	0.0042 (8)
C13	0.0327 (9)	0.0311 (10)	0.0378 (10)	-0.0029 (7)	-0.0008 (8)	-0.0027 (8)
C14	0.0461 (11)	0.0317 (10)	0.0503 (11)	-0.0104 (8)	-0.0157 (9)	-0.0015 (8)
C15	0.0448 (11)	0.0371 (10)	0.0347 (10)	-0.0087 (8)	0.0013 (8)	0.0020 (8)
C16	0.0423 (11)	0.0371 (11)	0.0385 (10)	-0.0016 (8)	0.0047 (8)	-0.0018 (8)
C17	0.0392 (10)	0.0418 (11)	0.0348 (10)	-0.0040 (8)	0.0030 (8)	-0.0016 (8)
C18	0.0349 (10)	0.0371 (10)	0.0380 (10)	-0.0026 (8)	-0.0013 (8)	-0.0070 (8)
C19	0.0513 (13)	0.0579 (14)	0.0596 (14)	0.0144 (11)	-0.0087 (11)	0.0092 (11)
N1	0.0678 (13)	0.0507 (12)	0.0688 (14)	-0.0265 (10)	-0.0316 (11)	0.0184 (10)
N2	0.0420 (10)	0.0529 (11)	0.0439 (10)	-0.0091 (8)	0.0012 (8)	-0.0076 (8)
N3	0.0368 (9)	0.0614 (12)	0.0420 (9)	-0.0061 (8)	-0.0046 (7)	-0.0111 (9)
N4	0.0385 (8)	0.0289 (8)	0.0338 (8)	-0.0064 (6)	-0.0031 (6)	-0.0028 (6)
N5	0.0363 (9)	0.0317 (9)	0.0407 (9)	-0.0006 (7)	-0.0023 (7)	-0.0001 (7)
O1	0.0513 (9)	0.0413 (9)	0.0904 (12)	-0.0006 (7)	-0.0138 (8)	-0.0095 (8)
O2	0.1085 (16)	0.0526 (11)	0.1104 (16)	0.0045 (11)	-0.0328 (12)	0.0230 (11)
O3	0.1177 (16)	0.0729 (12)	0.0587 (11)	-0.0338 (11)	-0.0321 (11)	0.0242 (9)
O4	0.0656 (10)	0.0825 (12)	0.0461 (9)	-0.0069 (9)	0.0138 (8)	-0.0138 (8)
O5	0.0638 (10)	0.0524 (10)	0.0677 (10)	0.0059 (8)	-0.0006 (8)	0.0038 (8)
O6	0.0623 (10)	0.0701 (11)	0.0503 (9)	-0.0117 (8)	0.0055 (7)	0.0075 (8)

O7	0.0693 (11)	0.0822 (12)	0.0591 (10)	-0.0126 (9)	0.0035 (8)	-0.0371 (9)
O8	0.0599 (9)	0.0308 (7)	0.0409 (7)	-0.0161 (6)	0.0051 (6)	-0.0049 (6)
O9	0.0948 (12)	0.0441 (9)	0.0388 (8)	-0.0271 (8)	0.0105 (8)	0.0014 (6)

Geometric parameters (Å, °)

C1—O1	1.244 (2)	C14—H14B	0.970
C1—C2	1.450 (3)	C15—N4	1.479 (2)
C1—C6	1.453 (3)	C15—C16	1.509 (3)
C2—C3	1.369 (3)	C15—H15A	0.970
C2—N1	1.466 (3)	C15—H15B	0.970
C3—C4	1.375 (3)	C16—N5	1.493 (2)
C3—H3	0.930	C16—H16A	0.970
C4—C5	1.382 (2)	C16—H16B	0.970
C4—N2	1.447 (2)	C17—N5	1.490 (2)
C5—C6	1.362 (2)	C17—C18	1.505 (2)
C5—H5	0.930	C17—H17A	0.970
C6—N3	1.456 (2)	C17—H17B	0.970
C7—O9	1.209 (2)	C18—N4	1.476 (2)
C7—O8	1.313 (2)	C18—H18A	0.970
C7—C8	1.494 (2)	C18—H18B	0.970
C8—C9	1.385 (2)	C19—N5	1.489 (2)
C8—C13	1.391 (2)	C19—H19A	0.960
C9—C10	1.382 (3)	C19—H19B	0.960
C9—H9	0.930	C19—H19C	0.960
C10—C11	1.390 (3)	N1—O2	1.215 (3)
C10—H10	0.930	N1—O3	1.226 (3)
C11—C12	1.386 (2)	N2—O4	1.225 (2)
C11—C14	1.509 (2)	N2—O5	1.227 (2)
C12—C13	1.386 (2)	N3—O7	1.223 (2)
C12—H12	0.930	N3—O6	1.226 (2)
C13—H13	0.930	N5—H5A	0.89 (2)
C14—N4	1.491 (2)	O8—H8	0.820
C14—H14A	0.970		
O1—C1—C2	125.88 (19)	C16—C15—H15A	109.3
O1—C1—C6	122.99 (19)	N4—C15—H15B	109.3
C2—C1—C6	111.10 (16)	C16—C15—H15B	109.3
C3—C2—C1	123.83 (18)	H15A—C15—H15B	108.0
C3—C2—N1	115.70 (19)	N5—C16—C15	111.14 (14)
C1—C2—N1	120.46 (19)	N5—C16—H16A	109.4
C2—C3—C4	120.19 (18)	C15—C16—H16A	109.4
C2—C3—H3	119.9	N5—C16—H16B	109.4
C4—C3—H3	119.9	C15—C16—H16B	109.4
C3—C4—C5	120.72 (17)	H16A—C16—H16B	108.0
C3—C4—N2	119.81 (17)	N5—C17—C18	110.62 (15)
C5—C4—N2	119.47 (17)	N5—C17—H17A	109.5
C6—C5—C4	118.97 (17)	C18—C17—H17A	109.5
C6—C5—H5	120.5	N5—C17—H17B	109.5
C4—C5—H5	120.5	C18—C17—H17B	109.5

supplementary materials

C5—C6—C1	125.15 (17)	H17A—C17—H17B	108.1
C5—C6—N3	115.99 (17)	N4—C18—C17	110.48 (14)
C1—C6—N3	118.84 (16)	N4—C18—H18A	109.6
O9—C7—O8	123.05 (17)	C17—C18—H18A	109.6
O9—C7—C8	123.12 (16)	N4—C18—H18B	109.6
O8—C7—C8	113.82 (15)	C17—C18—H18B	109.6
C9—C8—C13	119.15 (16)	H18A—C18—H18B	108.1
C9—C8—C7	118.80 (16)	N5—C19—H19A	109.5
C13—C8—C7	122.01 (15)	N5—C19—H19B	109.5
C10—C9—C8	120.65 (17)	H19A—C19—H19B	109.5
C10—C9—H9	119.7	N5—C19—H19C	109.5
C8—C9—H9	119.7	H19A—C19—H19C	109.5
C9—C10—C11	120.59 (16)	H19B—C19—H19C	109.5
C9—C10—H10	119.7	O2—N1—O3	122.7 (2)
C11—C10—H10	119.7	O2—N1—C2	119.6 (2)
C12—C11—C10	118.64 (16)	O3—N1—C2	117.7 (2)
C12—C11—C14	120.67 (16)	O4—N2—O5	123.64 (17)
C10—C11—C14	120.69 (16)	O4—N2—C4	117.71 (17)
C13—C12—C11	121.00 (17)	O5—N2—C4	118.64 (16)
C13—C12—H12	119.5	O7—N3—O6	123.34 (18)
C11—C12—H12	119.5	O7—N3—C6	118.32 (18)
C12—C13—C8	119.95 (16)	O6—N3—C6	118.34 (17)
C12—C13—H13	120.0	C18—N4—C15	110.02 (14)
C8—C13—H13	120.0	C18—N4—C14	112.50 (13)
N4—C14—C11	115.44 (15)	C15—N4—C14	111.92 (14)
N4—C14—H14A	108.4	C17—N5—C19	111.33 (15)
C11—C14—H14A	108.4	C17—N5—C16	109.59 (15)
N4—C14—H14B	108.4	C19—N5—C16	112.03 (15)
C11—C14—H14B	108.4	C17—N5—H5A	106.5 (13)
H14A—C14—H14B	107.5	C19—N5—H5A	105.9 (13)
N4—C15—C16	111.45 (14)	C16—N5—H5A	111.3 (13)
N4—C15—H15A	109.3	C7—O8—H8	109.5
O1—C1—C2—C3	-176.28 (19)	C9—C8—C13—C12	0.4 (3)
C6—C1—C2—C3	2.0 (3)	C7—C8—C13—C12	-177.54 (16)
O1—C1—C2—N1	2.5 (3)	C12—C11—C14—N4	-90.9 (2)
C6—C1—C2—N1	-179.22 (16)	C10—C11—C14—N4	88.9 (2)
C1—C2—C3—C4	-1.1 (3)	N4—C15—C16—N5	-55.9 (2)
N1—C2—C3—C4	-179.94 (17)	N5—C17—C18—N4	59.5 (2)
C2—C3—C4—C5	0.3 (3)	C3—C2—N1—O2	-173.7 (2)
C2—C3—C4—N2	-179.94 (17)	C1—C2—N1—O2	7.4 (3)
C3—C4—C5—C6	-0.7 (3)	C3—C2—N1—O3	6.7 (3)
N2—C4—C5—C6	179.59 (16)	C1—C2—N1—O3	-172.20 (18)
C4—C5—C6—C1	1.9 (3)	C3—C4—N2—O4	2.3 (3)
C4—C5—C6—N3	-179.92 (16)	C5—C4—N2—O4	-177.91 (17)
O1—C1—C6—C5	175.93 (18)	C3—C4—N2—O5	-178.37 (18)
C2—C1—C6—C5	-2.4 (3)	C5—C4—N2—O5	1.4 (3)
O1—C1—C6—N3	-2.2 (3)	C5—C6—N3—O7	146.71 (18)
C2—C1—C6—N3	179.41 (16)	C1—C6—N3—O7	-35.0 (2)
O9—C7—C8—C9	13.3 (3)	C5—C6—N3—O6	-32.8 (2)

O8—C7—C8—C9	-165.86 (16)	C1—C6—N3—O6	145.57 (18)
O9—C7—C8—C13	-168.83 (19)	C17—C18—N4—C15	-58.31 (19)
O8—C7—C8—C13	12.1 (2)	C17—C18—N4—C14	176.17 (15)
C13—C8—C9—C10	-0.7 (3)	C16—C15—N4—C18	56.68 (19)
C7—C8—C9—C10	177.30 (16)	C16—C15—N4—C14	-177.48 (15)
C8—C9—C10—C11	-0.3 (3)	C11—C14—N4—C18	64.0 (2)
C9—C10—C11—C12	1.5 (3)	C11—C14—N4—C15	-60.5 (2)
C9—C10—C11—C14	-178.33 (16)	C18—C17—N5—C19	178.02 (17)
C10—C11—C12—C13	-1.8 (3)	C18—C17—N5—C16	-57.47 (19)
C14—C11—C12—C13	178.01 (16)	C15—C16—N5—C17	55.6 (2)
C11—C12—C13—C8	0.9 (3)	C15—C16—N5—C19	179.69 (17)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O8—H8 \cdots N4 ⁱ	0.82	1.79	2.6006 (19)	172
N5—H5A \cdots O1 ⁱⁱ	0.89 (2)	1.89 (2)	2.734 (2)	156.9 (18)

Symmetry codes: (i) $x, y+1, z$; (ii) $-x+1, -y+2, -z+1$.

Fig. 1

