

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 2,5-Dichloroanilinium chloride monohydrate

B. Thimme Gowda,<sup>a\*</sup> Sabine Foro,<sup>b</sup> B. S. Saraswathi,<sup>a</sup> Hiromitsu Terao<sup>c</sup> and Hartmut Fues<sup>b</sup>

<sup>a</sup>Department of Chemistry, Mangalore University, Mangalagangothri 574 199, Mangalore, India, <sup>b</sup>Institute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany, and <sup>c</sup>Faculty of Integrated Arts and Sciences, Tokushima University, Minamijosanjima-cho, Tokushima 770-8502, Japan  
Correspondence e-mail: gowdabt@yahoo.com


Received 4 February 2009; accepted 6 February 2009

Key indicators: single-crystal X-ray study;  $T = 299$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.049;  $wR$  factor = 0.143; data-to-parameter ratio = 13.4.

The title compound,  $\text{C}_6\text{H}_6\text{Cl}_2\text{N}^+\cdot\text{Cl}^-\cdot\text{H}_2\text{O}$ , is composed of discrete cations, chloride anions and water molecules, which are connected through  $\text{N}-\text{H}\cdots\text{Cl}$ ,  $\text{O}-\text{H}\cdots\text{Cl}$  and  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonding. Two H atoms of the positively charged  $-\text{NH}_3^+$  group have two chloride acceptors and the other one has the O atom of the water molecule as acceptor. The chloride anions form hydrogen bonds with two H atoms from two different water molecules and two H atoms from two positively charged  $-\text{NH}_3^+$  groups.

## Related literature

For water-free 2,5-dichloroanilinium chloride see: Gray & Jones (2002).



## Experimental

## Crystal data

 $\text{C}_6\text{H}_6\text{Cl}_2\text{N}^+\cdot\text{Cl}^-\cdot\text{H}_2\text{O}$  $M_r = 216.48$ 

Monoclinic,  $P2_1/n$   
 $a = 7.679$  (1) Å  
 $b = 6.476$  (1) Å  
 $c = 19.060$  (5) Å  
 $\beta = 96.95$  (3)°  
 $V = 940.9$  (3) Å<sup>3</sup>

$Z = 4$   
Cu  $K\alpha$  radiation  
 $\mu = 8.39$  mm<sup>-1</sup>  
 $T = 299$  (2) K  
 $0.35 \times 0.30 \times 0.10$  mm

## Data collection

Enraf–Nonius CAD-4 diffractometer  
Absorption correction:  $\psi$  scan (North *et al.*, 1968)  
 $T_{\min} = 0.109$ ,  $T_{\max} = 0.432$   
3331 measured reflections

1669 independent reflections  
1421 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.068$   
3 standard reflections  
frequency: 120 min  
intensity decay: 1%

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$   
 $wR(F^2) = 0.143$   
 $S = 1.10$   
1669 reflections  
125 parameters

3 restraints  
Only H-atom coordinates refined  
 $\Delta\rho_{\text{max}} = 0.39$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.49$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H11}\cdots\text{Cl3}^{\text{i}}$	0.92 (4)	2.24 (4)	3.123 (3)	162 (3)
$\text{N1}-\text{H12}\cdots\text{Cl3}$	0.94 (4)	2.16 (4)	3.099 (3)	172 (3)
$\text{N1}-\text{H13}\cdots\text{O1}^{\text{ii}}$	0.88 (4)	1.82 (4)	2.699 (4)	175 (4)
$\text{O1}-\text{H1A}\cdots\text{Cl3}$	0.85 (3)	2.37 (3)	3.212 (3)	172 (4)
$\text{O1}-\text{H1B}\cdots\text{Cl3}^{\text{iii}}$	0.83 (3)	2.34 (3)	3.158 (3)	169 (4)

Symmetry codes: (i)  $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $x + 1, y, z$ ; (iii)  $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$ .

Data collection: *CAD-4-PC* (Enraf–Nonius, 1996); cell refinement: *CAD-4-PC*; data reduction: *REDU4* (Stoe & Cie, 1987); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2003); software used to prepare material for publication: *SHELXL97*.

BTG thanks the Alexander von Humboldt Foundation, Bonn, Germany, for extensions of his research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2865).

## References

- Enraf–Nonius (1996). *CAD-4-PC*. Enraf–Nonius, Delft, The Netherlands.  
Gray, L. & Jones, P. G. (2002). *Z. Naturforsch. Teil B*, **57**, 73–82.  
North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.  
Stoe & Cie (1987). *REDU4*. Stoe & Cie, Darmstadt, Germany.

**supplementary materials**

*Acta Cryst.* (2009). E65, o516 [ doi:10.1107/S1600536809004395 ]

## 2,5-Dichloroanilinium chloride monohydrate

B. T. Gowda, S. Foro, B. S. Saraswathi, H. Terao and H. Fuess

### Comment

The crystal structure of water free 2,5-dichloroanilinium chloride has been reported (Gray & Jones, 2002). We report herein the crystal structure of 2,5-dichloroanilinium chloride monohydrate. The title compound showed interesting H-bonding in its crystal structure (Fig. 1). Two H-atoms of the positively charged NH<sub>3</sub> group have two chloride acceptors and the other H has O atom acceptor of the water molecule, while chloride anions are linked by four-center hydrogen bonds, with each chloride forming H-bonding with two H-atoms, one each from two different water molecules and two H-atoms, one each from two positively charged NH<sub>3</sub> groups. This is in comparison with the usual set of hydrogen bonds from NH<sub>3</sub> to chloride leading to layer structure observed with water free 2,5-dichloroanilinium chloride (Gray & Jones, 2002), with a short Cl1...Cl3 contact. Further, the water free structure involved four weak interactions, namely the three hydrogen bonds H4...Cl3, H6...Cl2 and H3...Cl1 and the chlorine-chlorine interaction Cl2...Cl3. The crystal packing of (I) through N—H...Cl, O—H...Cl and N—H...O hydrogen bonding (Table 1) is shown in Fig.2

### Experimental

The solution of pure 2,5-dichloroaniline (0.02 mole) in ethanol (20 cc) was treated dropwise with dilute hydrochloric acid (>0.025 mole) with constant stirring. The resulting mixture was slowly evaporated at room temperature to obtain 2,5-dichloroanilinium hydrochloride monohydrate. The resultant solid was recrystallized to constant melting point from ethanol. The single crystals used in X-ray diffraction studies were grown in ethanolic solution by slow evaporation at room temperature.

### Refinement

H atoms were located in a difference map, and their positional parameters were refined freely except for the water H atoms which were refined with the O—H distances restrained to 0.85 (4) Å and H—H distance restrained to 1.37 (4)Å. All H atoms were refined with isotropic displacement parameters set to 1.2 times of the  $U_{eq}$  of the parent atom.

### Figures




Fig. 1. Molecular structure of the title compound, showing the atom labeling scheme. The displacement ellipsoids are drawn at the 50% probability level. The H atoms are represented as small spheres of arbitrary radii.




Fig. 2. Molecular packing of the title compound with hydrogen bonding shown as dashed lines.

## 2,5-Dichloroanilinium chloride monohydrate

### Crystal data

$C_6H_6Cl_2N^+ \cdot Cl^- \cdot H_2O$

$M_r = 216.48$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 7.679$  (1) Å

$b = 6.476$  (1) Å

$c = 19.060$  (5) Å

$\beta = 96.95$  (3)°

$V = 940.9$  (3) Å<sup>3</sup>

$Z = 4$

$F_{000} = 440$

$D_x = 1.528$  Mg m<sup>-3</sup>

Cu  $K\alpha$  radiation

$\lambda = 1.54180$  Å

Cell parameters from 25 reflections

$\theta = 6.0$ – $20.3$ °

$\mu = 8.39$  mm<sup>-1</sup>

$T = 299$  K

Plate, colourless

$0.35 \times 0.30 \times 0.10$  mm

### Data collection

Enraf–Nonius CAD-4  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 299$  K

$\omega/2\theta$  scans

Absorption correction:  $\psi$  scan  
(North *et al.*, 1968)

$T_{\min} = 0.109$ ,  $T_{\max} = 0.432$

3331 measured reflections

1669 independent reflections

1421 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.068$

$\theta_{\max} = 67.2$ °

$\theta_{\min} = 4.7$ °

$h = -9 \rightarrow 9$

$k = 0 \rightarrow 7$

$l = -22 \rightarrow 22$

3 standard reflections

every 120 min

intensity decay: 1.0%

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.049$

$wR(F^2) = 0.143$

$S = 1.10$

1669 reflections

Secondary atom site location: difference Fourier map

Hydrogen site location: difference Fourier map

Only H-atom coordinates refined

$w = 1/[\sigma^2(F_o^2) + (0.0831P)^2 + 0.169P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.012$

$\Delta\rho_{\max} = 0.39$  e Å<sup>-3</sup>

125 parameters

$$\Delta\rho_{\min} = -0.49 \text{ e } \text{\AA}^{-3}$$

3 restraints

Extinction correction: SHELXL97 (Sheldrick, 2008),

$$F_c^* = kFc[1+0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$$

Primary atom site location: structure-invariant direct methods

Extinction coefficient: 0.0126 (16)

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.64647 (11)	-0.35617 (13)	0.11963 (6)	0.0720 (4)
C12	0.84843 (12)	0.33419 (16)	-0.08256 (4)	0.0716 (4)
N1	0.8342 (3)	0.0359 (4)	0.16645 (11)	0.0476 (6)
H11	0.872 (4)	-0.079 (6)	0.1920 (19)	0.057*
H12	0.736 (5)	0.083 (5)	0.1868 (18)	0.057*
H13	0.917 (5)	0.128 (6)	0.179 (2)	0.057*
C1	0.7922 (3)	0.0096 (5)	0.09016 (13)	0.0441 (6)
C2	0.7056 (4)	-0.1657 (5)	0.06338 (18)	0.0529 (7)
C3	0.6664 (4)	-0.1873 (6)	-0.0093 (2)	0.0635 (9)
H3	0.604 (5)	-0.303 (7)	-0.024 (2)	0.076*
C4	0.7107 (4)	-0.0333 (6)	-0.05366 (16)	0.0626 (9)
H4	0.681 (5)	-0.046 (6)	-0.100 (2)	0.075*
C5	0.7943 (4)	0.1398 (5)	-0.02646 (15)	0.0534 (7)
C6	0.8371 (4)	0.1651 (5)	0.04598 (15)	0.0474 (6)
H6	0.905 (4)	0.288 (5)	0.065 (2)	0.057*
O1	0.0902 (3)	0.3197 (4)	0.19565 (15)	0.0697 (7)
H1A	0.200 (4)	0.301 (6)	0.206 (3)	0.084*
H1B	0.067 (5)	0.413 (6)	0.223 (2)	0.084*
Cl3	0.50259 (9)	0.21839 (12)	0.22100 (4)	0.0565 (3)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C11	0.0655 (6)	0.0603 (5)	0.0883 (6)	-0.0101 (3)	0.0009 (5)	0.0022 (4)
C12	0.0671 (6)	0.0981 (7)	0.0499 (4)	0.0101 (4)	0.0085 (4)	0.0145 (4)
N1	0.0485 (13)	0.0551 (13)	0.0379 (10)	0.0017 (11)	-0.0002 (9)	-0.0007 (10)
C1	0.0377 (13)	0.0528 (14)	0.0402 (11)	0.0065 (11)	-0.0015 (10)	-0.0062 (11)

## supplementary materials

C2	0.0400 (13)	0.0555 (16)	0.0615 (17)	0.0031 (12)	-0.0004 (13)	-0.0084 (13)
C3	0.0500 (17)	0.0694 (19)	0.0677 (19)	0.0019 (15)	-0.0065 (15)	-0.0270 (17)
C4	0.0522 (16)	0.086 (2)	0.0470 (14)	0.0111 (16)	-0.0034 (13)	-0.0205 (15)
C5	0.0436 (14)	0.076 (2)	0.0405 (13)	0.0130 (13)	0.0032 (12)	-0.0027 (13)
C6	0.0415 (13)	0.0578 (15)	0.0419 (13)	0.0044 (12)	0.0009 (11)	-0.0054 (12)
O1	0.0571 (14)	0.0722 (16)	0.0776 (15)	-0.0017 (11)	-0.0004 (12)	-0.0175 (12)
Cl3	0.0529 (5)	0.0671 (5)	0.0484 (4)	-0.0018 (3)	0.0015 (3)	-0.0101 (3)

### Geometric parameters (Å, °)

C1—C2	1.731 (3)	C3—C4	1.377 (6)
C1—C5	1.735 (3)	C3—H3	0.92 (4)
N1—C1	1.461 (3)	C4—C5	1.362 (5)
N1—H11	0.92 (4)	C4—H4	0.88 (4)
N1—H12	0.94 (4)	C5—C6	1.390 (4)
N1—H13	0.88 (4)	C6—H6	1.00 (4)
C1—C2	1.382 (4)	O1—H1A	0.85 (3)
C1—C6	1.383 (4)	O1—H1B	0.83 (3)
C2—C3	1.388 (5)		
C1—N1—H11	117 (2)	C4—C3—H3	124 (3)
C1—N1—H12	111 (2)	C2—C3—H3	116 (3)
H11—N1—H12	105 (3)	C5—C4—C3	120.1 (3)
C1—N1—H13	114 (2)	C5—C4—H4	121 (3)
H11—N1—H13	104 (3)	C3—C4—H4	119 (3)
H12—N1—H13	105 (3)	C4—C5—C6	121.3 (3)
C2—C1—C6	121.2 (3)	C4—C5—Cl2	120.0 (2)
C2—C1—N1	120.1 (3)	C6—C5—Cl2	118.8 (3)
C6—C1—N1	118.7 (3)	C1—C6—C5	118.2 (3)
C1—C2—C3	119.2 (3)	C1—C6—H6	121 (2)
C1—C2—Cl1	120.5 (2)	C5—C6—H6	121 (2)
C3—C2—Cl1	120.3 (3)	H1A—O1—H1B	104 (4)
C4—C3—C2	120.0 (3)		
C6—C1—C2—C3	1.3 (4)	C3—C4—C5—C6	0.2 (5)
N1—C1—C2—C3	179.8 (3)	C3—C4—C5—Cl2	-179.5 (3)
C6—C1—C2—Cl1	-178.6 (2)	C2—C1—C6—C5	-0.6 (4)
N1—C1—C2—Cl1	-0.2 (4)	N1—C1—C6—C5	-179.1 (3)
C1—C2—C3—C4	-1.3 (5)	C4—C5—C6—C1	-0.2 (4)
Cl1—C2—C3—C4	178.7 (3)	Cl2—C5—C6—C1	179.6 (2)
C2—C3—C4—C5	0.5 (5)		

### Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N1—H11 $\cdots$ Cl3 <sup>i</sup>	0.92 (4)	2.24 (4)	3.123 (3)	162 (3)
N1—H12 $\cdots$ Cl3	0.94 (4)	2.16 (4)	3.099 (3)	172 (3)
N1—H13 $\cdots$ O1 <sup>ii</sup>	0.88 (4)	1.82 (4)	2.699 (4)	175 (4)
O1—H1A $\cdots$ Cl3	0.85 (3)	2.37 (3)	3.212 (3)	172 (4)
O1—H1B $\cdots$ Cl3 <sup>iii</sup>	0.83 (3)	2.34 (3)	3.158 (3)	169 (4)

Symmetry codes: (i)  $-x+3/2, y-1/2, -z+1/2$ ; (ii)  $x+1, y, z$ ; (iii)  $-x+1/2, y+1/2, -z+1/2$ .

Fig. 1




Fig. 2

