

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

[*N,N'*-Bis(4-bromobenzylidene)-2,2-dimethylpropane- κ^2N,N']iodidocopper(I). Corrigendum

Reza Kia,^a Hoong-Kun Fun^{a*} and Hadi Kargar^b

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bDepartment of Chemistry, School of Science, Payame Noor University (PNU), Ardakan, Yazd, Iran
Correspondence e-mail: hkfun@usm.my

Received 16 June 2009; accepted 17 June 2009

The chemical name of the title compound in the paper by Kia, Fun & Kargar [*Acta Cryst.* (2009), **E65**, m289] is corrected.

In the paper by Kia, Fun & Kargar [*Acta Cryst.* (2009), **E65**, m289], the chemical name given in the *Title* should be '*[N,N'*-Bis(4-bromobenzylidene)-2,2-dimethylpropane-1,3-diamine- κ^2N,N']iodidocopper(I)'

[*N,N'*-Bis(4-bromobenzylidene)-2,2-dimethylpropane- κ^2 *N,N'*]iodidocopper(I)Reza Kia,^a Hoong-Kun Fun^{a*} and Hadi Kargar^b^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bDepartment of Chemistry, School of Science, Payame Noor University (PNU), Ardakan, Yazd, Iran

Correspondence e-mail: hkfun@usm.my


Received 21 January 2009; accepted 12 February 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.012$ Å; R factor = 0.050; wR factor = 0.112; data-to-parameter ratio = 16.0.

The title compound, $[\text{CuI}(\text{C}_{19}\text{H}_{20}\text{Br}_2\text{N}_2)]$, lies across a crystallographic mirror plane. The coordination around the copper centre is distorted trigonal planar, with a bite angle of $94.7(3)^\circ$. A six-membered chelate ring in a chair conformation is formed by the coordination of the imine N atoms of the bidentate ligand to the Cu^{I} atom. This conformation is required by the crystallographic mirror symmetry. The interplanar angle between the benzene rings is $74.85(19)^\circ$. The crystal structure exhibits weak intermolecular $\text{C}-\text{H}\cdots\pi$ interactions, which link the molecules into chains along the b axis.

Related literature

For the puckering parameters, see: Cremer & Pople (1975). For related literature and the catalytic applications see, for example: Killian *et al.* (1996); Jung *et al.* (1996); Small *et al.* (1998). For a related structure, see: Kia *et al.* (2009). For the stability of the temperature controller, see Cosier & Glazer (1986).

**Experimental***Crystal data*

$[\text{CuI}(\text{C}_{19}\text{H}_{20}\text{Br}_2\text{N}_2)]$
 $M_r = 626.63$
 Monoclinic, $C2/m$
 $a = 16.2224(15)$ Å
 $b = 12.2807(12)$ Å
 $c = 10.6292(12)$ Å
 $\beta = 91.599(6)^\circ$

$V = 2116.8(4)$ Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 6.27$ mm⁻¹
 $T = 100$ K
 $0.58 \times 0.09 \times 0.05$ mm

Data collection

Bruker SMART APEXII CCD
 area-detector diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 2005)
 $T_{\text{min}} = 0.119$, $T_{\text{max}} = 0.714$

10374 measured reflections
 1936 independent reflections
 1474 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.099$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.112$
 $S = 1.16$
 1936 reflections

121 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 2.64$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.99$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the C1–C6 benzene ring.

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|---|-------|-------------|-------------|---------------|
| $\text{C8}-\text{H8A}\cdots\text{Cg1}^{\text{i}}$ | 0.99 | 2.83 | 3.631 (9) | 138 |

Symmetry code: (i) $-x + \frac{1}{2}, -y + \frac{1}{2}, -z$.

Data collection: APEX2 (Bruker, 2005); cell refinement: APEX2; data reduction: SAINT (Bruker, 2005); program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

HKF and RK thank the Malaysian Government and Universiti Sains Malaysia for Science Fund grant No. 305/PFIZIK/613312. RK thanks Universiti Sains Malaysia for a post-doctoral research fellowship. HK thanks PNU for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: EZZ2161).

References

- Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
 Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
 Jung, B., Karlin, K. D. & Zuberbühler, A. D. (1996). *J. Am. Chem. Soc.* **118**, 3763–3768.
 Kia, R., Fun, H.-K. & Kargar, H. (2009). *Acta Cryst.* **E65**, m197.
 Killian, C. M., Tempel, D. J., Johnson, L. K. & Brookhart, M. (1996). *J. Am. Chem. Soc.* **118**, 11664–11670.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Small, B. L., Brookhart, M. & Bennett, A. M. A. (1998). *J. Am. Chem. Soc.* **120**, 4049–4054.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2009). E65, m289 [doi:10.1107/S1600536809005078]

[*N,N'*-Bis(4-bromobenzylidene)-2,2-dimethylpropane- κ^2 *N,N'*]iodidocopper(I)

R. Kia, H.-K. Fun and H. Kargar

Comment

In recent years, an increasing amount of research has been focused on the design and preparation of mono- or dinuclear mixed ligand transition metal complexes with neutral, chelating nitrogen-containing ligands. Early and late transition metal complexes of this type have been extensively used as catalysts for a wide categories of reactions, including olefin polymerization (Killian *et al.*, 1996) and oxygen activation (Jung *et al.*, 1996). In this context, diverse chelating Schiff base type ligands, amines and pyridine derivatives (Small *et al.*, 1998) have successfully been applied in the preparation of these homogeneous catalysts. Here we report the crystal structure of an aldimine Schiff base ligand with copper(I) iodide. To the best of our knowledge, only one such related compound has been published (Kia *et al.*, 2009). The title compound is the second tricoordinate complex of an aldimine bis-Schiff base ligand with copper(I) iodide adopting trigonal planar geometry.

The title compound, I, Fig. 1, lies across a crystallographic mirror plane. Atoms I1, Cu1, C9, C10 and C11 lies on this mirror plane. The asymmetric unit of (I) is composed of one-half of the molecule. The coordination around the copper centre is distorted trigonal planar, with a bite angle of 94.7 (3)°. The deviation of the Cu atom from the N1/N1A/I1 plane is -0.105 (4) Å. A six-membered chelate ring with a chair conformation is formed by the coordination of iminic N atoms of the bidentate ligand to the Cu(I) atom, with ring puckering parameters (Cremer & Pople, 1975) of $Q = 0.696$ (7) Å, $\Theta = 172.2$ (6)°, $\Phi = 180$ (5)°. This conformation is required if the local symmetry of the metal coordination site is in accordance with the mirror plane that passes through the metal atom normal to the line connecting the nitrogen atoms. The dihedral angle between the phenyl rings is 74.85 (19)°. The crystal structure is stabilized by weak intermolecular C—H $\cdots\pi$ interactions (Cg1 is the centroid of the C1–C6 benzene ring) which link the molecules into chains along the *b*-axis (Fig. 2 and Table 1).

Experimental

N,N'-Bis(4-bromobenzylidene)-2,2-dimethylpropane (783 mg, 2 mmol) was added dropwise to a suspension of CuI (380 mg, 2.0 mmol) in 50 ml of THF. After 15 min a clear yellowish solution was obtained. The volume of the reaction mixture was reduced until the formation of a yellow precipitate occurred. Single crystals suitable for X-ray diffraction were grown from the acetonitrile solution.

Refinement

All H atoms were positioned geometrically with C—H = 0.95–0.99 Å and refined in a riding model approximation with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The highest peak (2.64 e. Å⁻³) is located 1.02 Å from I1 and the deepest hole (-0.99 e. Å⁻³) is located 0.58 Å from H10A.

Figures




Fig. 1. The molecular structure of (I), showing 40% probability displacement ellipsoids and the atomic numbering. Symmetry code for A atoms; $x, -y + 1, z$.




Fig. 2. The crystal packing of (I), viewed down the c -axis, showing C—H... π interactions linking the molecules into chains along the b -axis.

[*N,N'*-Bis(4-bromobenzylidene)-2,2-dimethylpropane- κ^2 *N,N'*]iodidocopper(I)

Crystal data

[CuI(C₁₉H₂₀Br₂N₂)]

$M_r = 626.63$

Monoclinic, $C2/m$

Hall symbol: $-C 2y$

$a = 16.2224 (15) \text{ \AA}$

$b = 12.2807 (12) \text{ \AA}$

$c = 10.6292 (12) \text{ \AA}$

$\beta = 91.599 (6)^\circ$

$V = 2116.8 (4) \text{ \AA}^3$

$Z = 4$

$F_{000} = 1200$

$D_x = 1.966 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 4080 reflections

$\theta = 2.5\text{--}29.5^\circ$

$\mu = 6.27 \text{ mm}^{-1}$

$T = 100 \text{ K}$

Needle, yellow

$0.58 \times 0.09 \times 0.05 \text{ mm}$

Data collection

Bruker SMART APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 100 \text{ K}$

φ and ω scans

Absorption correction: multi-scan

(SADABS; Bruker, 2005)

$T_{\min} = 0.119, T_{\max} = 0.714$

10374 measured reflections

1936 independent reflections

1474 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.099$

$\theta_{\text{max}} = 25.0^\circ$

$\theta_{\text{min}} = 2.1^\circ$

$h = -19 \rightarrow 19$

$k = -14 \rightarrow 14$

$l = -12 \rightarrow 12$

Refinement

| | |
|--|--|
| Refinement on F^2 | Secondary atom site location: difference Fourier map |
| Least-squares matrix: full | Hydrogen site location: inferred from neighbouring sites |
| $R[F^2 > 2\sigma(F^2)] = 0.050$ | H-atom parameters constrained |
| $wR(F^2) = 0.112$ | $w = 1/[\sigma^2(F_o^2) + (0.0405P)^2 + 16.0151P]$ |
| $S = 1.16$ | where $P = (F_o^2 + 2F_c^2)/3$ |
| 1936 reflections | $(\Delta/\sigma)_{\max} < 0.001$ |
| 121 parameters | $\Delta\rho_{\max} = 2.64 \text{ e } \text{\AA}^{-3}$ |
| Primary atom site location: structure-invariant direct methods | $\Delta\rho_{\min} = -0.99 \text{ e } \text{\AA}^{-3}$ |
| | Extinction correction: none |

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|-------------|-------------|--------------|----------------------------------|
| I1 | 0.38486 (4) | 0.5000 | 0.14862 (6) | 0.0166 (2) |
| Br1 | 0.11472 (5) | 0.16378 (8) | 0.54336 (7) | 0.0309 (3) |
| Cu1 | 0.25251 (8) | 0.5000 | 0.02730 (11) | 0.0168 (3) |
| N1 | 0.1853 (3) | 0.3798 (6) | -0.0519 (5) | 0.0167 (14) |
| C1 | 0.1751 (5) | 0.3315 (7) | 0.2223 (7) | 0.0213 (18) |
| H1A | 0.2085 | 0.3923 | 0.2022 | 0.026* |
| C2 | 0.1683 (4) | 0.3023 (7) | 0.3463 (7) | 0.0210 (18) |
| H2A | 0.1957 | 0.3426 | 0.4113 | 0.025* |
| C3 | 0.1204 (5) | 0.2130 (7) | 0.3742 (6) | 0.0193 (18) |
| C4 | 0.0779 (5) | 0.1560 (7) | 0.2811 (8) | 0.026 (2) |
| H4A | 0.0442 | 0.0958 | 0.3021 | 0.031* |
| C5 | 0.0852 (4) | 0.1878 (7) | 0.1556 (7) | 0.0188 (18) |
| H5A | 0.0559 | 0.1496 | 0.0907 | 0.023* |
| C6 | 0.1354 (4) | 0.2756 (7) | 0.1260 (7) | 0.0172 (18) |
| C7 | 0.1428 (4) | 0.3030 (7) | -0.0085 (7) | 0.0184 (17) |
| H7A | 0.1130 | 0.2590 | -0.0675 | 0.022* |

supplementary materials

| | | | | |
|------|------------|------------|--------------|-------------|
| C8 | 0.1816 (5) | 0.3967 (7) | -0.1892 (7) | 0.0213 (18) |
| H8A | 0.1542 | 0.3330 | -0.2292 | 0.026* |
| H8B | 0.2387 | 0.3997 | -0.2198 | 0.026* |
| C9 | 0.1360 (7) | 0.5000 | -0.2327 (9) | 0.020 (3) |
| C10 | 0.0477 (6) | 0.5000 | -0.1869 (11) | 0.028 (3) |
| H10A | 0.0492 | 0.5000 | -0.0947 | 0.041* |
| H10B | 0.0187 | 0.4348 | -0.2177 | 0.041* |
| C11 | 0.1346 (7) | 0.5000 | -0.3771 (10) | 0.024 (3) |
| H11A | 0.1916 | 0.5000 | -0.4055 | 0.036* |
| H11B | 0.1061 | 0.5652 | -0.4091 | 0.036* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|------------|------------|------------|-------------|-------------|------------|
| II | 0.0191 (4) | 0.0171 (4) | 0.0134 (4) | 0.000 | -0.0029 (3) | 0.000 |
| Br1 | 0.0371 (5) | 0.0400 (6) | 0.0154 (4) | -0.0155 (4) | -0.0026 (3) | 0.0048 (4) |
| Cu1 | 0.0195 (7) | 0.0188 (8) | 0.0121 (7) | 0.000 | -0.0020 (5) | 0.000 |
| N1 | 0.020 (3) | 0.018 (4) | 0.012 (3) | 0.001 (3) | 0.001 (2) | -0.002 (3) |
| C1 | 0.024 (4) | 0.021 (5) | 0.019 (4) | -0.007 (3) | 0.002 (3) | -0.003 (4) |
| C2 | 0.025 (4) | 0.023 (5) | 0.015 (4) | -0.009 (4) | -0.004 (3) | -0.007 (4) |
| C3 | 0.031 (4) | 0.022 (5) | 0.005 (4) | -0.003 (4) | 0.001 (3) | 0.002 (3) |
| C4 | 0.031 (5) | 0.022 (5) | 0.025 (5) | -0.010 (4) | 0.002 (4) | 0.003 (4) |
| C5 | 0.018 (4) | 0.025 (5) | 0.013 (4) | -0.001 (3) | -0.002 (3) | -0.002 (4) |
| C6 | 0.019 (4) | 0.014 (5) | 0.020 (4) | 0.005 (3) | 0.004 (3) | -0.002 (4) |
| C7 | 0.022 (4) | 0.016 (5) | 0.017 (4) | 0.000 (3) | -0.005 (3) | -0.010 (4) |
| C8 | 0.031 (4) | 0.019 (5) | 0.014 (4) | -0.003 (4) | 0.001 (3) | -0.002 (4) |
| C9 | 0.029 (6) | 0.024 (7) | 0.007 (5) | 0.000 | 0.003 (4) | 0.000 |
| C10 | 0.022 (6) | 0.034 (8) | 0.027 (7) | 0.000 | -0.009 (5) | 0.000 |
| C11 | 0.038 (7) | 0.020 (7) | 0.015 (6) | 0.000 | -0.006 (5) | 0.000 |

Geometric parameters (\AA , $^\circ$)

| | | | |
|------------------------|-------------|--------------------|------------|
| II—Cu1 | 2.4735 (14) | C5—C6 | 1.392 (11) |
| Br1—C3 | 1.902 (7) | C5—H5A | 0.9500 |
| Cu1—N1 | 2.006 (6) | C6—C7 | 1.477 (11) |
| Cu1—N1 ⁱ | 2.006 (6) | C7—H7A | 0.9500 |
| N1—C7 | 1.263 (10) | C8—C9 | 1.533 (10) |
| N1—C8 | 1.474 (9) | C8—H8A | 0.9900 |
| C1—C2 | 1.374 (11) | C8—H8B | 0.9900 |
| C1—C6 | 1.378 (10) | C9—C10 | 1.527 (15) |
| C1—H1A | 0.9500 | C9—C8 ⁱ | 1.533 (10) |
| C2—C3 | 1.381 (11) | C9—C11 | 1.535 (14) |
| C2—H2A | 0.9500 | C10—H10A | 0.9800 |
| C3—C4 | 1.381 (11) | C10—H10B | 0.9800 |
| C4—C5 | 1.398 (11) | C11—H11A | 0.9800 |
| C4—H4A | 0.9500 | C11—H11B | 0.9801 |
| N1—Cu1—N1 ⁱ | 94.8 (4) | C5—C6—C7 | 117.3 (7) |
| N1—Cu1—II | 132.24 (18) | N1—C7—C6 | 125.7 (7) |

| | | | |
|----------------------------|-------------|--------------------------|------------|
| N1 ⁱ —Cu1—I1 | 132.24 (18) | N1—C7—H7A | 117.1 |
| C7—N1—C8 | 117.4 (6) | C6—C7—H7A | 117.1 |
| C7—N1—Cu1 | 133.8 (5) | N1—C8—C9 | 114.9 (7) |
| C8—N1—Cu1 | 108.5 (5) | N1—C8—H8A | 108.5 |
| C2—C1—C6 | 122.3 (8) | C9—C8—H8A | 108.5 |
| C2—C1—H1A | 118.9 | N1—C8—H8B | 108.5 |
| C6—C1—H1A | 118.9 | C9—C8—H8B | 108.5 |
| C1—C2—C3 | 118.3 (7) | H8A—C8—H8B | 107.5 |
| C1—C2—H2A | 120.9 | C10—C9—C8 ⁱ | 110.7 (6) |
| C3—C2—H2A | 120.9 | C10—C9—C8 | 110.7 (6) |
| C4—C3—C2 | 121.4 (7) | C8 ⁱ —C9—C8 | 111.7 (9) |
| C4—C3—Br1 | 118.7 (6) | C10—C9—C11 | 109.3 (9) |
| C2—C3—Br1 | 119.8 (6) | C8 ⁱ —C9—C11 | 107.1 (6) |
| C3—C4—C5 | 119.2 (8) | C8—C9—C11 | 107.1 (6) |
| C3—C4—H4A | 120.4 | C9—C10—H10A | 108.7 |
| C5—C4—H4A | 120.4 | C9—C10—H10B | 109.8 |
| C6—C5—C4 | 119.8 (7) | H10A—C10—H10B | 109.5 |
| C6—C5—H5A | 120.1 | C9—C11—H11A | 108.6 |
| C4—C5—H5A | 120.1 | C9—C11—H11B | 109.9 |
| C1—C6—C5 | 118.9 (7) | H11A—C11—H11B | 109.5 |
| C1—C6—C7 | 123.8 (7) | | |
| N1 ⁱ —Cu1—N1—C7 | -119.0 (7) | C4—C5—C6—C1 | -1.9 (11) |
| I1—Cu1—N1—C7 | 70.2 (8) | C4—C5—C6—C7 | 178.0 (7) |
| N1 ⁱ —Cu1—N1—C8 | 54.2 (5) | C8—N1—C7—C6 | -177.4 (7) |
| I1—Cu1—N1—C8 | -116.5 (4) | Cu1—N1—C7—C6 | -4.5 (12) |
| C6—C1—C2—C3 | 0.8 (12) | C1—C6—C7—N1 | -0.3 (12) |
| C1—C2—C3—C4 | -2.2 (12) | C5—C6—C7—N1 | 179.8 (7) |
| C1—C2—C3—Br1 | 175.9 (6) | C7—N1—C8—C9 | 109.6 (8) |
| C2—C3—C4—C5 | 1.6 (13) | Cu1—N1—C8—C9 | -64.9 (7) |
| Br1—C3—C4—C5 | -176.6 (6) | N1—C8—C9—C10 | -57.8 (9) |
| C3—C4—C5—C6 | 0.5 (12) | N1—C8—C9—C8 ⁱ | 66.0 (10) |
| C2—C1—C6—C5 | 1.3 (12) | N1—C8—C9—C11 | -176.9 (7) |
| C2—C1—C6—C7 | -178.6 (7) | | |

Symmetry codes: (i) $x, -y+1, z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|-----------------------------------|-------|-------------|-------------|---------------|
| C8—H8A \cdots Cg1 ⁱⁱ | 0.99 | 2.83 | 3.631 (9) | 138 |

Symmetry codes: (ii) $-x+1/2, -y+1/2, -z$.

Fig. 1




Fig. 2

