

catena-Poly[*cis*-dipyrimidine-*trans*-dithiocyanatoiron(II)]-di- μ -pyrimidine-[*trans*-dithiocyanatoiron(II)]-di- μ -pyrimidine]

Mario Wriedt, Sina Sellmer, Inke Jess and Christian Näther*

Institut für Anorganische Chemie, Christian-Albrechts-Universität Kiel, Max-Eyth-Strasse 2, D-24118 Kiel, Germany
Correspondence e-mail: mwriedt@ac.uni-kiel.de


Received 11 February 2009; accepted 16 February 2009

Key indicators: single-crystal X-ray study; $T = 170\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.051; wR factor = 0.124; data-to-parameter ratio = 14.7.

In the crystal structure of the title compound, $[\text{Fe}_3(\text{NCS})_6(\text{C}_4\text{H}_4\text{N}_2)_8]_n$, each iron(II) cation is coordinated by four *N*-bonded pyrimidine ligands and two *N*-bonded thiocyanate anions in a distorted octahedral environment. The asymmetric unit consists of one iron cation located on a crystallographic center of inversion, as well as one iron cation, three thiocyanate anions and four pyrimidine ligands occupying general positions. The structure consists of square secondary building units (SBUs) with an Fe atom at each corner, which are $\mu\text{-}N^1\text{:}N^3$ -bridged by the pyrimidine ligands. The SBUs are linked into infinite chains running in the *c*-axis direction *via* common opposite corners.

Related literature

For related pyrimidine structures, see: Lloret *et al.* (1998); Näther *et al.* (2007); Näther & Jess (2004). For general background, see: Näther & Greve (2003); Näther, Wriedt & Jess (2003); Wriedt *et al.* (2008, 2009).



Experimental

Crystal data

$[\text{Fe}_3(\text{NCS})_6(\text{C}_4\text{H}_4\text{N}_2)_8]$	$V = 2423.0(2)\text{ \AA}^3$
$M_r = 1156.77$	$Z = 2$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 18.256(1)\text{ \AA}$	$\mu = 1.20\text{ mm}^{-1}$
$b = 16.2855(9)\text{ \AA}$	$T = 170\text{ K}$
$c = 8.2765(4)\text{ \AA}$	$0.12 \times 0.10 \times 0.07\text{ mm}$
$\beta = 100.042(7)^\circ$	

Data collection

Stoe IPDS-1 diffractometer	20463 measured reflections
Absorption correction: numerical (<i>X-SHAPE</i> and <i>X-RED32</i> ; Stoe, 2008)	4615 independent reflections
$S_{\min} = 0.859$, $T_{\max} = 0.912$	3795 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.064$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$	314 parameters
$wR(F^2) = 0.124$	H-atom parameters constrained
$S = 1.13$	$\Delta\rho_{\max} = 0.95\text{ e \AA}^{-3}$
4615 reflections	$\Delta\rho_{\min} = -0.41\text{ e \AA}^{-3}$

Table 1
Selected bond lengths (\AA).

Fe1–N41	2.075 (3)	Fe2–N61	2.075 (3)
Fe1–N51	2.089 (3)	Fe2–N61 ⁱ	2.075 (3)
Fe1–N1	2.252 (3)	Fe2–N2	2.252 (3)
Fe1–N31	2.254 (3)	Fe2–N2 ⁱ	2.252 (3)
Fe1–N21	2.262 (3)	Fe2–N32 ⁱⁱ	2.291 (3)
Fe1–N11	2.304 (3)	Fe2–N32 ⁱⁱⁱ	2.291 (3)

Symmetry codes: (i) $-x, -y + 1, -z + 2$; (ii) $-x, -y + 1, -z + 1$; (iii) $x, y, z + 1$.

Data collection: *X-AREA* (Stoe, 2008); cell refinement: *X-AREA*; data reduction: *X-AREA*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *XCIF* in *SHELXTL*.

MW thanks the Stiftung Stipendien-Fonds des Verbandes der Chemischen Industrie and the Studienstiftung des deutschen Volkes" for a PhD scholarship. Moreover, we gratefully acknowledge financial support by the State of Schleswig-Holstein and we thank Professor Dr. Wolfgang Bensch for the opportunity to use his experimental facility.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IM2102).

References

- Lloret, F., Munno, G. D., Julve, M., Cano, J., Ruiz, R. & Caneschi, A. (1998). *Angew. Chem. Int. Ed. Engl.* **37**, 135–138.
- Näther, C., Bhosekar, G. & Jess, I. (2007). *Eur. J. Inorg. Chem.* pp. 5353–5359.
- Näther, C. & Greve, J. (2003). *J. Solid State Chem.* **176**, 259–265.
- Näther, C. & Jess, I. (2004). *Eur. J. Inorg. Chem.* pp. 2868–2876.
- Näther, C., Wriedt, M. & Jess, I. (2003). *Inorg. Chem.* **42**, 2391–2397.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Stoe (2008). *X-AREA, X-RED32 and X-SHAPE*. Stoe & Cie, Darmstadt, Germany.
- Wriedt, M., Jess, I. & Näther, C. (2008). *Eur. J. Inorg. Chem.* pp. 363–372.
- Wriedt, M., Jess, I. & Näther, C. (2009). *Eur. J. Inorg. Chem.* In the press.

supporting information

Acta Cryst. (2009). E65, m296–m297 [doi:10.1107/S1600536809005509]

catena-Poly[bis[*cis*-dipyrimidine-*trans*-dithiocyanatoiron(II)]-di- μ -pyrimidine-[*trans*-dithiocyanatoiron(II)]-di- μ -pyrimidine]

Mario Wriedt, Sina Sellmer, Inke Jess and Christian Näther

S1. Comment

In our ongoing investigation on the syntheses, structures and properties of new coordination polymers based on paramagnetic metal pseudohalides and N-donor ligands, we have demonstrated that new ligand deficient coordination polymers with interesting magnetic interactions can be conveniently prepared by thermal decomposition of suitable ligand rich precursor compounds (Näther & Greve, 2003 and Wriedt *et al.*, 2009). In further investigations we have reacted iron(II) sulfate heptahydrate and potassium thiocyanate with pyrimidine leading to the formation of single crystals of the title compound.


The 3:8 title compound $[(\text{Fe}(\text{SCN})_2)_3(\mu\text{-pyrimidine})_8(\text{pyrimidine})_2]_n$ structurally (Fig. 1) represents a chain-like coordination polymer consisting of square SBU's with shared opposite corners as the characteristic motif. Each square SBU contains four iron(II) cations in its corners that are μ -1,3-(*N,N'*) bridged by pyrimidine ligands. The commonly used corner iron cations are located on crystallographic centers of inversion and are each octahedrally coordinated by four bridging pyrimidine ligands and two symmetry related terminal *N*-bonded thiocyanate anions. The iron cations in general positions are each octahedrally coordinated by two bridging pyrimidine ligands as well as two terminal *N*-bonded pyrimidine ligands and two terminal *N*-bonded thiocyanate anions (Fig 2). The Fe—NCS distances amount to 2.075 (3) and 2.089 (3) Å and the Fe—N_{pyrimidine} distances range from 2.252 (3) to 2.304 (3) Å. The angles around the iron cations range from 86.9 (1) to 180.0°. The shortest intra- and interchain Fe···Fe distances measure to 6.2508 (6) and 8.7620 (8) Å, respectively.

S2. Experimental


FeSO₄·7H₂O and pyrimidine were obtained from Alfa Aesar, KSCN was obtained from Fluka. 0.25 mmol (69.5 mg) FeSO₄ × 7 H₂O, 0.5 mmol (48.6 mg) KSCN and 9 mmol (720.8 mg) pyrimidine were transferred into a test-tube. In a solvent-free reaction at room temperature without stirring red block-shaped single crystals of the title compound were obtained after 2 weeks.

S3. Refinement

All H atoms were located from difference map but were positioned with idealized geometry and were refined isotropic with $U_{\text{eq}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ of the parent atom using a riding model with C—H = 0.95 Å.

**Figure 1**

Molecular structure of the repeating unit of the title compound showing the labelling scheme. Displacement ellipsoids drawn at the 50% probability level. Symmetry codes: i = $-x, -y + 1, -z + 2$; ii = $x, y, z + 1$; iii = $-x, -y + 1, -z + 1$.

**Figure 2**

Crystal structure of the title compound viewed along the b -axis.

catena-Poly[bis[cis-dipyrimidine-*trans*- dithiocyanatoiron(II)]-di- μ -pyrimidine-[*trans*-dithiocyanatoiron(II)]- di- μ -pyrimidine]*Crystal data* $[\text{Fe}_3(\text{NCS})_6(\text{C}_4\text{H}_4\text{N}_2)_8]$ $M_r = 1156.77$ Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

 $a = 18.256 (1) \text{ \AA}$ $b = 16.2855 (9) \text{ \AA}$ $c = 8.2765 (4) \text{ \AA}$ $\beta = 100.042 (7)^\circ$ $V = 2423.0 (2) \text{ \AA}^3$ $Z = 2$ $F(000) = 1176$ $D_x = 1.586 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 8000 reflections

 $\theta = 8.3\text{--}27.2^\circ$ $\mu = 1.20 \text{ mm}^{-1}$ $T = 170 \text{ K}$

Block, red

 $0.12 \times 0.10 \times 0.07 \text{ mm}$ *Data collection*

Stoe IPDS-1

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ scans

Absorption correction: numerical

(X-SHAPE and X-RED32; Stoe, 2008)

 $T_{\min} = 0.859$, $T_{\max} = 0.912$

20463 measured reflections

4615 independent reflections

3795 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.064$ $\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 2.6^\circ$ $h = -22 \rightarrow 22$ $k = -20 \rightarrow 20$ $l = -10 \rightarrow 10$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.051$ $wR(F^2) = 0.124$ $S = 1.13$

4615 reflections

314 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0419P)^2 + 8.0294P]$
where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.95 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.41 \text{ e \AA}^{-3}$ Extinction correction: SHELXL97 (Sheldrick,
2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0024 (5)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^* / U_{\text{eq}}$
Fe1	0.26416 (3)	0.48712 (3)	0.61108 (6)	0.01115 (16)

Fe2	0.0000	0.5000	1.0000	0.01002 (18)
N41	0.26059 (19)	0.5829 (2)	0.7749 (4)	0.0202 (7)
C41	0.2859 (2)	0.6276 (2)	0.8790 (5)	0.0182 (8)
S41	0.32547 (8)	0.68848 (8)	1.02316 (17)	0.0418 (4)
N51	0.26521 (18)	0.3912 (2)	0.4439 (4)	0.0184 (7)
C51	0.26229 (19)	0.3726 (2)	0.3063 (5)	0.0127 (7)
S51	0.25824 (6)	0.35068 (7)	0.11413 (12)	0.0241 (3)
N61	0.00127 (18)	0.5933 (2)	0.8299 (4)	0.0167 (7)
C61	0.0140 (2)	0.6208 (2)	0.7072 (5)	0.0146 (8)
S61	0.03129 (7)	0.65765 (7)	0.53635 (14)	0.0292 (3)
N1	0.17147 (16)	0.42202 (19)	0.7062 (4)	0.0131 (6)
N2	0.07898 (17)	0.42630 (19)	0.8767 (4)	0.0131 (6)
C1	0.12769 (19)	0.4614 (2)	0.7952 (4)	0.0130 (7)
H1	0.1317	0.5195	0.8008	0.016*
C2	0.0754 (2)	0.3440 (2)	0.8720 (5)	0.0165 (8)
H2	0.0432	0.3166	0.9330	0.020*
C3	0.1170 (2)	0.2978 (2)	0.7813 (5)	0.0179 (8)
H3	0.1133	0.2397	0.7768	0.021*
C4	0.1643 (2)	0.3399 (2)	0.6977 (5)	0.0164 (8)
H4	0.1927	0.3099	0.6320	0.020*
N11	0.36347 (17)	0.5483 (2)	0.5208 (4)	0.0157 (7)
N12	0.4695 (2)	0.6360 (2)	0.5800 (5)	0.0264 (8)
C11	0.4082 (2)	0.6011 (3)	0.6157 (5)	0.0212 (9)
H11	0.3949	0.6149	0.7183	0.025*
C12	0.4872 (2)	0.6144 (3)	0.4348 (6)	0.0275 (10)
H12	0.5309	0.6369	0.4048	0.033*
C13	0.4452 (3)	0.5618 (3)	0.3286 (6)	0.0289 (10)
H13	0.4584	0.5482	0.2258	0.035*
C14	0.3827 (2)	0.5293 (3)	0.3769 (5)	0.0231 (9)
H14	0.3525	0.4923	0.3056	0.028*
N21	0.34829 (17)	0.4171 (2)	0.7928 (4)	0.0150 (7)
N22	0.4465 (2)	0.4159 (2)	1.0257 (5)	0.0285 (9)
C21	0.3925 (2)	0.4525 (3)	0.9200 (5)	0.0216 (9)
H21	0.3847	0.5093	0.9370	0.026*
C22	0.4557 (2)	0.3357 (3)	1.0026 (5)	0.0249 (9)
H22	0.4936	0.3074	1.0746	0.030*
C23	0.4125 (2)	0.2927 (3)	0.8790 (5)	0.0219 (9)
H23	0.4191	0.2354	0.8658	0.026*
C24	0.3587 (2)	0.3362 (2)	0.7736 (5)	0.0150 (8)
H24	0.3284	0.3080	0.6859	0.018*
N31	0.19048 (17)	0.5567 (2)	0.4101 (4)	0.0155 (7)
N32	0.09789 (17)	0.56059 (19)	0.1687 (4)	0.0143 (7)
C31	0.1369 (2)	0.5230 (2)	0.2998 (5)	0.0151 (8)
H31	0.1253	0.4670	0.3159	0.018*
C32	0.1156 (2)	0.6395 (2)	0.1491 (5)	0.0184 (8)
H32	0.0906	0.6684	0.0558	0.022*
C33	0.1689 (2)	0.6805 (3)	0.2591 (6)	0.0244 (9)
H33	0.1799	0.7368	0.2448	0.029*

C34	0.2051 (2)	0.6362 (2)	0.3899 (5)	0.0203 (9)
H34	0.2417	0.6627	0.4684	0.024*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Fe1	0.0097 (3)	0.0137 (3)	0.0102 (3)	-0.00061 (19)	0.0022 (2)	-0.0013 (2)
Fe2	0.0084 (3)	0.0124 (4)	0.0098 (4)	0.0004 (3)	0.0029 (3)	0.0014 (3)
N41	0.0200 (17)	0.0208 (18)	0.0212 (18)	-0.0004 (14)	0.0073 (15)	-0.0038 (14)
C41	0.0174 (18)	0.0167 (19)	0.020 (2)	0.0057 (15)	0.0032 (16)	-0.0026 (16)
S41	0.0447 (7)	0.0309 (7)	0.0414 (7)	0.0109 (5)	-0.0158 (6)	-0.0206 (6)
N51	0.0190 (17)	0.0198 (17)	0.0169 (17)	0.0001 (13)	0.0047 (14)	-0.0014 (14)
C51	0.0103 (16)	0.0106 (17)	0.0170 (19)	0.0005 (13)	0.0015 (14)	-0.0018 (14)
S51	0.0343 (6)	0.0272 (5)	0.0112 (5)	-0.0026 (4)	0.0052 (4)	-0.0027 (4)
N61	0.0162 (16)	0.0181 (17)	0.0161 (17)	0.0015 (12)	0.0035 (13)	0.0033 (13)
C61	0.0127 (17)	0.0131 (17)	0.0178 (19)	0.0037 (14)	0.0022 (15)	-0.0002 (15)
S61	0.0441 (7)	0.0281 (6)	0.0189 (5)	0.0067 (5)	0.0147 (5)	0.0085 (4)
N1	0.0085 (14)	0.0158 (16)	0.0155 (15)	-0.0002 (11)	0.0034 (12)	-0.0004 (12)
N2	0.0114 (14)	0.0153 (15)	0.0132 (15)	0.0004 (12)	0.0039 (12)	-0.0001 (12)
C1	0.0110 (16)	0.0159 (18)	0.0126 (17)	0.0012 (14)	0.0035 (14)	0.0007 (14)
C2	0.0151 (18)	0.0142 (18)	0.022 (2)	-0.0018 (14)	0.0075 (16)	0.0016 (15)
C3	0.0166 (18)	0.0114 (17)	0.026 (2)	-0.0026 (14)	0.0042 (16)	-0.0024 (15)
C4	0.0124 (17)	0.0175 (19)	0.020 (2)	0.0009 (14)	0.0039 (15)	-0.0038 (15)
N11	0.0107 (14)	0.0210 (17)	0.0153 (16)	-0.0021 (12)	0.0015 (13)	0.0017 (13)
N12	0.0231 (18)	0.032 (2)	0.027 (2)	-0.0136 (15)	0.0106 (16)	-0.0069 (16)
C11	0.0184 (19)	0.026 (2)	0.021 (2)	-0.0072 (16)	0.0072 (17)	-0.0056 (16)
C12	0.024 (2)	0.038 (3)	0.024 (2)	-0.0139 (19)	0.0140 (18)	-0.0027 (19)
C13	0.028 (2)	0.040 (3)	0.021 (2)	-0.012 (2)	0.0124 (19)	-0.0024 (19)
C14	0.024 (2)	0.031 (2)	0.015 (2)	-0.0097 (17)	0.0052 (17)	-0.0062 (17)
N21	0.0131 (15)	0.0181 (16)	0.0139 (15)	-0.0007 (12)	0.0026 (13)	0.0022 (13)
N22	0.0267 (19)	0.026 (2)	0.0274 (19)	-0.0014 (15)	-0.0108 (16)	0.0036 (16)
C21	0.025 (2)	0.021 (2)	0.0167 (19)	0.0001 (16)	-0.0026 (17)	0.0009 (16)
C22	0.021 (2)	0.027 (2)	0.026 (2)	0.0033 (17)	-0.0006 (18)	0.0084 (18)
C23	0.022 (2)	0.022 (2)	0.021 (2)	0.0063 (16)	0.0013 (17)	0.0055 (16)
C24	0.0157 (18)	0.0183 (19)	0.0114 (18)	0.0018 (14)	0.0036 (15)	0.0022 (14)
N31	0.0115 (15)	0.0181 (16)	0.0163 (16)	-0.0014 (12)	0.0011 (13)	0.0007 (13)
N32	0.0116 (15)	0.0154 (16)	0.0156 (16)	-0.0005 (12)	0.0012 (13)	-0.0015 (12)
C31	0.0112 (17)	0.0169 (18)	0.0165 (18)	0.0004 (14)	0.0010 (15)	-0.0004 (15)
C32	0.0155 (18)	0.0185 (19)	0.020 (2)	0.0003 (15)	-0.0008 (16)	0.0061 (15)
C33	0.023 (2)	0.017 (2)	0.031 (2)	-0.0046 (16)	-0.0027 (18)	0.0032 (17)
C34	0.0185 (19)	0.020 (2)	0.020 (2)	-0.0045 (15)	-0.0032 (16)	-0.0030 (16)

Geometric parameters (\AA , ^\circ)

Fe1—N41	2.075 (3)	N12—C11	1.334 (5)
Fe1—N51	2.089 (3)	N12—C12	1.344 (6)
Fe1—N1	2.252 (3)	C11—H11	0.9500
Fe1—N31	2.254 (3)	C12—C13	1.364 (6)

Fe1—N21	2.262 (3)	C12—H12	0.9500
Fe1—N11	2.304 (3)	C13—C14	1.378 (6)
Fe2—N61	2.075 (3)	C13—H13	0.9500
Fe2—N61 ⁱ	2.075 (3)	C14—H14	0.9500
Fe2—N2	2.252 (3)	N21—C21	1.340 (5)
Fe2—N2 ⁱ	2.252 (3)	N21—C24	1.345 (5)
Fe2—N32 ⁱⁱ	2.291 (3)	N22—C22	1.334 (6)
Fe2—N32 ⁱⁱⁱ	2.291 (3)	N22—C21	1.339 (5)
N41—C41	1.161 (5)	C21—H21	0.9500
C41—S41	1.621 (4)	C22—C23	1.371 (6)
N51—C51	1.171 (5)	C22—H22	0.9500
C51—S51	1.619 (4)	C23—C24	1.388 (5)
N61—C61	1.169 (5)	C23—H23	0.9500
C61—S61	1.618 (4)	C24—H24	0.9500
N1—C1	1.341 (5)	N31—C31	1.335 (5)
N1—C4	1.344 (5)	N31—C34	1.338 (5)
N2—C1	1.335 (5)	N32—C31	1.338 (5)
N2—C2	1.343 (5)	N32—C32	1.343 (5)
C1—H1	0.9500	N32—Fe2 ^{iv}	2.291 (3)
C2—C3	1.380 (6)	C31—H31	0.9500
C2—H2	0.9500	C32—C33	1.382 (6)
C3—C4	1.379 (6)	C32—H32	0.9500
C3—H3	0.9500	C33—C34	1.372 (6)
C4—H4	0.9500	C33—H33	0.9500
N11—C14	1.336 (5)	C34—H34	0.9500
N11—C11	1.342 (5)		
N41—Fe1—N51	178.69 (13)	N1—C4—H4	118.9
N41—Fe1—N1	90.82 (13)	C3—C4—H4	118.9
N51—Fe1—N1	88.50 (12)	C14—N11—C11	116.2 (3)
N41—Fe1—N31	91.24 (13)	C14—N11—Fe1	122.3 (3)
N51—Fe1—N31	87.72 (12)	C11—N11—Fe1	121.4 (3)
N1—Fe1—N31	96.10 (11)	C11—N12—C12	115.4 (4)
N41—Fe1—N21	92.26 (13)	N12—C11—N11	126.3 (4)
N51—Fe1—N21	88.85 (12)	N12—C11—H11	116.8
N1—Fe1—N21	89.70 (11)	N11—C11—H11	116.8
N31—Fe1—N21	173.18 (12)	N12—C12—C13	123.0 (4)
N41—Fe1—N11	90.18 (13)	N12—C12—H12	118.5
N51—Fe1—N11	90.56 (13)	C13—C12—H12	118.5
N1—Fe1—N11	176.83 (11)	C12—C13—C14	117.1 (4)
N31—Fe1—N11	86.88 (11)	C12—C13—H13	121.5
N21—Fe1—N11	87.25 (11)	C14—C13—H13	121.5
N61—Fe2—N61 ⁱ	180.000 (1)	N11—C14—C13	122.0 (4)
N61—Fe2—N2	89.96 (12)	N11—C14—H14	119.0
N61 ⁱ —Fe2—N2	90.04 (12)	C13—C14—H14	119.0
N61—Fe2—N2 ⁱ	90.04 (12)	C21—N21—C24	115.9 (3)
N61 ⁱ —Fe2—N2 ⁱ	89.96 (12)	C21—N21—Fe1	123.5 (3)
N2—Fe2—N2 ⁱ	180.000 (1)	C24—N21—Fe1	120.6 (2)

N61—Fe2—N32 ⁱⁱ	90.10 (12)	C22—N22—C21	116.0 (4)
N61 ⁱ —Fe2—N32 ⁱⁱ	89.90 (12)	N22—C21—N21	126.6 (4)
N2—Fe2—N32 ⁱⁱ	89.27 (11)	N22—C21—H21	116.7
N2 ⁱ —Fe2—N32 ⁱⁱ	90.73 (11)	N21—C21—H21	116.7
N61—Fe2—N32 ⁱⁱⁱ	89.90 (12)	N22—C22—C23	122.4 (4)
N61 ⁱ —Fe2—N32 ⁱⁱⁱ	90.10 (12)	N22—C22—H22	118.8
N2—Fe2—N32 ⁱⁱⁱ	90.73 (11)	C23—C22—H22	118.8
N2 ⁱ —Fe2—N32 ⁱⁱⁱ	89.27 (11)	C22—C23—C24	117.4 (4)
N32 ⁱⁱ —Fe2—N32 ⁱⁱⁱ	180.000 (1)	C22—C23—H23	121.3
C41—N41—Fe1	154.5 (3)	C24—C23—H23	121.3
N41—C41—S41	177.0 (4)	N21—C24—C23	121.6 (4)
C51—N51—Fe1	146.5 (3)	N21—C24—H24	119.2
N51—C51—S51	177.7 (3)	C23—C24—H24	119.2
C61—N61—Fe2	153.7 (3)	C31—N31—C34	116.8 (3)
N61—C61—S61	179.3 (4)	C31—N31—Fe1	124.8 (3)
C1—N1—C4	116.2 (3)	C34—N31—Fe1	118.2 (2)
C1—N1—Fe1	121.5 (2)	C31—N32—C32	115.6 (3)
C4—N1—Fe1	121.5 (3)	C31—N32—Fe2 ^{iv}	123.0 (3)
C1—N2—C2	116.5 (3)	C32—N32—Fe2 ^{iv}	121.2 (2)
C1—N2—Fe2	122.4 (2)	N31—C31—N32	126.1 (4)
C2—N2—Fe2	120.9 (3)	N31—C31—H31	117.0
N2—C1—N1	125.9 (3)	N32—C31—H31	117.0
N2—C1—H1	117.1	N32—C32—C33	122.7 (4)
N1—C1—H1	117.1	N32—C32—H32	118.7
N2—C2—C3	122.1 (4)	C33—C32—H32	118.7
N2—C2—H2	119.0	C34—C33—C32	116.9 (4)
C3—C2—H2	119.0	C34—C33—H33	121.6
C4—C3—C2	117.0 (4)	C32—C33—H33	121.6
C4—C3—H3	121.5	N31—C34—C33	121.9 (4)
C2—C3—H3	121.5	N31—C34—H34	119.0
N1—C4—C3	122.2 (4)	C33—C34—H34	119.0

Symmetry codes: (i) $-x, -y+1, -z+2$; (ii) $-x, -y+1, -z+1$; (iii) $x, y, z+1$; (iv) $x, y, z-1$.